



Journal of Health Disparities Research and Practice
Volume 9, Special Edition 1, Summer 2016, pp. 28-29
© 2011 Center for Health Disparities Research
School of Community Health Sciences
University of Nevada, Las Vegas

Elucidating the Expression Profile of EZH2 Isoforms in Endometriosis: An immunohistochemical study

John L. Soto-Vargas

Idhaliz Flores, PhD; Ponce Health Sciences University/Ponce Research Institute

Mariano Colón-Caraballo, BS; Ponce Health Science University/Ponce Research Institute

Andrés Velázquez-García, BS; Ponce Health Science University Medical School

Coordinating Center: Charles R. Drew University of Medicine and Science

ABSTRACT

Endometriosis is an estrogen-dependent gynecological disease that affects 1 out of 10 women of reproductive age causing severe pelvic pain and infertility. Factors including genetics, environment, inflammation, and recently epigenetics have been shown to play roles in the pathophysiology of this disease. Histone methylation is an epigenetic modification that modulates gene expression by causing changes in the chromatin structure. Trimethylation of histone 3 at lysine residue 27 (H3K27me₃) is a histone mark related to gene repression. EZH2 is the histone methyltransferase (HMT) responsible of catalyzing H3K27me₃. It has been shown that the EZH2 is involved in carcinogenesis; however, the specific role of EZH2 in endometriosis is unknown. This is important because there are drugs available that block this enzyme's functioning, and could serve as a potential new treatment.

We have previously shown that endometriotic lesions are characterized by high H3K27me₃ nuclear immunostaining. Therefore, we hypothesize that EZH2 will be highly expressed in lesions compared to endometrium of patients and controls. Using immunohistochemistry (IHC) of an endometriosis Tissue Micro Array (TMA), EZH2 α and β nuclear intensity were assessed using specific antibodies. We observed that pelvic endometriotic lesions (peritoneal and fallopian tube) have higher EZH2 α intensity scores compared to control tissues. EZH2 β nuclear immunostaining analysis is ongoing. We expect to observe a higher nuclear intensity score in endometriotic lesions compared to endometriosis free endometria.

This study is the first to analyze the expression profile of EZH2 isoforms in endometriosis. These studies will help better understand the role of EZH2 in this disease.

Keywords: EZH2, H3K27me₃, endometriosis, epigenetic

29 Elucidating the Expression Profile of EZH2 Isoforms in Endometriosis
Soto-Vargas et al.

ACKNOWLEDGEMENTS

The STEP-UP HS program is supported by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health, Grant Number: 5R25DK078384-09.