

11-1-2011

Evaluating the impact of social determinants on pedestrian injury in Clark County, Nevada

Jonathon LaValley
University of Nevada, Las Vegas

Courtney Ann Coughenour
University of Nevada, Las Vegas

Michelle Chino
University of Nevada, Las Vegas, michelle.chino@unlv.edu

Follow this and additional works at: https://digitalscholarship.unlv.edu/hs_grad_pubs_pres



Part of the [Community-Based Research Commons](#), [Medicine and Health Commons](#), and the [Public Health Commons](#)

Repository Citation

LaValley, J., Coughenour, C. A., Chino, M. (2011). Evaluating the impact of social determinants on pedestrian injury in Clark County, Nevada.

https://digitalscholarship.unlv.edu/hs_grad_pubs_pres/1

This Poster is protected by copyright and/or related rights. It has been brought to you by Digital Scholarship@UNLV with permission from the rights-holder(s). You are free to use this Poster in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/or on the work itself.

This Poster has been accepted for inclusion in Graduate Publications & Presentations (HS) by an authorized administrator of Digital Scholarship@UNLV. For more information, please contact digitalscholarship@unlv.edu.

Evaluating the Impact of Social Determinants on Pedestrian Injury in Clark County, Nevada

Jonathon LaValley, MPH, Courtney Coughenour, MPH, Michelle Chino, PhD

School of Community Health Sciences, University of Nevada Las Vegas, 4505 S. Maryland Pkwy, Las Vegas, NV 89154



BACKGROUND

- Pedestrian crashes are a major public health concern. In 2009 there were over 4,000 pedestrian deaths and 59,000 pedestrian injuries¹.
- In 2008, NV had the 5th highest pedestrian fatality rate in the nation².
- The Las Vegas metropolitan area accounts for 82% of the fatalities³ and 75% of pedestrian injuries that take place in NV⁴.
- Las Vegas is an urban, sprawling, metropolis which increased in population size by 83% between 1990 and 2000⁵ and 41.8% from 2000 to 2010⁶.
- Pedestrian fatalities have been positively correlated with urban sprawl⁷, lack of pedestrian facilities, liquor outlet density, and rapid population growth⁸.
- To ensure sustainable communities, pedestrian injury issues in Clark County, NV warrant immediate attention.

OBJECTIVES

In order to understand the social determinants which may impact pedestrian injury rates in an urban, sprawling, western community we conducted an ecologic investigation to compare pedestrian crash rates by social determinants available for census tracts in Clark County, NV.

METHODS

Data Sources: Electronic Uniform Police Traffic Crash Reports, 2004 – 2008, Census 2000 Data

Data elements included: neighborhood demographics, socioeconomic factors, and characteristics of the built environment

Pedestrian crash and fatality rates per 1,000 population were calculated for each census tract and merged by tract onto Census 2000 data. Stepwise multiple linear regressions (MLR) were applied to explore the relationship between each of the predictor variables, and pedestrian crash and fatality rates using SAS 9.3. Prior to MLR, tests for linearity of the criterion variable, normality, and multicollinearity were applied.



Table 1. Mean Pedestrian Crash Rates by Census Tract

Variable	Mean	SD	Min	Max
Total Crash Rate per 1,000 pop	1.95	2.40	0.13	16.78
Fatal Rate per 1,000 pop	0.39	0.30	0.04	7.31

Table 2. Influence of neighborhood demographics, socioeconomic factors, and built environment characteristics on pedestrian crash rates.

Variable	β	P-value
% households without vehicle	9.18	p<0.001
liquor outlet density	1.85	p<0.001
number transit stops per square meter	0.83	p<0.001
% population walk or take public transit	0.16	p<0.001
population density	- 4.6	p<0.001

RESULTS

2,262 pedestrian crashes in 287 census tracts were identified. Of these, 92.4% were injury crashes, and 4.5% resulted in fatality, 3.1% involved property damage only.

Variables which related to poverty and income, home ownership, education, employment, age, and race were removed from the final model.

The final, selected model for pedestrian crash rate was significant and described 58.2% of the variance in the per capita pedestrian crash rate ($r^2=0.582$, $p<0.001$) with VIF all less than two.

The final, selected model for pedestrian fatality rate was significant but described only 8.5% of the variance in the per capita pedestrian fatality rate ($r^2=0.085$, $p<0.001$). Given the low amount of variance described by this model it was discarded.

CONCLUSION

The final model suggests that percentage of households without a vehicle, population density, liquor outlets per square meter, the number of transit stops per square meter, and the percent of the population who walk or use public transit in combination were useful in predicting crash rate. Contrary to previous research, population density was a protective factor for pedestrian crashes.

These results underscore the importance of neighborhood determinants in contributing to pedestrian injury. Results should have implications on injury prevention efforts, regional transit planning, and guide public policy. Public health professionals, city planners, and regional transit authorities must collaborate to successfully enhance pedestrian safety.

LIMITATIONS

The assumption that all individuals living in a Census tract have similar characteristics represents an ecological fallacy.

Three Census tracts were removed from the data set which had total pedestrian crashes greater than the tract population. These represent the Las Vegas 'strip', McCarran Airport, and Mt. Charleston, all popular tourist destinations.

Census data (2000) may not reflect current total population, environmental conditions, or population characteristics.

References

1. National Highway Traffic Safety Administration [NHTSA]. (2010). Traffic Safety Facts, 2009 data. Retrieved September 27, 2011 from <http://www-nrd.nhtsa.dot.gov/Pubs/811163.PDF>.
2. National Highway Traffic Safety Administration [NHTSA]. (2009). Traffic Safety Facts, 2008 data. Retrieved April 1, 2010 from <http://www-nrd.nhtsa.dot.gov/Pubs/811163.PDF>.
3. National Highway Traffic Safety Administration. Fatality analysis reporting system (FARS) web-based encyclopedia. Available at: www.fars.nhtsa.dot.gov. Accessed February 2011.
4. Pulugurtha, S. & Nambisan, S. (2002). Pedestrian safety engineering and intelligent transportation system-based countermeasures program for reducing pedestrian fatalities, injuries, conflicts, and other surrogate measures. Phase 1 Final Technical Report.
5. U.S. Census Bureau. (2001). Census 2000 data. Available at: www.census.gov. Accessed January 2011.
6. U.S. Census Bureau. (2010). Census 2010 data. Available at: www.census.gov. Accessed September 27, 2011.
7. Ewing, R., Schieber, R. & Zegeer, C. (2003). Urban sprawl as a risk factor in motor vehicle occupant and pedestrian fatalities. *Research and Practice*, 93(9).
8. Paulozzi, J. (2006). Is it safe to walk in the sunbelt? Geographic variation among pedestrian fatalities in the United States, 1999-2003. *Journal of Safety Research*, 37(5), 453-459.
9. Transportation for America. (2011). Dangerous by Design. Available at <http://t4america.org/docs/dbd2011/Dangerous-by-Design-2011.pdf>