

6-1-2022

Diabetes-Related Outcomes following Use of Basal-Bolus Insulin Vs Premixed Insulin In a VA Population With Type 2 Diabetes: A Retrospective Chart Review.

Ravisha Bai
University of Nevada, Las Vegas

Kavita Batra
University of Nevada, Las Vegas, kavita.batra@unlv.edu

Charmaine Yap
VA Southern Nevada Health Care System

Kenneth Izuora
University of Nevada, Las Vegas, kenneth.izuora@unlv.edu

Follow this and additional works at: https://digitalscholarship.unlv.edu/som_fac_presentations



Part of the [Endocrine System Diseases Commons](#), [Endocrinology Commons](#), and the [Endocrinology, Diabetes, and Metabolism Commons](#)

Repository Citation

Bai, R., Batra, K., Yap, C., Izuora, K. (2022). Diabetes-Related Outcomes following Use of Basal-Bolus Insulin Vs Premixed Insulin In a VA Population With Type 2 Diabetes: A Retrospective Chart Review.. 1-1. Available at: https://digitalscholarship.unlv.edu/som_fac_presentations/1

This Poster is protected by copyright and/or related rights. It has been brought to you by Digital Scholarship@UNLV with permission from the rights-holder(s). You are free to use this Poster in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/or on the work itself.

This Poster has been accepted for inclusion in School of Medicine Faculty Presentations by an authorized administrator of Digital Scholarship@UNLV. For more information, please contact digitalscholarship@unlv.edu.

DIABETES-RELATED OUTCOMES FOLLOWING USE OF BASAL-BOLUS INSULIN VS PREMIXED INSULIN IN A VA POPULATION WITH TYPE 2 DIABETES: A Retrospective Chart Review.



Authors: Ravisha Bai¹, Kavita Batra¹, Charmaine Yap², Kenneth Izuora¹
 Affiliations: 1 University of Nevada Las Vegas, 2 VA Southern Nevada Health Care System
 Conflict of interest disclosures: None



Background

- The complexity of basal bolus insulin (BB) therapy may adversely influence patient compliance.
- Premixed insulin (PM) is simpler and cheaper but is perceived to be less effective in managing diabetes (DM).
- We compared DM-related outcomes between BB and PM insulin regimens among a veteran population with type 2 DM.

Methods

- This is a retrospective chart review that enrolled 140 veterans (70 in each group) with type2 DM.
- Inclusion criteria: Age > 18 years and A1C ≥ 8%.
- Exclusion criteria: Type 1 DM and switching insulin regimen during the follow-up period.
- The included subjects were followed for a year, after starting 70/30 insulin NPH/regular (PM) OR detemir or glargine + aspart, lispro or regular insulin (BB).
- Our primary outcome was change in HbA1c one year of insulin therapy.
- Change in body mass index (BMI) and the rate of hypoglycemia were secondary outcomes
- Data were analyzed using Chi-square/Fisher exact tests and logistic regression.

| Variable | Characteristics | Basal Bolus Insulin Group n = 70 | Premixed Insulin Group n = 70 | P value |
|-----------------------------------|------------------------|-------------------------------------|----------------------------------|---------|
| Age (years) | - | 64.5±10.8 | 67.0±9.1 | 0.07 |
| Gender | Female | 3 (4.3) | 5 (7.1) | 0.5 |
| | Male | 67 (95.7) | 65 (92.9) | |
| Baseline BMI (kg/m ²) | - | 33.0±5.8 | 32.5±6.1 | 0.3 |
| Follow-up BMI | - | 33.8±5.7 | 33.0±6.3 | 0.2 |
| BMI difference | - | 0.84±3.1 | 0.40±2.2 | 0.2 |
| Baseline BMI Status | Normal (18.5-24.9) | 7 (10.0) | 7 (10.0) | 0.8 |
| | Overweight (25.0-29.9) | 14 (20.0) | 17 (24.3) | |
| | Obese (≥30.0) | 49 (70.0) | 46 (65.7) | |
| Follow-up BMI Status | Normal (18.5-24.9) | 4 (5.7) | 5 (7.1) | 0.4 |
| | Overweight (25.0-29.9) | 15 (21.4) | 21 (30.0) | |
| | Obese (≥30.0) | 51 (72.9) | 44 (62.9) | |
| Change in HbA1c | - | -1.9±1.8 | -2.1±1.9 | 0.3 |
| Hypoglycemia | Yes | 21 (30.0) | 15 (21.4) | 0.3 |
| | No | 49 (70.0) | 55 (78.6) | |

Following treatment initiation, there was no statistically significant difference between the BB and PM insulin groups for change in A1c (-1.9±1.8 vs -2.1±1.9%, p = 0.3).

Conclusion

- There were no significant differences in A1c change, rate of hypoglycemia, and change in body mass index in patients treated with either BB or PM insulin among our study population.
- These results suggest that PM insulin is equally effective and safe as BB insulin therapy.
- The use of PM insulin can potentially improve patient compliance and reduce the cost of care.