7-2001

A Cytochrome-b Perspective on Passerina Bunting Relationships

John Klicka

University of Nevada, Las Vegas, klicka@unlv.nevada.edu

Adam J. Fry

University of Minnesota

Robert M. Zink

University of Minnesota, rzink2@unl.edu

Christopher W. Thompson

University of Washington - Seattle

Follow this and additional works at: https://digitalscholarship.unlv.edu/hrc_ornithology

Part of the [Molecular Genetics Commons](#), [Ornithology Commons](#), and the [Population Biology Commons](#)

Repository Citation

Klicka, J., Fry, A. J., Zink, R. M., Thompson, C. W. (2001). A Cytochrome-b Perspective on Passerina Bunting Relationships. *The Auk*, 118(3), 610-623.

Available at: https://digitalscholarship.unlv.edu/hrc_ornithology/31

This Article is protected by copyright and/or related rights. It has been brought to you by Digital Scholarship@UNLV with permission from the rights-holder(s). You are free to use this Article in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/or on the work itself.

This Article has been accepted for inclusion in Ornithology Program (HRC) by an authorized administrator of Digital Scholarship@UNLV. For more information, please contact digitalscholarship@unlv.edu.

COLOR PLATE. The genus *Passerina*. The branching pattern reflects hypothesized relationships based on results of this study.

A CYTOCHROME-*b* PERSPECTIVE ON *PASSERINA* BUNTING RELATIONSHIPS

JOHN KLIKA,^{1,3} ADAM J. FRY,^{1,4} ROBERT M. ZINK,¹ AND CHRISTOPHER W. THOMPSON^{2,5}

¹J. F. Bell Museum of Natural History, University of Minnesota, 1987 Upper Buford Circle, St. Paul, Minnesota 55108, USA; and

²Burke Museum and Department of Zoology DB-10, University of Washington, Seattle, Washington 98195, USA

ABSTRACT.—We sequenced the complete mitochondrial cytochrome-*b* gene (1,143 nucleotides) for representatives of each species in the cardinalid genera *Passerina* (6 species), *Guiraca* (1 species), and *Cyanocopsa* (3 species), and used a variety of phylogenetic methods to address relationships within and among genera. We determined that *Passerina*, as presently recognized, is paraphyletic. Lazuli Bunting (*P. amoena*) is sister to the much larger Blue Grosbeak (*Guiraca caerulea*). Indigo Bunting (*P. cyanea*) and Lazuli Bunting are not sister taxa as generally thought. In all weighted parsimony trees and for the gamma-corrected HKY tree, Indigo Bunting is the sister of two sister groups, a “blue” (Lazuli Bunting and Blue Grosbeak) and a “painted” (Rosita’s Bunting [*P. rositae*], Orange-breasted Bunting [*P. leclancherii*], Varied Bunting [*P. versicolor*], and Painted Bunting [*P. ciris*]) clade. The latter two species form a highly supported sister pair of relatively more recent origin. Uncorrected (*p*) distances for ingroup (*Passerina* and *Guiraca*) taxa range from 3.0% (*P. versicolor*–*P. ciris*) to 7.6% (*P. cyanea*–*P. leclancherii*) and average 6.5% overall. Assuming a molecular clock, a bunting “radiation” between 4.1 and 7.3 Mya yielded four lineages. This timing is consistent with fossil evidence and coincides with a late-Miocene cooling during which a variety of western grassland habitats evolved. A reduction in size at that time may have allowed buntings to exploit that new food resource (grass seeds). We speculate that the Blue Grosbeak subsequently gained large size and widespread distribution as a result of ecological character displacement. Received 25 October 1999, accepted 16 September 2000.

THE BUNTINGS of the avian genus *Passerina* (family Cardinalidae) include six species whose composite range includes Mexico, the United States, and parts of southern Canada. They are small, heavy-billed finches, most of which (see Thompson and Leu 1995) exhibit pronounced sexual dichromatism with brightly colored males and mostly brownish or olive females. Traditional museum studies relying upon comparisons of study skins (e.g. Ridgway 1901), as well as a more recent study employing numerical phenetic analyses of both skins and skeletons (Hellack and Schnell 1977), have concluded that the members of *Passerina*, as pres-

ently recognized (Sibley and Monroe 1990, AOU 1998), form a natural group. Nevertheless, some authors (Phillips et al. 1964, Blake 1969, Mayr and Short 1970) place the monotypic Blue Grosbeak (*Guiraca caerulea*) within this genus, and others (Paynter 1970) further expand *Passerina* to include additional genera. Thus, despite the phenotypic similarities of the six recognized species, monophyly of *Passerina* remains unresolved.

Relationships within *Passerina* also remain unclear. Phillips et al. (1964) consider the Indigo Bunting (*P. cyanea*) and the Lazuli Bunting (*P. amoena*) conspecific. Hybridization on the Great Plains between *P. cyanea* and *P. amoena* has been well documented (Sibley and Short 1959, Emlen et al. 1975, Kroodsma et al. 1975). Similarly, the Painted Bunting (*P. ciris*) and Varied Bunting (*P. versicolor*) are known to have hybridized (Storer 1961). Whereas such hybridization might be taken as evidence of sister-species relationships, a known pairing between *P. cyanea* and *P. ciris* (Taylor 1974) muddles the picture. The ability to hybridize might be a mis-

³ Present address: Barrick Museum, Box 454012, University of Nevada Las Vegas, 4505 Maryland Parkway, Las Vegas, Nevada 89154-4012, USA. E-mail: klicka@nevada.edu

⁴ Present address: Department of Ecology and Evolutionary Biology, Box G-W, Brown University, Providence, Rhode Island 02912, USA.

⁵ Present address: Washington Department of Fish and Wildlife, 16018 Mill Creek Boulevard, Mill Creek, Washington 98012, USA.

leading indicator of relationship (Zink and McKittrick 1995), therefore the construction of an independent phylogeny is warranted.

Our understanding of the species in *Passerina* is uneven. *Passerina cyanea* is among our best known songbirds with song, display behaviors, mating strategies, molt, and migration having been studied extensively (see Payne's [1992] comprehensive review for references). At the other extreme are the endemic forms of southern Mexico, Leclancher's (Orange-breasted) Bunting (*P. leclancherii*) and Rosita's (Rose-bellied) Bunting (*P. rositae*), about which very little is known. A few comparative studies have involved most or all members of *Passerina*, examining molts and plumages (summarized in Thompson and Leu 1994), songs (Thompson 1968), and morphology (Hellack and Schnell 1977). However, interpretation of those studies is compromised by the lack of an explicit phylogenetic hypothesis (Brooks and McLennan 1991).

In this study, we used DNA sequences from the complete mitochondrial (mtDNA) protein coding cytochrome-*b* (cyt-*b*) gene to assess the monophyly of *Passerina* as presently recognized and to construct a phylogenetic hypothesis for the constituents of this group. Cyt-*b* and its flanking regions are well characterized in birds and its rate of evolution is particularly well suited for avian studies at the species level (Moore and DeFilippis 1997). The resulting phylogenetic tree provides a historical framework for a better understanding of the evolution of various life-history traits and for investigating historical biogeography.

METHODS

Taxon sampling.—Outgroup choice is a critical component of molecular phylogenetic analysis (Smith 1994). Outgroups that are too distantly related to the group of interest lead to spurious rooting of the ingroup topology (Wheeler 1990). Relationships among Cardinalid genera are poorly understood, therefore an initial generic level phylogenetic survey of the group was performed. A cyt-*b* (430 bp) segment from at least one representative of each putative Cardinalid genus (excepting *Cyanoloxia*) was sequenced, as were representatives of several Thrupid genera, which were used as outgroups. The detailed results of that initial study are to be published elsewhere; however, in all phylogenetic analyses the *Passerina* buntings and *G. caerulea* were determined to constitute a clade (ingroup), sister to *Cyanocompsa*

(outgroup). Those three genera were thus targeted for complete cyt-*b* gene sequencing. Because adding more taxa to the sister group increases the chance of obtaining the correct tree (Smith 1994), single representatives of all three members of *Cyanocompsa* were subsequently used. To obtain a measure of intraspecific genetic variation and to provide a check for potential sequencing errors, each of six species of *Passerina* and the monotypic *G. caerulea* (ingroup taxa) were represented by two different tissue specimens (Table 1).

Laboratory protocols.—Total DNA was extracted from fragments (~100 mg) of muscle or liver tissues following either a standard phenol/chloroform protocol (Hillis et al. 1996) or via incubation in Chelex/Proteinase K, a modification of Ellegren's (1992) method. Overlapping sequence fragments were amplified via the polymerase chain reaction (PCR) using various combinations of the following published primers: L14851 (Kornegay et al. 1993), L14841 (Kocher et al. 1989), H15299 (Hackett 1996), H4A (Harshman 1996), LCBA, LCBB, LCBC, HCBC (Klicka et al. 1999); and two designed (J. Klicka) for general use on New World nine-primaried oscines: HCBRB (GGAGAATGACTCCGAYGTTTCA), and HCBJT (GGCTGGGGTGAAATTTTCTGGGTCT). A map of primer locations is available by request. PCR reactions were performed in 50 μ L total volumes using a 1.5 mM concentration of MgCl₂, 1 μ M concentrations of each primer, 0.2 mM final concentration of dNTPs, 0.5 μ L of *Tfl* polymerase (*Thermus flavus*, Epicentre), 2.2 μ L of 20 $\times$ *Tfl* buffer (400 mM [NH₄]₂SO₄ and 1 M Tris-HCl [pH 9.0], Epicentre) and 10–1,000 ng (2 μ L) of DNA template. A typical double-stranded amplification began with a 2 min denaturation at 94°C followed by 38 cycles of denaturing (94°C, 45 s), annealing (54°C, 1 min, 30 s), and primer extension (72°C, 1 min, 30 s) with a final extension of 72°C for 10 min. The presence of the desired product was verified by electrophoresis of 5 μ L in a 1% agarose gel (Seakem LE, FMC) followed by Ethidium Bromide staining. Excess primers and dNTPs were then removed by enzymatic treatment of the PCR product with Exonuclease 1 and Shrimp Alkaline Phosphatase (U.S. Biochemical Corp.). To ensure sequencing accuracy, large fragment overlaps were used and wherever possible both light and heavy strands of the treated product were sequenced using either the original or nested primers. Sequencing reactions (Hillis et al. 1996) were done manually (product no. 70130, U.S.B.), run out on 6% acrylamide (Gene-Page, Amresco) gels, and visualized by autoradiography (³⁵S). Sequences (GenBank nos. AF301446–AF301462) were aligned visually using published chicken (*Gallus*; Desjardins and Morais 1990) and Snow Goose (*Chen [Anser] caerulescens*; Quinn and Wilson 1993) sequences for comparison.

Data analysis.—Parsimony (uniform and differential weighting schemes) and maximum-likelihood

TABLE 1. Species names, voucher numbers, and localities of origin of taxa sequenced, this study. UWBM = University of Washington Burke Museum; BMNH = J. F. Bell Museum of Natural History; LSUMNS = Louisiana State University Museum of Natural Science.

Species	Museum	Number	Origin	Date
<i>Passerina cyanea</i> 1	BMNH	X7250	USA, Wisconsin, St. Croix Co.	June 1994
<i>Passerina cyanea</i> 2	BMNH	JK94162	USA, Minnesota, Isanti Co.	5 August 1992
<i>Passerina amoena</i> 1	BMNH	JK95030	USA, Washington, Sherman Co., mouth of Deschutes River	29 June 1995
<i>Passerina amoena</i> 2	LSUMNS	B-4015	USA, Washington, Douglas Co., 5 mi N of Palisades	17 August 1987
<i>Passerina leclancherii</i> 1	UWBM	CWT034	Mexico, Chiapas	3 May 1993
<i>Passerina leclancherii</i> 2	UWBM	CWT040	Mexico, Chiapas	5 May 1993
<i>Passerina rostrata</i> 1	UWBM	CWT033	Mexico, Chiapas, Monte Bonito	2 May 1993
<i>Passerina rostrata</i> 2	UWBM	CWT036	Mexico, Chiapas	4 May 1993
<i>Passerina ciris</i> 1	LSUMNS	B-5693	USA, Louisiana, Cameron Parish, Monkey Island, near Cameron	29 April 1984
<i>Passerina ciris</i> 2	LSUMNS	B-5694	USA, Louisiana, Cameron Parish, Monkey Island, near Cameron	29 April 1984
<i>Passerina versicolor</i> 1	UWBM	CWT095	Mexico, Coahuila, 35 km N of Saltillo	24 May 1993
<i>Passerina versicolor</i> 2	UWBM	CWT097	Mexico, Coahuila, 35 km N of Saltillo	24 May 1993
<i>Guitraca caerulea</i> 1	LSUMNS	B-21812	USA, Texas, Jeff Davis Co., Davis Mountains State Park	7 September 1992
<i>Guitraca caerulea</i> 2	LSUMNS	B-20958	USA, Louisiana, Cameron Parish, Garner Ridge	12 November 1992
<i>Cyanocompsa brissonii</i>	LSUMNS	B-18658	Bolivia, Dept. Santa Cruz, Cordellara	8 September 1990
<i>Cyanocompsa cyanoides</i>	LSUMNS	B-12708	Bolivia, Dept. Santa Cruz, Velasco	27 July 1988
<i>Cyanocompsa parellina</i>	UWBM	CWT082	Mexico, Oaxaca, Chivela, 6 mi NW	21 May 1993

methods were used to estimate phylogenetic trees. For comparison, and to allow quick visual assessment of relative genetic distances, we also constructed neighbor-joining trees. All analyses were done using PAUP* (version 4.0b2; Swofford 1999). To assess saturation, absolute numbers of transitions and transversions at first, second, and third codon positions were plotted against Kimura 2-parameter (K2-P, Kimura 1980) distances for all pairwise comparisons. Deviations from linearity (see below) indicated saturation effects, suggesting that the weighting of the more slowly evolving and presumably less homoplastic transversions was warranted.

We used an array of maximum parsimony (MP) weighting schemes to explore the dynamics of *cyt-b* evolution (Voelker and Edwards 1998). All analyses used branch-and-bound or full heuristic searches with 10 random addition sequence replicates. Initial parsimony analyses including 1:1 (equal weights), 2:1, and 5:1 weighting of transversions over transitions resulted in two nearly identical topologies whose unweighted tree lengths differed by a single step. Composite and codon position-specific sequence statistics were generated in PAUP* using the weighted topology (but equal weights). Relative rates of change among codon positions, percentage nucleotide composition, and transition:transversion ratios were calculated in MacClade (Maddison and Maddison 1992) by independently reconstructing, for various sequence partitions (see below), the average number of each transformation type onto the same tree. Subsequently, an analysis was performed in which weights were assigned on the basis of the ratios calculated for each codon partition; thus, first, second, and third position sites were weighted 8.6, 1.5, and 4.9, respectively. The "native" transition to transversion ratio (Sturmbauer and Meyer 1992) was also approximated by plotting transitions versus transversions for third position sites (plot not shown, approximate ratio 9.5:1). Following Yoder et al. (1996), we also used a 5:1 weighting of third position transversions only.

Six-parameter weighting schemes account for an additional source of error, base composition biases. The logic is identical to that behind transversion weighting; frequent classes of character change are considered more likely than rare classes to have experienced homoplasy (Cunningham 1997). Each of six possible transformations ($A \leftrightarrow G$, $C \leftrightarrow T$, $A \leftrightarrow C$, $A \leftrightarrow T$, $C \leftrightarrow G$, $T \leftrightarrow G$) was assigned a weight on the basis of the observed frequency in the data (Williams and Fitch 1989). These weights were derived by first mapping changes (average of all possible reconstructions used) onto the equal-weight (1:1; [nearly identical results were obtained using the weighted tree]) parsimony tree using MacClade (Maddison and Maddison 1992). The frequencies of each of the six classes of change were then calculated and negative-

TABLE 2. Cytochrome-*b* substitution type proportions shown above the diagonal; corresponding six-parameter weights shown below. Values were obtained by reconstructing the data onto the shortest equally weighted tree. Corrections for triangle inequality shown in parentheses.

	A	C	G	T
A	—	11.6%	23.5%	4.2%
C	4.1	—	1.0%	59.3%
G	2.8	8.8 (6.9)	—	0.4%
T	6.0 (5.1)	1.0	10.6 (7.9)	—

natural-log (–ln) transformed to obtain the weights entered into a step matrix (Table 2).

Mutation rates are known to vary throughout the *cyt-b* gene, presumably due to functional constraints at the amino acid level (e.g. Howell 1989, Krajewski and King 1996, Griffiths 1997). We addressed that problem by using MP and a differential weighting scheme in which transitions located in putatively more variable regions were down-weighted. Three rate classes of residues were defined using Howell’s (1989) structural model for mouse *cyt-b* in which individual residues were classified as *evolutionarily conserved*, *intermediate*, or *variable*. Although similar, these partitions do not correspond exactly to those used in other published studies (e.g. Griffiths 1997). We performed an analysis in which the sequence constituting each of these three partitions was weighted according to the inherent transition:transversion ratio (calculated in MacClade as described above) in that data partition. The result was transversions weighted 3.0:1, 6.2:1, and 4.8:1 over transitions for conserved, intermediate, and variable sequence partitions respectively. Standard tree statistics, including length, consistency index (CI), retention index (RI), and rescaled consistency index (RC) were generated for all parsimony reconstructions.

All maximum-likelihood (ML) analyses used empirical base frequencies, with full heuristic searches. An initial analysis was based on Felsenstein’s (1981, [F81]) model of DNA evolution which corrects for unequal base frequencies. The data were subsequently analyzed with the HKY85 (Hasegawa et al. 1985) model which additionally accommodates transition biases. That model in particular has proven to be effective in describing the dynamics of many genes (Goldman 1993, Yang 1996). This analysis used a transversion transition ratio of 6.9 and a gamma distribution with a shape parameter (α) of 0.106. These parameters were estimated using the stable (1:1, 2:1) MP trees following the recommendation of Swofford et al. (1996). Because the models used are nested, we were able to assess statistically their relative values by comparing log-likelihood values using likelihood ratio tests (LRT, Huelsenbeck and Rannala 1997).

Node support for MP analyses was determined via bootstrapping (Felsenstein 1985) with 500 replications using full heuristic searches and random addition of taxa. One hundred such replicates were done for ML analyses. Decay indices (Bremer 1994) were generated for the most highly supported parsimony-derived topology using TREEROT (Sorenson 1996). MacClade (Maddison and Maddison 1992) was used to compare tree lengths for competing tree topologies. Alternative topologies were tested using the Kishino-Hasegawa (1989) maximum-likelihood ratio test.

Biogeographic interpretations are enhanced by invoking a molecular clock. The validity of assuming clock-like rates of evolution for those taxa was tested via a likelihood-ratio test (Huelsenbeck and Rannala 1997) by comparing log-likelihood values from ML trees constructed with and without a molecular clock constraint using the HKY85/gamma(Γ)-corrected model of sequence evolution. To allow direct comparison with the Fleischer et al. (1998) songbird-specific clock calibration, we generated a matrix of K2-P Γ -corrected pairwise distances following their protocol. The α value used (0.300) was estimated via the method of Sullivan et al. (1995) in PAUP* from the shortest unweighted MP tree.

RESULTS

Sequence variation.—All 1,143 nucleotides were aligned with no insertions, deletions, or nonsense codons in evidence. Although the amplification of nuclear pseudogenes of mitochondrial origin can confound phylogenetic analysis (Zhang and Hewitt 1996), a number of factors make it unlikely that our sequences are nuclear copies. In addition to straightforward alignment and lack of insertions or deletions, heme-ligating histidines and other conserved residues (Howell 1989) were identified, and no coamplified DNA product (i.e. ambiguous sites) was apparent on the sequencing gels. For birds, nuclear copies are most often associated with the use of blood as a DNA source (Johnson and Sorenson 1998); we used muscle or liver tissue exclusively. Furthermore, the nucleotide bias observed in gene partitions of this data set is consistent with that reported for other birds (Moore and DeFilippis 1997). Lastly, our results are reasonable from both a biological and geographical perspective, as we show below.

Of the 1,143 aligned sites, 240 were variable, and of those, 195 were potentially phylogenetically informative (Table 3). Among those sites, 164 (84%) were at third positions, 28 (14.5%) at first positions, and only 3 (1.5%) at second position sites. Relative rate calculations indicate

TABLE 3. Overall and codon position-specific dynamics of the cytochrome-*b* gene for all taxa. Mean base composition is averaged over all sequences using PAUP*. Transition-transversion ratio (Ts/Tv) values are the average number of changes reconstructed on the weighted parsimony topology.

Position	Number of sites	Variable sites	Parsimony informative	Relative rate	%A	%C	%G	%T	Ts/Tv	α
All	1143	240	195	16.0	27.3	34.7	13.3	24.7	5.08	0.106
1st	381	33	28	6.6	25.6	30.0	23.4	21.0	8.60	0.007
2nd	381	5	3	1.0	20.7	25.4	12.6	41.3	1.50	∞
3rd	381	202	164	40.4	35.5	48.8	3.8	11.9	4.91	1.07

that third-position sites are evolving 40 $\times$ faster than are second-position sites. Nucleotide compositional bias was most acute at third codon positions (0.173), followed by second (0.058) and first (0.006) positions. That was due mainly to the expected strong bias against G (3.8%) at that site. Overall, base composition (Table 3) and base-composition biases are similar to those recovered in other avian *cyt-b* studies (e.g. Kornegay et al. 1993, Nunn et al. 1996).

Most empirical gamma-shape parameter estimates (α) vary from 0.1 to 0.5 (Yang 1996) and the composite sequence estimate for those data falls within this range. However, α estimates for first and third positions differ by two orders of magnitude (0.007 vs. 1.07), suggesting that severe among-site rate heterogeneity exists at first positions, whereas the higher value associated with third positions suggests a more even distribution of substitutions. Despite the fact that the preponderance of phylogenetically informative sites are at third positions, we note that the overall α estimate (0.106) imposed differs from the estimate for only third positions by an order of magnitude.

Uncorrected percentage sequence divergence (p , Table 4) between *Passerina* species ranges from 3.0% (*P. versicolor*–*P. ciris*) to 7.6% (*P. cyanea*–*P. leclancherii*) and averages 6.5 ($\pm 1.0\%$, $n = 84$). Minimum within-species pairwise values range from 0.1% (*P. ciris*) to 0.4% (*P. cyanea* and *G. caerulea*). Overall, intraspecific divergence was 0.24 ($\pm 0.1\%$, $n = 7$), less than the empirical value of 0.7% that was calculated for a range of avian taxa (Moore 1995). Despite the relatively low divergence values, saturation plots for the combined dataset (*Passerina*, *G. caerulea*, and *Cyanocompsa*) provide evidence that multiple substitutions have begun to accumulate in third position transitions (Fig. 1, other [linear] plots not shown).

For these data, log-likelihoods derived from analyses with and without the assumption of a molecular clock (Table 5) did not differ significantly ($0.5 > P > 0.1$) suggesting that these taxa display an approximately uniform rate of substitution such that inference of divergence times is appropriate.

Phylogenetic analyses.—From the many analyses performed, two fully resolved topologies were consistently recovered (Fig. 2A, C). They differ only in the placement of *P. leclancherii*. Equally weighted parsimony, distance, and the F81 maximum-likelihood analysis yield a topology in which the Mexican endemics *P. leclancherii* and *P. rositae* are sisters (Fig. 2A), whereas in all weighted parsimony and other likelihood analyses, *P. leclancherii* is sister to the well supported *P. ciris*–*P. versicolor* clade (Fig. 2C). With the data weighted equally, those topologies differ by a single step (Fig. 2A: length [l] = 419, CI = 0.649, RI = 0.742, RC = 0.481; 3c: l = 420, CI = 0.648, RI = 0.740, RC = 0.479; Kishino-Hasegawa test [1989], $P = 0.3950$). Because the Figure 2C topology was recovered in all weighted-parsimony analyses and in the best fit maximum-likelihood analysis, we consider this our best estimate of relationships within that group. Lanyon (1993) advises that systematists identify both a “best estimate” and [italics his] a “reliable estimate” of phylogenetic relationships. In that spirit, we consider the more conservative, bootstrapped equally weighted MP tree (Fig. 2B) as our most reliable estimate.

In both fully resolved trees, *P. cyanea* is sister to all other ingroup taxa, although that node is poorly supported in several analyses. Two main clades are supported in both trees, a predominately temperate zone (“blue” = *G. caerulea* + *P. amoena*) clade and a more sedentary group (“painted” = *P. rositae* + *P. leclancherii* +

TABLE 4. Observed overall genetic pairwise distances for complete cytochrome-*b* gene. Values above the diagonal are Kimura two-parameter genetic distances corrected for among-site rate heterogeneity; uncorrected (*p*) sequence divergences below the diagonal.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
1 <i>P. cyanea</i> 2		0.004	0.091	0.088	0.088	0.090	0.096	0.096	0.089	0.090	0.100	0.099	0.081	0.082	0.106	0.093	0.106
2 <i>P. cyanea</i> 1	0.004		0.096	0.093	0.090	0.088	0.099	0.099	0.090	0.092	0.108	0.104	0.085	0.087	0.111	0.098	0.108
3 <i>G. caerulea</i> 1	0.069	0.072		0.004	0.061	0.062	0.090	0.093	0.088	0.090	0.104	0.100	0.092	0.094	0.124	0.116	0.120
4 <i>G. caerulea</i> 2	0.067	0.070	0.004		0.056	0.057	0.087	0.090	0.086	0.087	0.101	0.097	0.090	0.091	0.121	0.113	0.114
5 <i>P. amoena</i> 1	0.067	0.068	0.050	0.046		0.003	0.075	0.075	0.089	0.090	0.095	0.091	0.089	0.091	0.112	0.155	0.113
6 <i>P. amoena</i> 2	0.068	0.067	0.051	0.047	0.003		0.072	0.072	0.084	0.086	0.096	0.092	0.091	0.092	0.110	0.113	0.115
7 <i>P. rositae</i> 1	0.072	0.073	0.068	0.066	0.059	0.058		0.002	0.075	0.078	0.082	0.081	0.074	0.076	0.111	0.110	0.117
8 <i>P. rositae</i> 2	0.072	0.073	0.070	0.068	0.059	0.058	0.002		0.075	0.078	0.082	0.081	0.074	0.076	0.116	0.107	0.120
9 <i>P. versicolor</i> 2	0.067	0.068	0.067	0.066	0.067	0.065	0.059	0.059		0.002	0.085	0.084	0.034	0.033	0.130	0.118	0.123
10 <i>P. versicolor</i> 1	0.068	0.069	0.068	0.066	0.068	0.066	0.060	0.060	0.002		0.089	0.087	0.034	0.033	0.132	0.120	0.125
11 <i>P. leclancherii</i> 1	0.076	0.076	0.077	0.075	0.072	0.073	0.064	0.064	0.065	0.066		0.003	0.090	0.091	0.116	0.119	0.128
12 <i>P. leclancherii</i> 2	0.073	0.076	0.074	0.073	0.069	0.070	0.063	0.063	0.064	0.066	0.003		0.088	0.089	0.115	0.117	0.124
13 <i>P. ciris</i> 2	0.063	0.066	0.070	0.068	0.068	0.069	0.059	0.059	0.030	0.030	0.067	0.066		0.001	0.122	0.120	0.120
14 <i>P. ciris</i> 1	0.064	0.066	0.071	0.069	0.069	0.070	0.059	0.059	0.029	0.029	0.068	0.067	0.001		0.124	0.122	0.118
15 <i>P. parviflora</i>	0.078	0.080	0.087	0.085	0.080	0.080	0.080	0.082	0.089	0.090	0.083	0.082	0.086	0.087		0.091	0.099
16 <i>C. brissoni</i>	0.070	0.073	0.082	0.080	0.082	0.081	0.078	0.078	0.083	0.084	0.085	0.084	0.085	0.086	0.069		0.078
17 <i>C. cyanoides</i>	0.078	0.079	0.085	0.081	0.081	0.082	0.083	0.085	0.086	0.087	0.089	0.087	0.085	0.084	0.073	0.060

FIG. 1. Saturation plot for the cytochrome-*b* gene, third-position transitions only. Within-species comparisons are included. Similar plots of other codon positions and transformation types (not shown) were approximately linear.

P. versicolor + *P. ciris*) centered in the aridlands of the southwestern United States and western Mexico. Levels of support as measured by decay indices (Fig. 2A) and bootstrap values vary among nodes (Table 5) and between analytical methods. Perhaps the most extreme case of the latter is the node which unites the painted clade (C on Fig. 2C). Under the HKY Γ model, that node has 78% bootstrap support yet its counterpart on the most parsimonious equal-weights topology has a decay index of 1 and <50% bootstrap support. Similarly, support for a basal *P. cyanea* (node B) is moderately high under the equal and 2:1 parsimony analysis but is lacking under the HKY Γ maximum likelihood model.

Both trees demonstrate that the monotypic *G. caerulea*, despite having a mass approximately twice that of most *Passerina* buntings, is embedded within *Passerina*, rendering the genus paraphyletic. In fact, in all analyses *G. caerulea* and *P. amoena* were sister taxa with relatively high support from both bootstrap and decay values (Table 5). Constraining *P. cyanea* and *P. amoena* to be sisters (the traditional arrangement) with *G. caerulea* sister to all of *Passerina*, requires 17 additional (equal weight) steps and results in a significantly worse topology (Kishino-Hasegawa [1989] test, $P = 0.0498$). Likewise, forcing *G. caerulea* to be sister to the similarly sized spe-

TABLE 5. Levels of bootstrap support for select nodes (see Fig. 2c) for combined data using maximum parsimony and maximum likelihood analyses. All MP (×500) and ML (×100) replicates included full heuristic searches. For ML estimates, empirical nucleotide frequencies were used and α (=0.106) and T_s/T_v (=6.92) were estimated from the data under the HKY model of evolution.

Node	Parsimony ^a								Likelihood ^b			
	=W	×2	×5	×9.5	Y	6P	CS	How	F81	HKY	HKYT	HKYT + c
A	99	99	98	95	96	95	99	98	100	100	100	—
B	71	71	58	<50	58	64	59	54	74	60	51	—
C	—	52	71	75	72	61	74	67	—	63	78	—
D	—	52	75	82	76	<50	70	75	—	55	62	—
E	96	99	97	91	97	96	100	94	95	98	99	—
F	99	99	99	100	99	97	100	98	100	100	99	—
G	73	81	80	84	84	83	82	71	69	80	82	—
—Ln length	419								3,984.3*	3,756.6*	3,568.4	3,576.4
CI	0.649	0.671	0.715	0.747	0.703	0.688	0.719	0.717				
RI	0.742	0.756	0.786	0.809	0.781	0.767	0.791	0.791				
RC	0.481	0.508	0.562	0.605	0.549	0.528	0.569	0.567				

^a For MP, ×9.5 = native rate weighting, Y = (Yoder) ×5 weighting of third position transversions only, 6P = six parameter weighting, CS = codon specific weighting, How = weighted according to Howell's (1989) *cyt-b* variability categories (see text).
^b LRTs indicate that HKY is a significant improvement over F81 (null hypothesis: transition rate equals transversion rate is rejected [−2 log Δ = 455.4, df = 1, $P \ll .001$]); HkYΓ is significantly better than uncorrected HKY (null hypothesis: equal rates among sites is rejected [−2 log Δ = 376.4, df = 1, $P \ll .001$]). LRT for molecular clock is not significantly different (−2 log Δ = 16, df = 15, $0.5 < P < 0.01$). Unconstrained ML = 2985.44.

FIG. 2. Results of analyses. (A) Topology derived from equally weighted MP analysis; decay indices are shown above nodes. Branch lengths shown are proportional to genetic distances derived from an Γ -corrected (α = 0.106) Tamura-Nei (1993) neighbor-joining algorithm. (B) bootstrapped (×500) equal weights MP tree. (C) Bootstrapped (×100) ML tree with best likelihood score (HKYT); α = 0.106, T_s/T_v = 6.93. Lettered nodes refer to Table 5.

cies of *Cyanocompsa* yields an even worse tree (33 steps, $P = 0.0023$). Thus, the mitochondrial gene tree is unambiguous regarding the placement of *G. caerulea*. The other strongly supported terminal taxon pair is *P. versicolor*–*P. ciris*, receiving no less than 97% bootstrap support in any analysis.

DISCUSSION

Passerina paraphyly.—A surprising finding of this study was the high bootstrap support across all analyses (average of 96.5%) for a *G. caerulea*–*P. amoena* sister relationship, demonstrating that *G. caerulea* is derived from within the *Passerina* assemblage. This result indicates that a merging of the monotypic genus *Guiraca* into *Passerina* is warranted. The conventional “superspecies” groupings of *P. cyanea*–*P. amoena* with *P. versicolor* (Mengel 1970) or *P. cyanea*–*P. amoena* with *P. versicolor* and *P. ciris* (Mayr and Short 1970) are rejected by our data. Those workers who, on the basis of morphological and behavioral evidence, had considered *G. caerulea* to be a member of the genus *Passerina* (despite its much larger size; e.g. Phillips et al. 1964, Blake 1969, Mayr and Short 1970) are vindicated; although, we know of no one who hypothesized a *G. caerulea*–*P. amoena* sister relationship. Our result conflicts with conclusions drawn from a family-level allozyme study (Tamplin et al. 1993). Those authors concluded that *G. caerulea* might be more closely related to *Pheucticus* than to either *Passerina* or *Cyanocompsa*; however, they noted that the strong plumage and vocal similarities between *G. caerulea* and the *Passerina* buntings, seem unlikely to be merely shared primitive characters. Indeed, their (Tamplin et al. 1993; Fig. 2) consensus of 172 most parsimonious trees does not support a *Guiraca*–*Pheucticus* relationship. Although mtDNA and allozyme surveys are often congruent at the intrageneric level, the apparent lack of resolution in the Tamplin et al. (1993) study may reflect a taxon sampling problem or the limited resolving power of allozymes for higher (intergeneric) taxonomic levels.

We suggest that if *G. caerulea* were the size of the other buntings (adult males, approximately 15–20 g, see below), it would have long been classified as a member of *Passerina*, perhaps even as sister to *P. amoena*. Study-skin compar-

isons provided additional evidence in support of that relationship. Among members of this genus, only adult males of *P. amoena* and *G. caerulea* have conspicuous, complete wingbars in definitive plumage, and only in the plumage of this pair does a cinnamon (closest to Mikado Brown [no. 121C], of Smithe 1975) color occur (similar to that of *Cyanocompsa* females). Black streaking dorsally is also most conspicuous in this pair although it occurs to a lesser degree in *P. cyanea*. *Guiraca caerulea* (nearest Smalt Blue [no. 70], Smithe 1975) and *P. amoena* (nearest Turquoise Blue [no. 65], Smithe 1975) are rather different shades of blue. Curiously, both of those shades occur in *P. cyanea*, the blue of *G. caerulea* occurring on the head and throat and the blue of *P. amoena* occurring on the lower body (rump).

The “painted” clade (node C, Fig. 2C) is united similarly by plumage characteristics. All have some blue plumage, but it is the presence of patches of bright carotenoid pigments (yellow, pink, scarlet) and conspicuous eye rings that sets them apart. It is selection on precisely these sorts of sexually and socially selected traits that is presumed to lead to accelerated divergence (West-Eberhard 1983, Zink 1996), speciation, and ultimately taxonomic diversity. The short internode distances within the “painted” clade (e.g. node D, Fig. 2A, C) is consistent with that interpretation, that is, closely spaced speciation events.

Ecological implications.—The *Passerina* buntings (exclusive of *G. caerulea*) possess some of the smaller body sizes in the Cardinalidae, as shown by the morphometric phenetic analyses (PCA plots) of that group (Hellack and Schnell 1977, Tamplin et al. 1993). With the addition of *G. caerulea*, *Passerina* becomes composed of two discrete size classes (“bunting” and “grosbeak,” see Fig. 3). Notably, *P. amoena* and *G. caerulea* are closely related sister species, yet the mass of the former is approximately one-half that of the latter. A comparable size dichotomy is apparent within the sister genus *Cyanocompsa* where *C. parcellina* (15.9 ± 1.23 g, $n = 19$) and *C. cyanoides* (35.8 ± 2.77 g, $n = 13$) also exhibit a bunting versus grosbeak size relationship whereas a third *Cyanocompsa* member, *brissonii*, is reportedly of an intermediate size. Collectively, these observations suggest that body size (mass) is not a particularly conservative trait within this assemblage of birds.

FIG. 3. Approximate breeding distributions and "sizes" of all *Passerina* members shown in a phylogenetic context. Note that the distribution of *Passerina rositae* is restricted to the Pacific Slope of the Isthmus of Tehuantepec. Silhouette sizes reflect relative average mass, obtained from museum labels of adult male specimens: *P. cyanea* (14.8 ± 0.8 g, $n = 31$); *P. amoena* (16.8 ± 2.23 g, $n = 4$); *P. rositae* (20.07 ± 0.16 g, $n = 61$); *P. leclancherii* (13.70 ± 0.19 g, $n = 105$); *P. versicolor* (13.0 ± 1.78 g, $n = 104$); *P. ciris* (16.47 ± 2.11 g, $n = 9$); and *G. caerulea* (32.6 ± 2.87 g, $n = 15$).

The phylogeny shown (Fig. 2C) indicates that "bunting" size is the ancestral state for *Passerina* with *G. caerulea* secondarily acquiring a "grosbeak" size. Similarities in the size and shape of *Passerina* buntings, taken together with their relative lack of range overlap (Fig. 3), suggest the possibility that many, or all, are "ecological equivalents" and by extension, competitors for resources. A composite map of all bunting breeding distributions (not shown) indicates that virtually all suitable bunting habitat throughout North America is occupied by one bunting species or another. The relatively huge and overlapping range of *G. caerulea* with respect to those of the smaller buntings (Fig. 3) is likely a consequence of its now dissimilar size and food resource needs. We do not know whether that morphological shift was the

direct result of competition with congeners (ecological character displacement) or if it occurred in response to alternative selection pressures. Character displacement as a result of interspecific competition is difficult to demonstrate (Schluter et al. 1985) but may be tested by correlating shifts in morphology with degree of sympatry (or allopatry) with competitors. *Guiraca caerulea*, with seven described subspecies (Storer and Zimmerman 1959), varies in size across its range and should be studied in that context. A hypothesis of character displacement predicts that the largest *G. caerulea* should occur where the species is sympatric with high *Passerina* densities and the smallest in regions unoccupied (or sparsely populated) by any *Passerina* species.

Little is known about intrageneric competitive interactions for this group. Examination of breeding-bird survey maps (Price et al. 1995) suggests that actual overlaps in breeding distribution may be less extensive than that typically depicted in maps of breeding distributions (such as those shown, Fig. 3), and where overlap does occur, little is known about partitioning of habitat and food resources. In areas of overlap on the western Great Plains, the similarly sized but nonsister species *P. cyanea* and *P. amoena* defend interspecific territories (Emlen et al. 1975). Interspecific territoriality does not occur between the more broadly overlapping pair *P. cyanea* and *P. ciris* (Forsythe 1974), suggesting perhaps a diminution of competitive interactions over time and phylogenetic distance. Also, *P. cyanea* does not defend territories against the visually similar but much larger *G. caerulea* (Payne 1992). Indeed, the emphasis on breeding distributions may be overstated; competitive interactions between species of *Passerina* might be best understood by a winter study in southwestern Mexico where six of the seven taxa occur in sympatry.

Approximate times of divergence.—A temporal interpretation of the phylogeny requires a roughly uniform substitution rate among taxa. The likelihood-ratio test for molecular clock indicates base substitutions are accumulating in a sufficiently clock-like manner to allow such interpretations. We acknowledge that application of a molecular clock remains controversial (e.g. Hillis et al. 1996, Klicka and Zink 1997) but we are also aware of the heuristic value of assigning tentative divergence dates to phyloge-

netic branching events. We use two independently derived clock calibrations. The commonly used rate of 2% divergence per million years is the rate obtained by independent fossil calibrations from an array of avian orders (Klicka and Zink 1997). That this same rate is also derived for mammals (Wilson et al. 1985) and arthropods (Brower 1994) lends some credibility to its generality. Because variation in evolutionary rates has been shown to be related to variation in body size and its correlates (e.g. Martin and Palumbi 1993), the rate of 1.6% per million years calculated by Fleischer et al. (1998) is relevant to our study. It is based on *cyt-b* (only) divergences of small songbirds (Hawaiian Honeycreepers, Drepanididae), with multiple, well founded calibration dates based on emergence times of three different islands. Thus, we have two evolutionary rate estimates that provide a reasonable range of potential divergence times, a 2% rate with uncorrected *p* distances and, following the method of Fleischer et al. (1998), a 1.6% rate using corrected (K2-P) distances (Table 4).

These rates suggest that *Cyanocompsa* and *Passerina* buntings diverged from a common ancestor approximately 4.1 to 7.3 Mya. This coincides with a Late Miocene period of accelerated cooling and drying during which forests and woodlands gave way to an increasing variety of grassland habitats in western North America (Riddle 1995 and references therein). A reduction in size and an increased ability to exploit grassland food resources may played a role in a *Passerina* radiation (diversification and widespread distribution) at that time. This time frame is supported by a *Passerina* fossil (Steadman and McKittrick 1982) from Chihuahua, Mexico, dated to ~4 Mya. The fossil fragments identify an extinct *Passerina* species that was intermediate between *P. amoena* and *G. caerulea* both qualitatively and in size, and from a region where both presently occur.

Among the other well-resolved nodes, the calibrations indicate that *G. caerulea*–*P. amoena* diverged approximately 2.4 to 3.7 Mya and the most recent split within the group, *P. ciris*–*P. versicolor*, occurred 1.5 to 2.1 Mya. The bunting "pair" *P. amoena*–*P. cyanea* has figured prominently (see Sibley and Short 1959) in the development of evolutionary models of songbird evolution. Typically, those (e.g. Mengel 1970, Hubbard 1973) invoke recent glacial advances

(i.e. the last 250,000 years) that reconfigure habitats in such a way that populations fragment and subsequently speciate while in isolation. This "Late Pleistocene Origins" model was rejected using molecular data (Klicka and Zink 1997, 1998), a conclusion supported by this study. Here we show that *P. amoena*–*P. cyanea* are not even sister taxa and that it is unlikely that any members of this genus originated as a consequence of Late Pleistocene glaciations.

Collectively, the lack of consistent bootstrap support at nodes B, D, and E (Fig. 2C) suggest a burst of simultaneous speciation among four *Passerina* lineages, with the *P. amoena*–*G. caerulea* and *P. versicolor*–*P. ciris* lines splitting again at some later time. Zink et al. (1998) note that "star phylogenies" such as this might also be arrived at via bursts of extinction. Alternatively, additional data may change that interpretation entirely by strengthening the nodes in question. We are currently investigating that possibility by sequencing additional gene regions (ND6 and control region) for all *Cyanocompsa* and *Passerina* taxa. A thorough biogeographic assessment requires the teasing apart of these alternatives.

Hybridization as a measure of relationship.—Because a well-documented and extensive hybrid zone exists where they are sympatric on the Great Plains (Sibley and Short 1959, Emlen et al. 1975, Kroodsmma 1975), *P. amoena* and *P. cyanea* have been treated as conspecifics by some (e.g. Phillips et al. 1964) and considered a "superspecies" of very recent origin by others (Mayr and Short 1970). A single hybrid specimen is known for another partly sympatric bunting sister pair, *P. ciris* and *P. versicolor* (Storer 1961). The inference drawn from that observation is that *P. amoena* and *P. cyanea* are the more closely related of those pairs (e.g. Storer 1961). The results of this study challenge that interpretation. *Passerina amoena* and *P. cyanea* are not sister species, whereas *P. ciris* and *P. versicolor* have the most recent origins of any extant members of the genus. It is becoming increasingly clear that for birds, the ability to hybridize and the degree to which hybridization occurs are not necessarily good measures of relatedness (Prager and Wilson 1975). We add *P. amoena*–*P. cyanea* to the growing list of nonsister taxa that not only retain the ancestral capacity to hybridize, but also to form an ap-

parent hybrid swarm (e.g. *Colaptes*, Moore et al. 1991; *Icterus*, Freeman and Zink 1995).

ACKNOWLEDGMENTS

This paper is dedicated to the life and memory of Allan Galante who loved Mexico, its land, people, culture, music, and natural history, including its bird life. C.W.T. is grateful for financial and logistical support from (1) the Eddy Endowment for Excellence in Ornithology, (2) C. Wood, C. Spaw, and the Rohwer Lab Group at the Burke Museum, University of Washington, (3) two Chicago Zoological Society Society Conservation and Research Fund awards, (4) a Center for the Study of Tropical Birds award, (5) Adolfo Navarro and Octavio Rojas at the Museo de Zoología at the Universidad Nacional Autónoma de México, (6) Eduardo Morales and Eduardo Espinosa of the Instituto de Historia Natural, Gobierno de Chiapas, México, and (7) the Western Foundation of Vertebrate Zoology. We thank the curators and staff at the Louisiana State University Museum of Zoology and the University of Washington Burke Museum for acquiring and loaning tissue specimens. A. Phillips provided valuable field assistance as did T. Brashear. J.K. acknowledges funding support from the Dayton and Wilkie Funds of the J. F. Bell Museum of Natural History. Funding for laboratory work was provided by R.M.Z.'s NSF grant DEB-9317945.

LITERATURE CITED

- AMERICAN ORNITHOLOGISTS' UNION. 1998. Check-list of North American birds, 7th ed. American Ornithologists' Union, Washington, D. C.
- BLAKE, C. H. 1969. Notes on the Indigo Bunting. *Bird-Banding* 40:133-139.
- BREMER, K. 1994. Branch support and tree stability. *Cladistics* 10:295-304.
- BROOKS, D. R., AND D. A. McLENNAN. 1991. *Phylogeny, Ecology, and Behavior: A Research Program in Comparative Biology*. University of Chicago Press, Chicago.
- BROWER, A. V. Z. 1994. Rapid morphological radiation and convergence among races of the butterfly *Heliconius erato* inferred from patterns of mitochondrial DNA evolution. *Proceedings of the National Academy of Sciences USA* 91:6491-6495.
- CUNNINGHAM, C. 1997. Is congruence between data partitions a reliable predictor of phylogenetic accuracy? Empirically testing an iterative procedure for choosing among phylogenetic methods. *Systematic Biology* 46:464-478.
- DESJARDINS, P., AND R. MORAIS. 1990. Sequence and gene organization of the chicken mitochondrial genome: A novel gene order in higher vertebrates. *Journal of Molecular Biology* 212:599-634.
- ELLEGREN, H. 1992. Polymerase-chain-reaction (PCR) analysis of microsatellites—A new approach to studies of genetic relationships in birds. *Auk* 109:886-895.
- EMLEN, S. T., J. D. RISING, AND W. THOMPSON. 1975. A behavioral and morphological study of sympatry in the Indigo and Lazuli buntings of the Great Plains. *Wilson Bulletin* 87:145-179.
- FELSENSTEIN, J. 1981. Evolutionary trees from DNA sequences: A maximum likelihood approach. *Journal of Molecular Evolution* 17:368-376.
- FELSENSTEIN, J. 1985. Confidence limits on phylogenies: An approach utilizing the bootstrap. *Evolution* 39:783-791.
- FLEISCHER, R. C., C. E. MCINTOSH, AND C. L. TARR. 1998. Evolution on a volcanic conveyor belt: Using phylogeographic reconstructions and K-Ar based ages of the Hawaiian Islands to estimate molecular evolutionary rates. *Molecular Ecology* 7:533-545.
- FORSYTHE, D. M. 1974. Song characteristics of sympatric and allopatric Indigo and Painted bunting populations in the southeastern United States. Ph.D. dissertation, Clemson University, Clemson, South Carolina.
- FREEMAN, S., AND R. M. ZINK. 1995. A phylogenetic study of the blackbirds based on variation in mitochondrial restriction sites. *Systematic Biology* 44:409-420.
- GOLDMAN, N. 1993. Statistical tests of models of DNA substitution. *Journal of Molecular Evolution* 36:182-198.
- GRIFFITHS, C. S. 1997. Correlation of functional domains and rates of nucleotide substitution in cytochrome *b*. *Molecular Phylogenetics and Evolution* 7:352-365.
- HACKETT, S. J. 1996. Molecular phylogenetics and biogeography of tanagers in the genus *Ramphocelus* (Aves). *Molecular Phylogenetics and Evolution* 5:368-382.
- HARSHMAN, J. 1996. Phylogeny, evolutionary rates, and ducks. Ph.D. dissertation, University of Chicago, Chicago.
- HASEGAWA, M., M. KISHINO, AND T. YANO. 1985. Dating of the human-ape splitting by a molecular clock of mitochondrial DNA. *Journal of Molecular Evolution* 22:160-174.
- HELLACK, J. J., AND G. SCHNELL. 1977. Phenetic analysis of the subfamily Cardinalinae using external and skeletal characters. *Wilson Bulletin* 89:130-148.
- HILLIS, D. M., B. K. MABLE, A. LARSON, S. K. DAVIS, AND E. A. ZIMMER. 1996. Nucleic Acids IV: Sequencing and cloning. Pages 321-381 in *Molecular Systematics* (D. M. Hillis, C. Moritz, and B. K. Mabel, Eds.). Sinauer Associates, Sunderland, Massachusetts.
- HOWELL, N. 1989. Evolutionary conservation of protein regions in the protonmotive cytochrome *b*

- and their possible roles in redox catalysis. *Journal of Molecular Evolution* 29:157–169.
- HUBBARD, J. P. 1973. Avian evolution in the aridlands of North America. *Living Bird* 12:155–196.
- HUELSENBECK, J. P. AND B. RANNALA. 1997. Phylogenetic methods come of age: Testing hypotheses in an evolutionary context. *Science* 276:227–232.
- JOHNSON, K. P. AND M. D. SORENSON. 1998. Comparing molecular evolution in two mitochondrial protein coding genes (cytochrome *b* and ND2) in the dabbling ducks (Tribe: Anatini). *Molecular Phylogenetics and Evolution* 10:82–94.
- KIMURA, M. 1980. A simple method for estimating evolutionary rates of base substitution through comparative studies of nucleotide sequences. *Journal of Molecular Ecology* 16:11–120.
- KISHINO, H., AND M. HASEGAWA. 1989. Evaluation of the maximum likelihood estimate of the evolutionary tree topologies from DNA sequence data, and the branching order in Hominoidea. *Journal of Molecular Evolution* 29:170–179.
- KLICKA, J., J. C. BARLOW, W. B. MCGILLIVRAY, T. J. DOYLE, AND R. M. ZINK. 1999. Evidence supporting the recent origin and species status of the Timberline Sparrow. *Condor* 101:577–588.
- KLICKA, J., AND R. M. ZINK. 1997. The importance of recent ice ages in speciation: A failed paradigm. *Science* 277:1666–1669.
- KLICKA, J., AND R. M. ZINK. 1998. Pleistocene speciation and the mitochondrial DNA clock: A response to Arbogast and Slowinski. *Science* 282:1955.
- KOCHER, T. D., W. K. THOMAS, A. MEYER, S. V. EDWARDS, S. PAABO, F. X. VILLABLANCA, AND A. C. WILSON. 1989. Dynamics of mitochondrial DNA evolution in animals: Amplification and sequencing with conserved primers. *Proceedings of the National Academy of Sciences USA* 86: 6196–6200.
- KORNEGAY, J. R., T. D. KOCHER, L. A. WILLIAMS, AND A. C. WILSON. 1993. Pathways of lysozyme evolution inferred from the sequences of cytochrome *b* in birds. *Journal of Molecular Evolution* 37:367–379.
- KRAJEWSKI, C., AND D. G. KING. 1996. Molecular divergence and phylogeny: Rates and patterns of cytochrome-*b* evolution in cranes. *Molecular Biology and Evolution* 13:21–30.
- KROODSMA, R. L. 1975. Hybridization in buntings (*Passerina*) in North Dakota and eastern Montana. *Auk* 92:66–80.
- LANYON, S. M. 1993. Phylogenetic frameworks: Towards a firmer foundation for the comparative approach. *Biological Journal of the Linnean Society* 49:45–61.
- MADDISON, W. P., AND D. R. MADDISON. 1992. MacClade: Analysis of phylogeny and character evolution, 3.04. Sinauer Associates, Sunderland, Massachusetts.
- MARTIN, A. P., AND S. R. PALUMBI. 1993. Body size, metabolic rate, generation time, and the molecular clock. *Proceedings of the National Academy of Sciences USA* 90:4087–4091.
- MAYR, E., AND L. L. SHORT. 1970. Species taxa of North American birds, A contribution to comparative systematics. Publications of the Nuttall Ornithological Club, no. 9.
- MENGEL, R. M. 1970. The North American central plains as an isolating agent in bird speciation. Pages 279–340 in *Pleistocene and recent environments of the central Great Plains* (W. Dort, Jr., and J. Knox Jones, Jr., Eds.). University Press of Kansas, Lawrence.
- MOORE, W. S. 1995. Inferring phylogenies from mtDNA variation: Mitochondrial-gene trees versus nuclear-gene trees. *Evolution* 49:718–726.
- MOORE, W. S., AND V. R. DE FILIPPIS. 1997. The window of taxonomic resolution for phylogenies based on mitochondrial cytochrome *b*. Pages 83–119 in *Avian Molecular Evolution and Systematics* (D. P. Mindell, Ed.). Academic Press, New York.
- MOORE, W. S., J. H. GRAHAM, AND J. T. PRICE. 1991. Mitochondrial DNA variation in the Northern Flicker (*Colaptes auratus*). *Molecular Biology and Evolution* 8:327–344.
- NUNN, G. B., J. COOPER, P. JOUVENTIN, C. J. R. ROBERTSON, AND G. G. ROBINSON. 1996. Evolutionary relationships among extant albatrosses (Procellariiformes: Diomedidae) established from complete cytochrome-*b* gene sequences. *Auk* 113:784–801.
- PAYNE, R. B. 1992. Indigo Bunting (*Passerina cyanea*). In *The Birds of North America*, no. 4 (A. Poole, P. Stettenheim, and F. Gill, Eds.). Academy of Natural Sciences, Philadelphia, and American Ornithologists' Union, Washington, D.C.
- PAYNTER, R. A. 1970. Subfamily Cardinalinae. Pages 216–245 in *Check-list of Birds of the World*, vol. 13. (R. A. Paynter and R. W. Storer, Eds.). Museum of Comparative Zoology, Heffernan Press, Worcester, Massachusetts.
- PHILLIPS, A. R., J. T. MARSHALL, JR., AND G. MONSON. 1964. *The Birds of Arizona*. University of Arizona Press, Tucson.
- PRAGER, E. M., AND A. C. WILSON. 1975. Slow evolutionary loss of the potential for interspecific hybridization in birds: A manifestation of slow regulatory evolution. *Proceedings of the National Academy of Sciences USA* 72:200–204.
- PRICE, J., S. DROEGE, AND A. PRICE. 1995. *The Summer Atlas of North American Birds*. Academic Press, San Diego, California.
- QUINN, T. W., AND A. C. WILSON. 1993. Sequence evolution in and around the mitochondrial control region. *Journal of Molecular Evolution* 37:417–425.

- RIDDLE, B. R. 1995. Molecular biogeography in the pocket mice (*Perognathus* and *Chaetodipus*) and grasshopper mice (*Onychomys*): The late cenozoic development of a North American aridlands rodent guild. *Journal of Mammalogy* 76: 283–301.
- RIDGWAY, R. 1901. The birds of North and Middle America. *Bulletin of the United States National Museum*, no. 50, Part 1.
- SCHLUTER, D., T. D. PRICE, AND P. R. GRANT. 1985. Ecological character displacement in Darwin's finches. *Science* 227:1056–1059.
- SIBLEY, C. G., AND B. L. MONROE. 1990. *Distribution and Taxonomy of Birds of the World*. Yale University Press, New Haven, Connecticut.
- SIBLEY, C. G., AND L. SHORT, JR. 1959. Hybridization in the buntings (*Passerina*) of the Great Plains. *Auk* 76:443–463.
- SMITH, A. B. 1994. Rooting molecular trees: Problems and strategies. *Biological Journal of the Linnean Society* 51:279–292.
- SMITHE, F. B. 1975. *Naturalists' Color Guide*. The American Museum of Natural History, New York.
- SORENSEN, M. D. 1996. "Treeroot." University of Michigan, Ann Arbor.
- STEADMAN, D. W., AND M. C. MCKITRICK. 1982. A Pliocene bunting from Chihuahua, Mexico. *Condor* 84:240–241.
- STORER, R. W. 1961. A hybrid between the Painted and Varied buntings. *Wilson Bulletin* 73:209.
- STORER, R. W., AND D. A. ZIMMERMAN. 1959. Variation in the Blue Grosbeak (*Guiraca caerulea*) with special reference to the Mexican populations. *Occasional Papers of the Museum of Zoology University of Michigan* 609:1–13.
- STURMBAUER, C., AND A. MEYER. 1992. Genetic divergence, speciation and morphological stasis in a lineage of African cichlid fishes. *Nature* 358: 578–581.
- SULLIVAN, J., K. E. HOLSINGER, AND C. SIMON. 1995. Among-site rate variation and phylogenetic analysis of 12S rRNA in sigmodontine rodents. *Molecular Biology and Evolution* 12:988–1001.
- SWOFFORD, D. L. 1999. *PAUP: Phylogenetic analysis using parsimony (* and other methods)*, version 4.0b2. Sinauer Associates, Sunderland, Massachusetts.
- SWOFFORD, D. L., G. J. OLSEN, P. J. WADELL, AND D. M. HILLIS. 1996. Phylogenetic inference. Pages 407–514 in *Molecular Systematics* (D. M. Hillis, C. Moritz, and B. K. Mabel, Eds.). Sinauer Associates, Sunderland, Massachusetts.
- TAMPLIN, J. W., J. W. DEMASTES, AND J. V. REMSEN, JR. 1993. Biochemical and morphometric relationships among some members of the Cardinalinae. *Wilson Bulletin* 105:93–113.
- TAMURA, K., AND M. NEI. 1993. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution* 10:512–526.
- TAYLOR, W. K. 1974. A new hybrid bunting (*Passerina cyanea* × *Passerina ciris*). *Auk* 91:485–487.
- THOMPSON, C. W. AND M. LEU. 1994. Determining homology of molts and plumages to address evolutionary questions: A rejoinder regarding Emberizid finches. *Condor* 96:769–782.
- THOMPSON, C. W., AND M. LEU. 1995. Molts and plumages of Orange-breasted Buntings (*Passerini leclancherii*): Implications for theories of delayed plumage maturation. *Auk* 112:1–19.
- THOMPSON, W. L. 1968. The songs of five species of *Passerina*. *Behaviour* 31:261–287.
- VOELKER, G., AND S. V. EDWARDS. 1998. Can weighting improve bushy trees? Models of cytochrome *b* evolution and the molecular systematics of pipits and wagtails (Aves: Motacillidae). *Systematic Biology* 47:589–603.
- WEST-EBERHARD, M. J. 1983. Sexual selection, social competition, and speciation. *Quarterly Review of Biology* 58:155–183.
- WHEELER, W. C. 1990. Nucleic acid sequence phylogeny and random outgroups. *Cladistics* 6:363–368.
- WILLIAMS, P. L., AND W. M. FITCH. 1989. Finding the minimal change in a given tree. Pages 453–470 in *The Hierarchy of Life* (B. Fernholm, K. Bremer, and H. Jornvall, Eds.). Elsevier Press, New York.
- WILSON, A. C., R. L. CAAN, S. M. CARR, M. GEORGE, JR., U. B. GYLLENSTEN, K. M. HELM-BYCHOWSKI, R. G. HIGUCHI, S. R. PALUMBI, E. M. PRAGER, R. D. SAGE, AND M. STONEKING. 1985. Mitochondrial DNA and two perspectives on evolutionary genetics. *Biological Journal of the Linnean Society* 26:375–400.
- YANG, Z. 1996. Among-site rate variation and its impact on phylogenetic analyses. *Trends in Ecology and Evolution* 11:367–372.
- YODER, A. D., R. VILGALYS, AND M. RUVOLO. 1996. Molecular evolutionary dynamics of cytochrome *b* in Strepsirrhine primates: The phylogenetic significance of third position transversions. *Molecular Biology and Evolution* 13:1339–1350.
- ZHANG, D.-X., AND G. M., HEWITT. 1996. Nuclear integrations: Challenges for mitochondrial DNA markers. *Trends in Ecology and Evolution* 11: 247–251.
- ZINK, R. M. 1996. Species concepts, speciation, and sexual selection. *Journal of Avian Biology* 27:1–6.
- ZINK, R. M., AND M. C. MCKITRICK. 1995. The debate over species concepts and its implications for ornithology. *Auk* 112:701–719.
- ZINK, R. M., S. J. WELLER, AND R. C. BLACKWELL. 1998. Molecular phylogenetics of the avian genus *Pipilo* and a biogeographic argument for taxonomic uncertainty. *Molecular Phylogenetics and Evolution* 10:191–201.