

Aug 6th, 9:00 AM - 12:00 PM

## Stereospecificity in glucose consumption: A new approach to martian life detection

Vienna R. Saccomanno  
*Desert Research Institute*

Henry J. Sun  
*Desert Research Institute, Henry.Sun@dri.edu*

Follow this and additional works at: [https://digitalscholarship.unlv.edu/cs\\_urop](https://digitalscholarship.unlv.edu/cs_urop)



Part of the [Soil Science Commons](#), and the [The Sun and the Solar System Commons](#)

### Repository Citation

Saccomanno, Vienna R. and Sun, Henry J., "Stereospecificity in glucose consumption: A new approach to martian life detection" (2008). *Undergraduate Research Opportunities Program (UROP)*. 33.  
[https://digitalscholarship.unlv.edu/cs\\_urop/2008/aug6/33](https://digitalscholarship.unlv.edu/cs_urop/2008/aug6/33)

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Scholarship@UNLV with permission from the rights-holder(s). You are free to use this Event in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/or on the work itself.

This Event has been accepted for inclusion in Undergraduate Research Opportunities Program (UROP) by an authorized administrator of Digital Scholarship@UNLV. For more information, please contact [digitalscholarship@unlv.edu](mailto:digitalscholarship@unlv.edu).


**Vienna R. Saccomanno****Mentor - Henry J. Sun**

In 1976, the Viking mission made a remarkable discovery: Martian soil was capable of decomposing an organic nutrient broth to carbon dioxide as if it contained live microorganisms. However, a biological interpretation of this finding is in apparent contradiction with the gas chromatograph-mass spectrometer aboard the Viking landers, which showed Martian soil to be devoid of indigenous organics. To reconcile these findings, it has been hypothesized that unknown abiotic oxidants, such as peroxide and superoxide, are present on Mars and that they were responsible for its soil reactivity. The objective of this research is to develop a life detection method that can distinguish biological reactivity from abiotic mimicry.

# Stereospecificity in glucose consumption: a new approach to Martian life detection



Vienna R. Saccomanno and Henry J. Sun  
Desert Research Institute, Las Vegas.



## Is Mars a biological planet or a sterile (oxidized) world?


In 1976, the Viking mission made a remarkable discovery: Martian soil was capable of decomposing an organic nutrient broth to carbon dioxide as if it contained live microorganisms. However, a biological interpretation of this finding is in apparent contradiction with the gas chromatograph-mass spectrometer aboard the Viking landers, which showed Martian soil to be devoid of indigenous organics. To reconcile these findings, it has been hypothesized that unknown abiotic oxidants, such as peroxide and superoxide, are present on Mars and that they were responsible for its soil reactivity. The objective of this research is to develop a life detection method that can distinguish biological reactivity from abiotic mimicry.

## Our approach to distinguishing between biological and chemical reactivity




We hypothesize that organic consumption by living organisms is stereospecific, whereas abiotic destruction is not. In other words, if both D- and L-isomers of the same compound are fed to soil but only one is consumed, the soil is considered biologically reactive. On the other hand, if both isomers are destroyed, abiotic oxidants are assumed present.

Below, we provide experimental verification for this theory by demonstrating that: 1) only D-glucose, not L-glucose, is used by terrestrial organisms, and 2) non-life processes are not stereospecific and destroy both D- and L-glucose.

### 1. Glucose consumption by terrestrial life is stereospecific



### 2. Abiotic chemical oxidation is not stereospecific



## Discussion

Our data indicates that stereospecificity is a distinguishing character of life. With the exception of one of the archaea, which is apparently incapable of metabolizing glucose, all organisms studied consumed only D-glucose, not L-glucose. In contrast, chemical oxidation by potassium permanganate destroyed both D- and L-glucose. Further tests with more organisms will determine whether or not stereospecificity is a universal property of terrestrial life.

Future work will also consider lactic acid and amino acids. If these substrates are also utilized in a stereospecific manner like glucose, then they should also be used to maximize the probability of success of life detection. We envision that, once fully established, this new approach can be implemented on Mars to determine whether or not life exists on our neighboring planet.

## Acknowledgements:

This work is supported by a NASA Grant to HJS. Funding also came from NASA EPSCoR and Space Grant. We thank Dr. Brian Hedlund for providing the archaea and for help with their growth, Dr. Kurt Regner for providing the yeast, and Mr. Alex Michaud for technical assistance with the flow cytometer.