

2015

# Propagation of Lightning Through Thick Thunderclouds

Isaiah Henry

*University of Nevada, Las Vegas*, [isiahenry7@gmail.com](mailto:isiahenry7@gmail.com)

Dieudonné D. Phanord

*University of Nevada, Las Vegas*, [dieudonne.phanord@unlv.edu](mailto:dieudonne.phanord@unlv.edu)

Follow this and additional works at: [http://digitalscholarship.unlv.edu/mcnair\\_posters](http://digitalscholarship.unlv.edu/mcnair_posters)

 Part of the [Atmospheric Sciences Commons](#), and the [Physics Commons](#)

---

## Repository Citation

Henry, I., Phanord, D. D. (2015). Propagation of Lightning Through Thick Thunderclouds.

**Available at:** [http://digitalscholarship.unlv.edu/mcnair\\_posters/69](http://digitalscholarship.unlv.edu/mcnair_posters/69)

This Poster is brought to you for free and open access by the McNair Scholars Institute at Digital Scholarship@UNLV. It has been accepted for inclusion in McNair Poster Presentations by an authorized administrator of Digital Scholarship@UNLV. For more information, please contact [digitalscholarship@unlv.edu](mailto:digitalscholarship@unlv.edu).

Isaiah Henry: McNair Scholar; Mechanical Engineering  
 Advised By: Dr. Dieudonné D. Phanord; Professor

## Introduction

❖ The phenomenon that is lightning, have sparked the interest of physicists and scientists for centuries. The journey to understanding this phenomenon of high-current electric discharge, can unlock the secret to other lightning related phenomena. Lightning is a major source of interference in many types of radio communications. The effects of lightning on space crafts, nuclear power plants, and sophisticated military equipment, are problems of increasing concern. The purpose of this research is to study the propagation of lightning through optically thick thunderclouds by applying knowledge of cloud micro-physics, the physics of lightning, diffusion approximations, and an understanding of the scattering problem.

## Research Objectives

- ❖ Simulation of lightning propagation, using a cylindrical shape to model the cloud.
- ❖ Study of the multiple scattering effect for propagation of radiation in random media.



## Method

- ❖ The cloud was treated as a nuclear reactor in order to obtain forms that can be computable.
- ❖ Also, a one-speed Boltzmann transport theory using diffusion approximations is applied to examine the radiative transfer properties of lightning in optically thick thunder clouds.
- ❖ Lastly, the cloud is modeled geometrically, and the effects of multiple scattering are considered.


## Formation of Thick Thunderclouds

- ❖ There are three conditions that are necessary for the formation of a thundercloud: sunlight, moist air, and an unstable atmosphere (Cooray, 2014).

$$CAPE = \int_{Z_f}^{Z_e} g \frac{T_p(Z) - T_e(Z)}{T_e(Z)} dz$$





## Propagation of Lightning



1. **Charge Distribution**
2. **Concentration of negative charges:** 6 to 8 km above the mean sea level (-10 to -20°C)
3. **Preliminary breakdown:** Location; occur between the distribution of + and - charges in the cloud.
4. **Stepped-leader:** Highly branched discharge that propagates horizontally and downward.
5. Electric field just above the surface becomes very large as Stepped-leader approaches the ground.
6. **Attachment process:** Upward discharge begin at the ground and connects with the leader channel at a junction point.
7. **Return stroke:** Begins when contact occurs at the junction point.
8. An intense positive wave of ionization propagates up the leader channel at about one third the speed of light.
9. The current carries the ground potential upward and effectively neutralizes most of the leader channel and a portion of the cloud charge.
10. **Dart leader:** A new leader produced by most cloud-to-ground flashes approximately 40 to 80 milliseconds after the return stroke (Krider, 1986).

## Data Analysis/Results

$$I(\vec{r}, \hat{\Omega}, t) = \frac{2}{\pi V} \sum_{l=1}^{\infty} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left[ \Psi_{lmn}(\vec{r}) - \frac{3D}{c} \hat{\Omega} \right]$$



- ❖ For simplicity, a cylindrical shape is used to model the cloud.
- ❖ The differential equation was solved. Data was obtained directly from its solution.
- ❖ The curve represents the intensity of the lightning over time.
- ❖ Observe that a change in the angle of observation from  $\theta = 45$  to  $\theta = 90$ , results in a higher peak intensity in the propagation of radiation.

## Future Research

- ❖ Analyze data recorded by LIS (lightning Imaging Sensor), a space-based lightning sensor aboard the EOS TRMM satellite. Perform a numerical analysis to help create a more realistic model.

## References

- ❖ Cooray, Vernon. *An Introduction to lightning*. Netherlands: Springer Netherlands, 2014. 71-77. Print.
- ❖ Krider, E. P. "The Earth's Electric Environment (Studies in Geophysics)." *The Physics of Lightning*. Washington DC: National Academy Press, 1986. 30-39. Print.
- ❖ Koshak, W.J., R.J. Solakiewicz, D.D. Phanord, and R.J. Blakeslee (1994). *Diffusion Model for Lightning Radiative Transfer*

## Acknowledgements

- ❖ I will like to thank God for his Mercy and countless blessings. My parents for their love, support, and encouragement. My Advisor and Mentor Dr. Phanord for all the knowledge he has shared with me since I first met him at Space Camp; my sophomore year of high school. Last but not least, a special thank you to the McNair Scholars Institute, for providing this wonderful opportunity to conduct undergraduate research.