

1991

Littoral and limnetic zooplankton communities in Lake Mead, Nevada-Arizona, USA

Patrick J. Sollberger
University of Nevada, Las Vegas

Larry J. Paulson
University of Nevada, Las Vegas

Follow this and additional works at: https://digitalscholarship.unlv.edu/water_pubs

 Part of the [Biology Commons](#), [Environmental Health and Protection Commons](#), [Environmental Indicators and Impact Assessment Commons](#), [Environmental Monitoring Commons](#), [Fresh Water Studies Commons](#), [Natural Resources and Conservation Commons](#), [Terrestrial and Aquatic Ecology Commons](#), and the [Water Resource Management Commons](#)

Repository Citation

Sollberger, P. J., Paulson, L. J. (1991). Littoral and limnetic zooplankton communities in Lake Mead, Nevada-Arizona, USA. *Hydrobiologia* 1-11.

Available at: https://digitalscholarship.unlv.edu/water_pubs/83

This Article is brought to you for free and open access by the Water Resources at Digital Scholarship@UNLV. It has been accepted for inclusion in Publications (WR) by an authorized administrator of Digital Scholarship@UNLV. For more information, please contact digitalscholarship@unlv.edu.

Littoral and limnetic zooplankton communities in Lake Mead, Nevada-Arizona, USA

Patrick J. Sollberger¹ & Larry J. Paulson

Lake Mead Limnological Research Center, University of Nevada, Las Vegas, Las Vegas, 89154 NV, USA; ¹Current address; Southern Nevada Water System, 243 Lakeshore Rd., Boulder City, NV 89005, USA

Received 27 March 1990; in revised form 9 July 1991; accepted 26 July 1991

Key words: littoral, limnetic, species richness, zooplankton abundance, phytoplankton biomass, fish abundance

Abstract

Zooplankton were collected from adjacent littoral and limnetic sites in Lake Mead, Nevada-Arizona, USA. Limnetic species dominated both littoral and limnetic zooplankton communities; littoral species rarely exceeded 2% of monthly total zooplankton densities. Low species richness of littoral taxa and high similarity in species composition between littoral and limnetic habitats appeared to result from uniform horizontal physical and chemical environments, due to horizontal mixing, and from the absence aquatic macrophytes.

Significant differences in spatial distribution occurred in phytoplankton biomass, total zooplankton density, and fish abundances; highest concentrations of these factors occurred nearest an inflow high in nutrients and progressively declined farther below the inflow. These factors generally showed no significant difference between adjacent littoral and limnetic sites. Large variation also occurred in seasonal zooplankton community structure among some sites.

Introduction

Zooplankton are typically considered either littoral or limnetic (Edmondson, 1959; Hutchinson, 1967; Wetzel, 1983; Pennak, 1978). Limnetic communities are frequently dominated by cladocerans, copepods, and rotifers (Pennak, 1957, 1978; Hutchinson, 1967). Particular cladocerans (Smyly, 1952; Straskraba, 1964; Lemly & Dimmick, 1982a) and rotifers (Pennak, 1966) generally dominate littoral habitats where macrophytes are present. Calanoid copepods are less abun-

dant among vegetation and Gerhs (1974) suggested that secretions or physical effects of *Potamogeton* inhibit survival and reproduction of *Diatomus clavipes*. Straskraba (1964) and Pennak (1966) found that cyclopoid copepods generally occurred in low abundance in weedbeds. However, Cryer & Townsend (1988) found cyclopoid copepods had greater concentrations in littoral areas than in limnetic areas. The density of aquatic vegetation largely determines the diversity and abundance of littoral zooplankton (Straskraba, 1964; Pennak, 1966; Lemly & Dim-

mick, 1982a). Common littoral species are usually poorly represented in littoral areas lacking vegetation and in such areas species composition and abundances resemble that of adjacent limnetic areas (Smyly, 1952; Smirnov, 1963; Straskraba, 1964; Stolbunova & Stolbunov, 1981; Lemly & Dimmick, 1982a, b).

Variability in zooplankton horizontal distribution, however, may be greatly influenced by environmental conditions other than aquatic vegetation. Meyer (1984) reported equal numbers of *Daphnia pulex* in the limnetic zone and the littoral-limnetic interface during times of unlimited food availability. When phytoplankton became scarce, it was found grazing predominantly in outer margins of the littoral zone. *Chydorus sphaericus* followed a similar pattern, but ranged farther into the vegetation zone when food was scarce.

Advection by wind can concentrate plankton in down-wind areas along lake edges (George & Edwards, 1976). Hart (1976) noted that copepodite and adult *Pseudodiaptomus hessei* were benthic during the day and were generally undisturbed by wind-induced surface currents. During their migration upward at night, *P. hessei* was dispersed down-wind by surface currents. Naupliar stages that always lived near the surface were influenced by wind activated surface currents during the night and by slightly deeper counter currents during the day (Hart, 1976).

Effects of fish predation also can influence the distribution of zooplankton (Jakobsen & Johnsen, 1987; Cryer & Townsend, 1988). Cryer & Townsend (1988) found that limnetically associated taxa abundantly occupied the lake periphery as a result of large numbers of fish and the ability of fish to feed more efficiently in openwater than in stands of vegetation. During years having fewer fish, zooplankton showed a greater limnetic distribution (Cryer & Townsend, 1988). Siebeck (1980) suggested that limnetic species which become horizontally disoriented and wander inshore migrate back offshore during the day by optically distinguishing differences in light intensities between openwater and shoreline areas.

The objective of our study was to examine ro-

tifer and microcrustacean zooplankton communities in littoral and limnetic areas of Lake Mead, Nevada and Arizona, a large desert impoundment with a littoral zone that is sparsely vegetated. We also discuss possible influences of physical and chemical limnological conditions, relative phytoplankton biomass, and relative fish abundance on zooplankton horizontal distribution and seasonal dynamics.

Study sites

Lake Mead, bordering Nevada and Arizona, USA, is a major reservoir along the Colorado River. It has a surface area of about 66 096 ha, maximum depth of about 180 m, and mean depth of approximately 55 m (Hoffman & Jonez, 1973). Zooplankton were collected from five stations in three areas in the lower basin (Fig. 1). Samples were collected at inner Las Vegas Bay (ILVB), a site consisting of only littoral zone. ILVB had no close limnetic, similarly fertile habitat. Limnetic samples at middle Las Vegas Bay (MLVB) were collected mid-channel at a depth of approximately 40 m and littoral samples were collected in an adjacent nearby cove. Samples were collected at the offshore site at Boulder Basin (BB) having a depth of over 100 m and the adjacent littoral sampling station was located in a large cove on Saddle Island.

Aquatic macrophytes occurred sparsely throughout the reservoir (Haley *et al.*, 1987). In particular, ILVB was depauperate of rooted aquatic vegetation and in MLVB vegetation was very sparse and generally located less than 2 m deep (Haley *et al.*, 1987). Water clarity was greatest in Boulder Basin (summer Secchi depth ranged from 6 to 7 m) and no macrophytes were seen in the littoral area. Because very little vegetation existed in Lake Mead, and the distribution of rooted aquatic vegetation determines the extent of littoral zone (Wetzel, 1983), we identified the littoral zone as the area from shore to a depth of 10 m. Only about 11% of the total surface area of the lower basin then is considered as littoral zone.

Sampling Stations




Fig. 1. Map of Lake Mead showing littoral and limnetic sampling stations in the lower basin.

Methods

Littoral and limnetic zooplankton communities were collected at 1, 3, 5, 7, and 10 m. Littoral samples were collected where bottom depth was 10 m. Samples were collected using a pump sampler with a clearance rate of 20 l min^{-1} . One end of a hose (1.6 cm inner diameter) was attached to the pump and the opposite end had a double plexiglas plate with a 2.5 cm gap to collect an even draw of water. Twenty liters of water were filtered through an $80 \mu\text{m}$ mesh plankton net and zooplankton were preserved in 4% formalin-sucrose solution. Before samples from succeeding depths were collected, the first 20 l of water were discarded to flush the hose of organisms from the previous depth. Samples were collected monthly from July 1984 to June 1985 and between 0800 and 1300 hrs. Entire contents of each sample were counted except for particularly dense samples, where three separate 1 ml subsamples were counted and an average count was taken. Zooplankton communities were compared from water column averages (0 to 10 m).

Chlorophyll fluorescence was measured for each sample using a Turner Designs flow-through fluorometer to estimate relative phytoplankton biomass (see Heaney, 1978). Water temperature, dissolved oxygen, conductivity, and pH were measured with a Hydrolab Model 8000 Water Quality Analyser at each sampling depth. A Furuno Model FM-22A echolocator was used to record fish at each station. Echolocation transects of at least 500 m were run from adjacent limnetic to littoral sites at a constant boat velocity of about 8 km h^{-1} . Relative abundance of fish was then estimated from echograms following the procedure of Wilde & Paulson (1989) by scoring from 1 (no fish) to 5 (maximum relative fish abundance).

Friedman two-way ANOVA by ranked blocks (Zar, 1974) was used to evaluate monthly spatial differences among physical and chemical parameters, relative phytoplankton fluorescence, relative fish abundance, and zooplankton densities. Newman-Kuels multiple range test using standard errors appropriate for ranked blocks were

used to compare conditions between adjacent littoral and limnetic sampling sites. Relationships of zooplankton density with limnological parameters, relative phytoplankton fluorescence, and relative fish abundance were determined using Spearman's rank correlation. A Sorensen Similarity Index was used to determine percent similarity in species composition among stations.

Results

Physical and chemical conditions

Average (0 to 10 m) temperature, dissolved oxygen, and pH showed no spatial heterogeneity ($P > 0.05$ for each parameter). Thermal stratification developed during summer with the thermocline between 7 and 12 m. Summer epilimnetic temperature averaged $25.2\text{C} \pm 0.7$ (SD) and pH averaged 7.9 ± 0.4 (SD). Dissolved oxygen averaged $8.9 \text{ mg l}^{-1} \pm 0.9$ (SD) and at no time did the upper 10 m become anoxic at any station. Almost complete vertical mixing occurred during winter. Temperature (0–10 m) averaged $12.9\text{C} \pm 0.7$ (SD) and dissolved oxygen and pH averaged $9.2 \text{ mg l}^{-1} \pm 0.7$ (SD) and 7.7 ± 0.3 (SD), respectively.

Conductivity, however, varied significantly among stations ($P < 0.001$). ILVB had the highest summer conductivity averaging $1275 \mu\text{mhos cm}^{-1} \pm 79$ (SD). Littoral and limnetic MLVB stations, however, showed no statistical difference ($P > 0.20$) and summer values averaged $1043 \mu\text{mhos cm}^{-1} \pm 64$ (SD). Conductivity was uniform between littoral and limnetic BB sites ($P > 0.50$) and during summer averaged $985 \mu\text{mhos cm}^{-1} \pm 71$ (SD). During winter, conductivity was low at all sites and averaged $906 \mu\text{mhos cm}^{-1} \pm 44$ (SD).

Relative phytoplankton biomass and fish abundance

Phytoplankton fluorescence showed significant variation among sites ($P < 0.001$). Fluorescence was highest at ILVB and progressively declined

at MLVB sites and BB sites. However, adjacent littoral and limnetic sites at MLVB ($P > 0.20$) and at BB ($P > 0.20$) sites had similar values. Seasonal fluorescence varied with maxima occurring during late spring and summer (Table 1).

Relative fish abundance also showed significant spatial differences ($P < 0.001$). Adjacent littoral and limnetic MLVB ($P > 0.20$) and BB ($P > 0.50$) sites showed no difference. Fish were more abundant at ILVB and progressively declined at MLVB stations and BB stations (Table 2).

Zooplankton species composition

A diverse zooplankton community existed in Lake Mead. Species richness was greater for limnetic associated taxa (27 species) (Edmondson, 1959; Pennak, 1978) than littoral associated taxa (15 species). Littoral species comprised 2% or less of the total zooplankton density and occurred in about equal abundance between adjacent littoral and limnetic sites. Only during June did littoral zooplankton, primarily the rotifer *Trichocerca cylindrica*, increase; littoral taxa accounted for 40% of the total zooplankton density at ILVB and 10% and 11% at limnetic and littoral MLVB, respectively. No littoral species were found at BB sites during June.

Species composition generally was similar

Table 1. Relative phytoplankton fluorescence among littoral and limnetic sampling stations in Lake Mead. Values are actual fluorescence measurements in arbitrary units.

Month	ILVB	Litt MLVB	Limn MLVB	Litt BB	Limn BB
Jul.	34.2	8.0	5.0	0.5	0.7
Sept.	2.2	0.8	0.7	0.6	0.8
Nov.	0.9	1.1	1.0	0.9	1.0
Dec.	1.0	0.8	0.8	0.6	0.7
Jan.	0.9	0.8	0.7	0.7	0.8
Feb.	1.1	0.8	0.6	0.5	0.8
Mar.	3.0	1.2	1.0	0.8	0.7
Apr.	1.7	1.0	0.7	0.6	0.6
May	4.0	2.8	3.4	3.1	2.4
Jun.	-	1.9	3.0	0.8	1.7

Table 2. Relative fish abundance among littoral and limnetic sampling stations in Lake Mead. A one indicates no fish present and a 5 indicates the maximum relative abundance of fish recorded.

Month	ILVB	Litt MLVB	Limn MLVB	Litt BB	Limn BB
Jul.	4	4	5	2	2
Aug.	4	3	3	3	2
Sept.	4	3	2	2	2
Oct.	3	3	3	2	2
Nov.	5	3	1	2	2
Dec.	5	2	1	2	2
Jan.	5	2	2	2	2
Feb.	4	3	2	2	2
Mar.	4	2	3	2	2
Apr.	5	3	3	2	1
May	5	3	3	3	2
Jun.	-	3	3	3	2

among all littoral and limnetic sites (Table 3). The most common were the copepods *Diacyclops bicuspidatus thomasi*, *Diaptomus ashlandi*, and *Mesocyclops edax*; the cladocerans *Bosmina longirostris*, *Daphnia galeata mendotae*, and *D. pulex*; and the rotifers *Polyarthra* and *Synchaeta*.

Zooplankton abundance

Total average zooplankton densities throughout the water column are presented in Fig. 2. Seasonal patterns were, for the most part, similar between littoral and limnetic stations; peaks occurred during autumn and late winter and de-

Table 3. Sorensen Similarity Index showing the percent similarity in species composition among littoral and limnetic sampling stations in Lake Mead.

	ILVB	Limn MLVB	Litt MLVB	Limn BB	Litt BB
ILVB	100	73	85	86	79
Limn MLVB		100	78	79	79
Litt MLVB			100	78	81
Limn BB				100	87
Litt BB					100




Fig. 2. Monthly total weighted-average zooplankton density among littoral and limnetic sampling stations in Lake Mead.

clined in late spring and summer. Densities were greatest at ILVB. Adjacent littoral and limnetic MLVB stations significantly differed ($P < 0.05$) in total zooplankton density, however, both BB sites had a similar total zooplankton density ($P > 0.20$).

Table 3 shows results from Spearman's correlation of zooplankton species with biological, physical, and chemical factors. Few species correlated with phytoplankton fluorescence and fish abundance, however, more species significantly correlated with physical and chemical factors.

Most species showed significant differences in density among sampling stations ($P < 0.05$). Rotifers (mainly *Synchaeta* spp., *Polyarthra* spp., and, at times, *T. cylindrica*) accounted for a large portion of the zooplankton community during late summer and autumn at ILVB (Fig. 3). Rotifers were far less important at MLVB and BB and reached similar community dominance in adjacent littoral and limnetic sites. Copepods (mainly nauplii, copepodites, and to a lesser extent adult *Diaptomus ashlandi*) dominated during winter and most of spring at ILVB and throughout the year at littoral and limnetic MLVB and limnetic BB (Fig. 3). Cladocerans dominated only at littoral BB when *Bosmina longirostris* and *Daphnia galeata mendotae* increased in abundance during summer and early autumn (Fig. 3). However, densities of *Mesocyclops edax* ($P > 0.10$), *Diaptomus ashlandi* ($P > 0.995$), *D. reighardi* ($P > 0.25$), *Daphnia pulex* ($P > 0.10$), and *D. galeata mendotae*

($P > 0.90$) showed no significant difference among stations.

Figure 4 summarizes relative phytoplankton fluorescence, relative fish abundance, and total zooplankton densities averaged during the study. These factors were greatest at ILVB and progressively declined at MLVB and even further at BB.

Discussion

Zooplankton species composition in littoral and limnetic areas in Lake Mead was similar with few littoral taxa in either. Similar horizontal species distribution might result from several factors. The area of littoral zone is relatively small with little structural complexity and wind-generated currents can easily mix water between inshore and offshore areas. This may result in homogeneous physical and chemical environments and aid in the transport and mixing of zooplankton (George & Edwards, 1976; Hart, 1976; Kairesalo, 1980). The absence of rooted aquatic vegetation, however, appears to be a major contributor in reducing the number of littoral zooplankton species (Straskraba, 1964; Stolbunova & Stolbunov, 1981; Lemly & Dimmick, 1982a, b). Green (1986) suggested that vegetation is more important in increasing species diversity than merely location (i.e., inshore or offshore areas) because of greater habitat complexity than openwater. Williams (1982) found 24 littoral chydorid species in a lake dominated by littoral vegetation, and Quade (1969) found over 20 littoral cladoceran species amongst vegetation in each of several lakes. Lakes with extensively vegetated littoral zones have greater habitat heterogeneity contributing to greater species richness than lakes that lack vegetation (Pennak, 1966; Stolbunova & Stolbunov, 1981; Lemly & Dimmick, 1982a). Large, dense macrophyte stands also reduce horizontal water mixing and transport of plankton between littoral and limnetic zones (Kairesalo, 1980).

Although zooplankton species composition was horizontally similar, there was considerable variation among other environmental conditions. Significant difference in phytoplankton biomass




Fig. 3. Percent occurrence of major zooplankton groups (copepod, cladoceran, rotifer) at littoral and limnetic sampling stations in Lake Mead.

occurred horizontally from ILVB to BB, however, no difference was found between adjacent littoral and limnetic zones. ILVB was located near Las Vegas Wash inflow containing treated wastewater high in phosphorus that enhances phytoplankton biomass and growth (Baker & Paulson,

1981; Paulson & Baker, 1984). Diffusion of nutrients primarily due to a deep-water density current reduces growth and standing stock of algae in middle Las Vegas Bay and Boulder Basin (Baker & Paulson, 1981). Zooplankton abundances also were greater at ILVB and spatially

Table 4. Spearman's correlation coefficients of common zooplankton species abundance in Lake Mead with phytoplankton biomass, fish abundance, dissolved oxygen, temperature, pH, and conductivity. An asteric indicates a significant difference ($P < 0.05$).

Species ⁵	Phytopl	Fish	DO	Temp	pH	Conduct
naup	0.098	0.317*	0.274*	-0.146	0.279*	0.053
C cop	0.080	0.133	0.233	-0.361*	0.290*	-0.181
M cop	0.341*	0.341*	-0.299*	0.472*	0.506*	0.401*
D cop	-0.261	-0.114	0.354*	-0.797*	-0.048	-0.580*
Dbt	-0.054	0.077	0.269*	-0.729*	0.061	-0.452*
Me	0.229	0.201	-0.093	-0.131	0.392*	-0.064
Da	-0.169	-0.061	0.366*	-0.859*	-0.031	-0.671*
Dr	-0.148	-0.014	0.030	-0.413*	0.072	-0.288*
Ds	0.107	0.018	0.216	-0.668*	0.105	-0.545*
Bl	0.347*	0.321*	-0.361*	0.779*	0.196	0.694*
Dgm	0.166	0.272*	0.023	0.277*	0.423*	0.255*
Dp	-0.078	0.017	0.230	-0.737*	0.207	-0.550*
Rot	0.491*	0.409*	0.357*	0.153	0.167	0.239
Total	0.049	0.366*	0.329*	-0.105	0.263*	0.128

⁵ naup = nauplii; C cop = *Diatylops* copepodites; M cop = *Mesocyclops* copepodites; D cop = *Diatomus* copepodites; Dbt = *Diatylops bicuspidatus thomasi*; Me = *Mesocyclops edax*; Da = *Diatomus ashlandi*; Dr = *D. reighardi*; Ds = *D. siciloides*; Bl = *Bosmina longirostris*; Dgm = *Daphnia galeata mendotae*; Dp = *D. pulex*; Rot = rotifers; Total = total zooplankton abundance.

paralleled phytoplankton fluorescence (see Fig. 4). However, monthly values did not significantly correlate suggesting the uncoupling of links between grazers and phytoplankton indicating other factors showing stronger interactions to affect zooplankton dynamics. Wilde (1984), though, found a positive correlation in zooplankton abundance and phytoplankton biomass (measured as chlorophyll-*a*).

The dominant open-water planktivorous fish in Lake Mead is threadfin shad (*Dorosoma petenense*) (Allan & Roden, 1978). Planktivores in the littoral zone include bluegill sunfish (*Lepomis macrochirus*), green sunfish (*L. cyanellus*) and many other larval and juvenile fishes that become abundant in spring and summer (Allan & Roden, 1978). During our study, fish abundance paralleled total zooplankton abundance. Fish such as threadfin shad migrate from deep-water areas to Las Vegas Bay to spawn in spring and summer (Paulson & Espinosa, 1975; Allan & Roden, 1978). This suggests that these fish utilize areas of high food abundances that can support more fish and increase larval survival.

Zooplankton community structure differed monthly among sites indicating variability in fac-

tors influencing these patterns. For example, cladocerans dominated the littoral zone of BB in summer and autumn. *Bosmina longirostris* and *Daphnia galeata mendotae* densities totaled nearly $60\ l^{-1}$ in the littoral zone and less than $10\ l^{-1}$ in the limnetic zone (Sollberger, 1987). It is possible that greater food resources and production occurred in the littoral zone. Porter (1977) noted




Fig. 4. Annual average zooplankton density, relative phytoplankton fluorescence, and relative fish abundance among littoral and limnetic sampling stations in Lake Mead. Vertical lines represent standard deviation.

that in oligotrophic lakes phytoplankton production may be high enough to support a greater number of zooplankton than suggested by phytoplankton biomass. Wind-induced water currents also can transport zooplankton from offshore to inshore areas (George & Edwards, 1976). Kairesalo & Penttila (1990) found that *Bosmina longispina* has relatively low resistance to wind-induced water currents which may control its horizontal distribution.

Rotifers, which positively correlated with phytoplankton biomass, were most abundant at ILVB and dominated the zooplankton community during summer and autumn. Wilde (1984) also found that rotifer spatial pattern paralleled those of chlorophyll-*a* concentrations throughout Lake mead. Pace (1986) and Zankai (1989) reported that rotifer abundances corresponded to algal biomass and production rates and were most abundant in eutrophic areas of lakes with variable degrees of fertility. Since fish were more abundant at ILVB than other sites, selective predation may have eliminated large competitive zooplankton (such as *Daphnia*) lending to the dominance of rotifers (Brooks & Dodson, 1965; Hurlbert & Mulla, 1981; MacIsaac & Gilbert, 1989). Rotifer abundances, however, never greatly increased until October and November when densities reached about 308 l^{-1} and 63 l^{-1} , respectively (Sollberger, 1987). This increase might have resulted from a shift in larval fish diets to larger prey items (Wilde & Paulson, 1988) or from more favorable conditions occurring for rotifer production to offset predation losses (Hutchinson, 1967; Orcutt & Pace, 1984).

Many zooplankton species significantly correlated with temperature, DO, conductivity, and pH. Physical and chemical environments can influence zooplankton seasonal abundances (Hutchinson, 1967), but from this study it is difficult to determine the extent these factors constrain species abundances.

In summary, most zooplankton species showed differences in horizontal distribution between littoral and limnetic habitats. The lack of littoral vegetation and similar limnological conditions probably resulted in the dominance of limnetic

zooplankton in littoral areas and low species richness of littoral taxa. If the littoral zone contained dense weedbeds, then perhaps zooplankton species richness would be greater in Lake Mead. Our data suggests that other factors, particularly relative abundance of fish, greatly influence zooplankton horizontal abundances and percent species composition.

Acknowledgements

We greatly appreciate help from Lisa Heki during field sampling. Gene Wilde, Mike Meador, Bill Richardson, and Martin Cryer made valuable comments for improvements to the manuscript. Thanks also to Philip Dixon for suggesting appropriate statistical analyses. The drafting of Fig. 4 came from support of the United States Department of Energy and the Research Foundation of the University of Georgia, contract DE-AC-09-76SR00819.

References

- Allan, R. C. & D. L. Roden, 1978. Fish of Lakes Mead and Mohave. Nev. Dept. Wildl. Biol. Bull. No. 7. 105 pp.
- Baker, J. R. & L. J. Paulson, 1981. Influences of Las Vegas Wash density current on nutrient availability and phytoplankton growth in Lake Mead. pp. 1639-1647. In: H. G. Stefan (ed.), Symposium on surface water impoundments ASCE. June 2-5, 1980. Minneapolis, MN.
- Brooks, J. L. & S. I. Dodson, 1965. Predation, body size and competition of plankton. *Science* 150: 552-564.
- Cryer, M. & C. R. Townsend, 1988. Spatial distribution of zooplankton in a shallow eutrophic lake, with a discussion on its relation to fish predation. *J. Plankton Res.* 10: 487-501.
- DeBernardi, R., G. Giussani, E. L. Pedretti & T. Ruffoni, 1985. Population dynamics of pelagic cladocerans in three lakes with different trophy. *Verh. int. Ver. Limnol.* 22: 3035-3039.
- Edmondson, W. T. (ed.), 1959. Fresh water biology. John Wiley and Sons. New York, NY. 1248 pp.
- Gehrs, C. W., 1974. Horizontal distribution and abundance of *Diaptomus clavipes* Schacht in relation to *Potamogeton foliosus* in a pond and under experimental conditions. *Limnol. Oceanogr.* 19: 100-104.
- George, D. G. & R. W. Edwards, 1976. The effects of wind on the distribution of chlorophyll A on crustacean plank-

- ton in a shallow eutrophic reservoir. *J. appl. Ecol.* 13: 667-690.
- Green, J., 1986. Associations of zooplankton in six crater lakes in Arizona, Mexico and New Mexico. *J. Zool.* 208: 138-159.
- Haley, J. S., S. Leavitt, L. Paulson & D. H. Bacpler, 1987. Lake Mead cover enhancement project. Lake Mead Limnol. Res. Cent., Las Vegas, Nevada. Tech. Rept. No. 18. 117 pp.
- Hart, R. C., 1976. Horizontal distribution of the copepod *Pseudodiaptomus hessi* in subtropical Lake Sibaya. *Freshwat. Biol.* 8: 415-421.
- Heaney, S. I., 1978. Some observations on the *in vivo* fluorescence technique to determine chlorophyll-*a* in natural populations and cultures of freshwater phytoplankton. *Freshwat. Biol.* 8: 115-126.
- Hoffman, D. A. & A. R. Jonez, 1973. Lake Mead, a case history. pp. 220-233. In: W. C. Ackerman, G. C. White & E. B. Worthington (eds.), *Man-Made lakes: their problems and environmental effects*. Geographical Monograph Series No. 17.
- Hurlbert, S. H. & M. S. Mulla, 1981. Impacts of mosquitofish (*Gambusia affinis*) predation on plankton communities. *Hydrobiologia* 83: 125-151.
- Hutchinson, G. E., 1967. A treatise on limnology. Vol. II. Introduction to lake biology and the limnoplankton. John Wiley and Sons. New York, NY. 1115 pp.
- Jakobsen, P. J. & G. H. Johnsen, 1987. The influence of predation on horizontal distribution of zooplankton species. *Freshwat. Biol.* 17: 501-507.
- Kairesalo, T., 1980. Diurnal fluctuations within a littoral plankton community in oligotrophic Lake Paajarvi, southern Finland. *Freshwat. Biol.* 10: 533-537.
- Kairesalo, T. & S. Penttila, 1990. Effects of light and water flow on the spatial distribution of littoral *Bosmina longispina* LEYDIG (Cladocera). *Verh. int. Ver. Limnol.* 24: 682-687.
- Lemly, A. D. & J. F. Dimmick, 1982a. Structure and dynamics of zooplankton communities in the littoral zone of some North Carolina lakes. *Hydrobiologia* 88: 299-307.
- Lemly, A. D. & J. F. Dimmick, 1982b. Growth of young-of-the-year and yearling centrarchids in relation to zooplankton in the littoral zone of lakes. *Copeia* (1982): 305-321.
- MacIsaac, H. J. & J. J. Gilbert, 1989. Competition between rotifers and cladocerans of different body sizes. *Oecologia* 81: 295-301.
- McCauley, E. & J. Kalf, 1980. Empirical relationships between phytoplankton and zooplankton biomass in lakes. *Can. J. Fish. aquat. Sci.* 38: 458-463.
- Meyer, D. G., 1984. Habitat, shifting, feeding mode versatility, and alternate resource exploitation by herbivorous cladoceran zooplankton in a montane lake. pp. 309-345. In: D. G. Meyer & J. R. Strickler (eds.), *Trophic interactions within aquatic ecosystems*. Westview Press. Boulder, CO.
- Orcutt, J. D., Jr. & M. L. Pace, 1984. Seasonal dynamics of rotifers and crustacean zooplankton populations in a eutrophic, monomictic lake with a note on rotifer sampling techniques. *Hydrobiologia* 119: 73-80.
- Pace, M. L., 1986. An empirical analysis of zooplankton community size structure across lake trophic gradients. *Limnol. Oceanogr.* 31: 45-55.
- Paulson, L. J. & J. R. Baker, 1984. Interrelationships among nutrients, plankton and striped bass in Lake Mead. Lake Mead Limnol. Res. Cent., Las Vegas, Nevada, Tech. Rept. No. 11. 276 pp.
- Paulson, L. J. & F. A. Espinosa, 1975. Fish trapping: a method of evaluating fish species composition in limnetic areas of a reservoir. *Calif. Fish Game* 61: 209-214.
- Pennak, R. W., 1957. Species composition of limnetic zooplankton communities. *Limnol. Oceanogr.* 2: 222-232.
- Pennak, R. W., 1966. Structure of zooplankton in the littoral macrophyte zone of some Colorado lakes. *Trans. am. Microsc. Soc.* 85: 329-349.
- Pennak, R. W., 1978. Fresh-water invertebrates of the United States. John Wiley & Sons, New York, NY. 803 pp.
- Porter, K. G., 1977. The plant-animal interface in freshwater ecosystems. *Am. Sci.* 65: 155-170.
- Quade, H. W., 1969. Cladoceran faunas associated with aquatic macrophytes in some lakes in northeastern Minnesota. *Ecology* 50: 170-179.
- Siebeck, O., 1980. Optical orientation of pelagic crustaceans and its consequences in the pelagic and littoral zone. pp. 28-38. In: W. C. Kerfoot (ed.), *Evolution and ecology of zooplankton communities*. Univ. Press of New England. Hanover, NH.
- Smirnov, N. N., 1963. On inshore Cladocera of the Volga Water reservoirs. *Hydrobiologia* 21: 166-176.
- Smyly, W. J. P., 1952. The entomostraca of the weeds of a moorland pond. *J. anim. Ecol.* 2: 1-11.
- Sollberger, P. J., 1987. Comparison of littoral and limnetic zooplankton communities of Lake Mead. unpublished M.S. thesis. Univ. Nev., Las Vegas. 105 pp.
- Stolbonova, V. N. & A. K. Stolbonov, 1981. The natural life of the littoral zone of a reservoir and its effect on the pelagic zone (with reference to the bacterioplankton and zooplankton of Ivan'Kovo Reservoir). *Hydrobiol. J.* 16: 1-16.
- Straskraba, M., 1964. Contributions to the productivity of the littoral region of ponds. I. Quantitative study of the littoral zooplankton of rich vegetation of the backwater Labicko. *Hydrobiologia* 26: 421-443.
- Wetzel, R. G., 1983. *Limnology*. 2nd edition. Saunders. Philadelphia, Penn. 743 pp.
- Wilde, G. R., 1984. Seasonal and spatial heterogeneity in the limnetic zooplankton community of Lake Mead. unpublished M.S. thesis. Univ. Nev., Las Vegas. 95 pp.
- Wilde, G. R. & L. J. Paulson, 1988. Food habits of young-of-year largemouth bass in Lake Mead and Lake Mohave, Arizona-Nevada. *Great Basin Nat.* 48: 485-488.
- Wilde, G. R. & L. J. Paulson, 1989. Temporal and spatial variation in pelagic fish abundance in Lake Mead determined from echograms. *Calif. Fish Game* 75: 218-223.

Williams, J. B., 1982. Temporal and spatial patterns of abundances of the Chydoridae (Cladocera) in Lake Itasca, Minnesota. *Ecology* 63: 345-353.

Zankai, N. P., 1989. Horizontal distribution of rotifer plankton along a trophic gradient in Lake Balaton: changes of

community structure and abundances during the past 20 years. *Arch. Hydrobiol.* 115: 111-123.

Zar, J. H., 1974. *Biostatistical analysis*. Prentice-Hall, Inc., Englewood cliffs, N.J. 620 pp.