

Summer 2009

Synthesis completed of post-fire recovery of native perennials in the Mojave, Sonoran Deserts

Scott R. Abella

University of Nevada, Las Vegas, scott.abella@unlv.edu

Follow this and additional works at: https://digitalscholarship.unlv.edu/sea_fac_articles



Part of the [Environmental Health and Protection Commons](#), [Environmental Indicators and Impact Assessment Commons](#), and the [Other Environmental Sciences Commons](#)

Repository Citation

Abella, S. R. (2009). Synthesis completed of post-fire recovery of native perennials in the Mojave, Sonoran Deserts. *Mojave Applied Ecology Notes*, 2(2), https://digitalscholarship.unlv.edu/sea_fac_articles/374

This Newsletter is protected by copyright and/or related rights. It has been brought to you by Digital Scholarship@UNLV with permission from the rights-holder(s). You are free to use this Newsletter in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/or on the work itself.

This Newsletter has been accepted for inclusion in Public Policy and Leadership Faculty Publications by an authorized administrator of Digital Scholarship@UNLV. For more information, please contact digitalscholarship@unlv.edu.

Synthesis Completed of Post-Fire Recovery of Native Perennials in the Mojave, Sonoran Deserts

Scott Abella, Ph.D.

scott.abella@unlv.edu

Literature syntheses to develop status of knowledge reports are important to integrate and summarize the scattered scientific literature on a particular topic. The isolation and fragmentation of scientific literature on a topic is not necessarily a shortcoming of science. Rather, it is simply a consequence of having (1) research published in a diverse array of journals, (2) articles build on each other and therefore articles relevant to a particular topic can be published decades apart, and (3) funding virtually impossible to secure to do these periodic assessments of what we know and don't know (competitive science grants want researchers to jump into novel, new research, and land management agencies, although wishful for syntheses, cannot typically target funding out of their budgets for syntheses).

I was able to recently complete the third in a series of literature reviews (the first one was on burro grazing effects on Mojave vegetation, the second on Mojave revegetation). To synthesize post-fire recovery of perennial vegetation in the Mojave and Sonoran Deserts, I analyzed data systematically extracted from published literature to address several questions about post-fire recovery patterns.

Sprouting by desert perennials after being burned is generally limited but varies among species. For example, only 3-37% of *Larrea tridentata* sprouted compared to 64-86% of *Yucca schidigera* (the best-sprouting *Yucca* species examined). Four of five studies measuring recovery of perennial cover reported close relationships ($r^2 = 0.67-0.99$) between time since fire and cover (Fig. 1). In fact, three studies measuring the longest time since fire (≥ 37 years) found that cover had returned to within 10% cover of unburned areas within approximately 40 years. Conversely, post-fire species composition exhibited little convergence with unburned composition in five of six studies even 47 years after fire. *Sphaeralcea ambigua*, *Gutierrezia* spp., *Achnatherum speciosum*, *Encelia* spp., *Hymenoclea salsola*, and *Baileya multiradiata* had the highest abundances on burns relative to unburned areas, meaning that these species actually increased their relative abundance compared to other species after fire. Analyzing the literature as a whole suggested some generalities about recovery after fire (e.g., that perennial cover reestablishes faster than composition), but more work is required for improving specific knowledge about plant recovery among different fires, sites, species, and climates.

The citation of the full article published in *Journal of Arid Environments* follows, and a PDF is freely available from <http://faculty.unlv.edu/abellas2/>:

Abella, S.R. 2009. Post-fire plant recovery in the Mojave and Sonoran Deserts of western North America. *Journal of Arid Environments* 73:699-707.




Fig. 1. Example showing one of the five studies that measured the natural reestablishment of perennial plant cover following wildfire in the Mojave Desert. This graph shows the difference between cover on burned and unburned areas (with zero at the top meaning that burned cover has recovered to levels on adjacent unburned areas) as a function of time since wildfire. The study used a chronosequence approach where each point on the graph represents a different aged wildfire. The studies generally found that native perennial cover on burns had returned to unburned levels within 40 years after fire, even though the cover was supplied by different perennial species on burned compared to unburned areas.