

12-18-2018

mRNA, rRNA and DNA Quantitative Stable Isotope Probing with H218O Indicates Use of Old rRNA among Soil Thaumarchaeota

Katrina Papp
University of Nevada, Las Vegas, katerina.papp@unlv.edu

Bruce A. Hungate
Northern Arizona University

Egbert Schwartz
Northern Arizona University

Follow this and additional works at: https://digitalscholarship.unlv.edu/fac_articles



Part of the [Biology Commons](#)


Repository Citation

Papp, K., Hungate, B. A., Schwartz, E. (2018). mRNA, rRNA and DNA Quantitative Stable Isotope Probing with H218O Indicates Use of Old rRNA among Soil Thaumarchaeota. *Soil Biology and Biochemistry*, 130 159-166. Elsevier.

<http://dx.doi.org/10.1016/j.soilbio.2018.12.016>

This Article is protected by copyright and/or related rights. It has been brought to you by Digital Scholarship@UNLV with permission from the rights-holder(s). You are free to use this Article in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/or on the work itself.

This Article has been accepted for inclusion in Civil & Environmental Engineering and Construction Faculty Publications by an authorized administrator of Digital Scholarship@UNLV. For more information, please contact digitalscholarship@unlv.edu.



mRNA, rRNA and DNA quantitative stable isotope probing with H₂¹⁸O indicates use of old rRNA among soil Thaumarchaeota

Katerina Papp^{a,b,*}, Bruce A. Hungate^{a,b}, Egbert Schwartz^{a,b}

^a Center for Ecosystem Science and Society, Northern Arizona University, Flagstaff, AZ, USA

^b Department of Biological Sciences, Northern Arizona University, Flagstaff, AZ, USA



ARTICLE INFO

Keywords:

mRNA
rRNA and DNA quantitative stable isotope probing
Thaumarchaeota
amoA gene

ABSTRACT

rRNA is considered to be a short-lived molecule, indicative of cellular metabolic activity, whereas DNA is thought to turn over more slowly because living cells do not always grow and divide. To explore differences in the rates of synthesis of these nucleic acids, we used H₂¹⁸O quantitative stable isotope probing (qSIP) to measure the incorporation of ¹⁸O into 16S rRNA, the 16S rDNA, *amoA* mRNA and the *amoA* gene of soil Thaumarchaeota.

Incorporation of ¹⁸O into the thaumarchaeal *amoA* mRNA pool was faster than into the 16S rRNA pool, suggesting that Thaumarchaea were metabolically active while using rRNA molecules that were likely synthesized prior to H₂¹⁸O addition. Assimilation rates of ¹⁸O into 16S rDNA and *amoA* genes were similar, which was expected because both genes are present in the same thaumarchaeal genome. The Thaumarchaea had significantly higher rRNA to rDNA ratios than bacteria, though the ¹⁸O isotopic signature of thaumarchaeal rRNA was lower than that of bacterial rRNA, further suggesting preservation of old non-labeled rRNA. Through qSIP of soil with H₂¹⁸O, we showed that ¹⁸O incorporation into thaumarchaeal nucleic acids was generally low, indicating slower turnover rates compared to bacteria, and potentially suggesting thaumarchaeal capability for preservation and efficient reuse of biomolecules.

1. Introduction

It is axiomatic in biology that DNA replicates in proportion to cellular division (Marstorp and Witter, 1999). The production of ribosomal RNA (rRNA) is closely coupled to DNA replication because the dividing cells require high rates of protein synthesis (Ruimy et al., 1994), though new rRNA may also be synthesized to maintain a subset of proteins when a cell is not dividing (van Bodegom, 2007). Dormant cells generally have relatively low levels of rRNA because they synthesize fewer new proteins (Lennon and Jones, 2011). In pure cultures, rRNA can be relatively stable during exponential growth (Meselson et al., 1964), during stationary phase (Pirr et al., 2011), as intact 70S ribosomes in prokaryotes (Zundel et al., 2009), or as 100S dimers (Wada, 1998). Alternatively, rRNA may be degraded in starved cells or at the transition point into stationary phase, when it can be used as a source of nutrients (Deutscher, 2003). As a cell transitions from dividing to a maintenance state, rRNA declines while the DNA concentration remains constant (Deutscher, 2003; Hsu et al., 1994). Messenger RNA (mRNA) turns over faster than rRNA in a cell (Brenner et al., 1961), and its

instability is important for regulation of gene expression and rapid responses to environmental changes (Belasco and Higgins, 1988; Jain, 2002; Steege, 2000). Regulatory RNAs that control persistence of specific mRNAs are ubiquitous in microorganisms. While there are several mechanisms through which mRNA can be temporarily stabilized (Wong and Chang, 1986; Kushner, 2002; Hambræus et al., 2003; Lee et al., 2003; Baker and Condon, 2004), degradation of mRNA by ribonucleases occurs continuously (Deutscher, 2006). Decay and synthesis of mRNA molecules are carefully controlled to optimize cell survival, growth, performance or metabolic activity.

Some studies (Malik et al., 2015) have examined the turnover of nucleic acids in soil but generally these studies are rare. Characterizing nucleic acids synthesis in soil is challenging because soil has many microhabitats, which can provide highly diverse physico-chemical conditions with notable effects on microbial growth and metabolism. Unlike a cell culture, an environmental sample will contain both older and newly formed nucleic acids. Upon cell death in the environment, the DNA can degrade, be assimilated, or adsorb to surrounding particles (Dlott et al., 2015; Morrissey et al., 2015). Ribosomal RNA is likely less

* Corresponding author. 755, E. Flamingo Road, Las Vegas, NV, 89119, USA.

E-mail addresses: katerina.papp@dri.edu, katerina.papp@unlv.edu (K. Papp).

¹ Present address: Department of Civil and Environmental Engineering and Construction, University of Las Vegas, Las Vegas, NV, USA.

² Present address: Division of Hydrologic Sciences, Desert Research Institute, Las Vegas, NV, USA.

persistent than DNA in environmental samples. Rates of mRNA synthesis in environmental samples are likely higher than rates of DNA synthesis, because some protein synthesis is related to cellular maintenance and activity, not growth. For these reasons, turnover of mRNA in environmental samples is presumed to be rapid, more rapid than turnover of DNA, though empirical tests of this hypothesis are rare. Here, we used quantitative stable isotope probing (qSIP) experiments with H_2^{18}O (Hungate et al., 2015) to measure synthesis rates of microbial nucleic acids in environmental samples. We quantified the ^{18}O incorporation into nucleic acids of individual thaumarchaeal taxa and used it to estimate nucleic acid synthesis rates.

Ammonia oxidizing Thaumarchaeota (Könneke et al., 2005; Schleper et al., 2005; Venter et al., 2004) have a cosmopolitan distribution, are found in a wide range of habitats including aquatic (Kirchman et al., 2007; Teira et al., 2006; Yakimov et al., 2009) and terrestrial ecosystems (Treusch et al., 2005; Kemnitz et al., 2007), and have a complex chemolithoautotrophic physiology (Hatzenpichler, 2012). They have been found to outnumber ammonia-oxidizing bacteria (AOB) in many ecosystems (Leininger et al., 2006; Adair and Schwartz, 2008; Chen et al., 2008; Shen et al., 2008), suggesting an important role in the nitrogen cycle. They can be detected and quantified through measurements of 16S rRNA or *amoA* gene sequence abundance (Francis et al., 2005). By studying this group of organisms in a quantitative stable isotope probing experiment, it is possible to measure the turnover of DNA, rRNA and mRNA in individual microbial taxa. In the present study, we investigated transcription and replication of the 16S rDNA and the *amoA* gene of soil Thaumarchaeota through H_2^{18}O mRNA, rRNA and DNA-qSIP. Specifically, we measured the incorporation of ^{18}O into newly synthesized 16S rRNA and 16S rDNA as well as into the *amoA* gene and its transcripts. For Thaumarchaeota, *amoA* gene transcription indicates cellular activity related to ammonia oxidation, which is central to metabolism for this particular group of organisms. In contrast, 16S rRNA transcription is required for any prokaryotic cell to construct protein synthesis machinery. We hypothesized that replication of both genes would be highly correlated, but that transcription rates would differ.

2. Materials and methods

2.1. Sample collection

Soil samples were collected from three separate locations (N: 34° 57'21.879" W: 111° 45'14.859", N: 34° 57'21.289" W: 111° 26'45'14.189" and N: 34° 57'21.591" W: 111° 45'14.683") in a semi-arid ecosystem near Sedona, Arizona, USA in April 2014. The soil was characterized as a sandy loam (average pH of 6.95 ± 0.42 , soil moisture content of $4.11 \pm 0.24\%$ at the time of sampling, organic matter content of $13.7 \pm 1.7\%$, and concentrations of $\text{NO}_3\text{-N}$ and total P of 13.5 ± 7.0 and 14.3 ± 0.6 ppm respectively (AB-DPTA test, Soltanpour and Schwab, 1977)).

2.2. Incubation and ultracentrifugation

Two grams of overnight air-dried soil were incubated in triplicate with 400 μl of 95 atom % H_2^{18}O or with 400 μl of water with natural abundance ^{18}O for 1, 4 or 8 days at room temperature, resulting in a final atom % ^{18}O of approximately 80%. Total RNA and DNA were simultaneously extracted from 1 g of soil using RNA PowerSoil Total RNA Isolation Kit and DNA Elution Accessory Kit (MO BIO Laboratories, Carlsbad, CA). RNA extracts were digested with DNA-free DNase Treatment Removal Reagents (Ambion, Life Technologies, Grand Island, NY) to remove any traces of genomic DNA. Nucleic acids were quantified using the Qubit 2.0 Fluorimeter (Life Technologies, Grand Island, NY), separated by ultracentrifugation at $127,000 \times g$ in 3.3 ml Optiseal polyallomer tubes (Beckman Coulter, Fullerton, CA) on CsTFA (initial buoyant density of 1.79 g/ml, for RNA) or CsCl (initial buoyant

density of 1.73 g/ml, for DNA) density gradients. Samples were separated into approximately 120–150 μl fractions and purified as described in Papp et al. (2018a and 2018b). Briefly, RNA fractions were incubated with 200 μl of isopropanol and 100 ng glycogen at -20°C overnight, and DNA fractions were incubated with 200–300 μl of water, 300–450 μl of isopropanol and 100 ng of ultra-pure glycogen at room temperature overnight. Fractions were then precipitated with 70% ethanol and resuspended in 20–50 μl 1X TE buffer. RNA from purified fractions and from non-fractionated total RNA extracts was reverse transcribed into complementary DNA (cDNA) using Maxima H-Minus First Strand cDNA synthesis kit (Thermo Fisher Scientific, Waltham, MA) and random pentadecamer primers (Eurofins MWG Operon, Huntsville, AL). Extracted nucleic acids and corresponding fractions were stored at -80°C until further processing.

2.3. Sequencing

Sequencing of 16S rRNA gene fragments in cDNA and DNA samples was performed on the Illumina MiSeq platform at the Environmental Genetics and Genomics Facility (EnGGen) at Northern Arizona University using the 2×150 paired-end read chemistry and 515F (5' - GTGCCAGCMGCCGCGTAA - 3') and 806R (5' - GGACTACVSGGGT-ATCTAAT - 3') primers. Sequencing data were analyzed using a Quantitative Insights into Microbial Ecology (QIIME)-based (Caporaso et al., 2010) chained workflow (Krohn, 2016). A detailed description of sequencing data processing steps is available at <https://github.com/alk224/akutils-v1.2> and in Papp et al. (2018a and 2018b).

2.4. Arch-*amoA* qPCR

Standards for the archaeal *amoA* qPCR assay were made with DNA purified from qSIP fractions. The archaeal *amoA* gene was amplified in 20 μl reactions containing 2 μl of template and 7.32 μl of RNase-free water, 1X Phusion Mastermix (Water, 10X RedJuice (40% 1M Tris, pH 8.5/phenol red and 60% glycerol), 5X HF Buffer, 10 mM dNTPs, Phusion HSII polymerase), 1.5X MgCl_2 and 0.2 μM of a modified archaeal *amoA* primer: Arch-amoAFadapter (5' -caagcagaagacggcctacacagatSTAATGGTCTGGCTTAGACG-3') and Arch-amoARadapter (5' -caagcagaagacggcctacacagatGCGGCCATCCATCTGTATGT-3'). These primers were modified by attaching the 5'caagcagaagacggcctacacagat 3' adapter sequence to the 5' end of the archamoA primer to ensure that the *amoA* primer binding sites were not at the ends of the standard templates. Amplification was carried out in a 96-well plate with cycling conditions starting at 95°C for a 2-min denaturation step, followed by 30 cycles of 30 s at 95°C , 1 min at 55°C , and 1 min at 72°C . After 30 cycles there was a final extension for 3 min at 72°C followed by an indefinite hold at 4°C . A subset of 18 PCR products were randomly selected and visualized by gel electrophoresis. All 96 products were subsequently pooled into a 5 ml tube and purified with magnetic beads (0.1% carboxyl-modified Sera-Mag Magnetic Speed-beads, Thermo Fisher Scientific, Fremont, CA) in 18% PEG following a standard bead cleanup protocol with two ethanol washes (DeAngelis et al., 1995), and eluted into 200 μl Tris-Cl (pH 8.0). These PCR products were visualized by gel electrophoresis, and purified using Qiaquick Gel Extraction Kit (Qiagen, Valencia, CA, USA) according to manufacturer's instructions. Purified products were again visualized by gel-electrophoresis and quantified with Quant-it PicoGreen double-stranded DNA assay kit (Life Technologies, Grand Island, NY) on a BioTek HT Plate reader (BioTek, Winooski, VT). Serial dilutions, ranging from 10^8 to 10^2 gene copies per μl , were used as the standard curve for assay optimization. They were analyzed in triplicate using a quantitative PCR assay performed on a CFX384 Touch Real-Time PCR Detection System (BIORAD, Hercules, CA, USA). Each 10- μl reaction contained 1 μl of template (i.e. diluted standard), or no template control (NTC), 1X Phusion Mastermix, 1X EvaGreen dye (Biotium), 1.5X MgCl_2 and 0.2 μM of each primer: Arch-amoAF (5'

-STAATGGTCTGGCTTAGACG-3') and Arch-amoAR (5'-GCGGCCATCC ATCTGTATGT-3') (Francis et al., 2005). Cycling conditions started with a denaturation step at 95 °C for 2 min, followed by 40 cycles of 30 s at 95 °C, 30 s at 55 °C and 1 min at 72 °C. All runs to quantify the *amoA* gene or its transcripts in fractions were carried out on the CFX384 Detection System, fractions were quantified in triplicate, and standards in quintuplicate (10- μ l reactions) with the same components and cycling conditions as describe above.

2.5. Incorporation of ^{18}O into nucleic acids

Fraction density within a sample was corrected based on ^{18}O - IRMS data (Table S1) as described in detail in Papp et al. (2018a). Briefly, nucleic acids were quantified, diluted with a salmon sperm DNA solution to 200 μ g oxygen and pipetted into silver capsules (Costech Analytical Technologies Inc., Valencia, CA) in a tray, which was set on a heat block at 50 °C for water to evaporate. Capsules were then closed, weighted, and sent to the Stable Isotope Facility at University of California, Davis, for isotopic analysis. The corrected densities were used for further calculations. Calculations of (1) taxon-specific weighted average densities (WAD_i), (2) weighted averages density shifts (WAD_i shift) and (3) atom percent excess ^{18}O (APE ^{18}O) of nucleic acids, including *amoA* mRNA, *amoA* gene, 16S rRNA, and 16S rDNA, were carried out manually. Subsequently, they were also confirmed in R (R Core Team, 2014), using a code available at https://bitbucket.org/QuantitativeSIP/qsip_repo, and described in Hungate et al. (2015).

2.5.1. Weighted average density shifts of nucleic acids

All calculations and equations used for determining weighted average density (WAD) shifts of microbial nucleic acids (16S rRNA and rDNA) were previously published in Papp et al. (2018a). The same equations were used for determining the WAD shifts of 16S rRNA and rDNA for the thaumarchaeal taxa. We calculated WAD of thaumarchaeal *amoA* gene from a given taxon as the weighted average across density fractions. For a given fraction, we multiplied the relative abundance of that thaumarchaeal taxon (*rel.ab._{thaum}*) by the number of *amoA* copies detected via qPCR in that fraction (*amoA copy number*):

$$ABS\ amoA\ copy = rel.\ ab._{thaum} * amoA\ copy\ number \quad (1)$$

yielding an estimate of the absolute abundance of *amoA* copies for a given taxon in a given fraction (*ABS amoA copy*). The WAD for a given thaumarchaeal taxon (WAD_{thaum}) was then calculated as the sum of products of density times the absolute *amoA* copy number abundance in each fraction, divided by the total abundance across all fractions:

$$WAD_{thaum.} = \frac{\sum_{k=1}^K (ABS\ amoA\ copy * density)}{\sum_{k=1}^K (ABS\ amoA\ copy)} \quad (2)$$

where:

- k = fraction of a nucleic acid sample
- K = total number of fractions from a nucleic acid sample
- density* = density of a nucleic acid in each fraction

To obtain a WAD shift, we took the difference between WAD of nucleic acids from a sample incubated with H_2^{18}O and the paired sample incubated with water containing natural abundance levels of ^{18}O . The WAD shift reflects the increase in density of nucleic acids following incubation with isotopically labeled water. The procedure for estimating *amoA* mRNA was analogous but the *amoA* copy number was derived from fractions of complementary DNA (cDNA).

2.5.2. Atom percent excess ^{18}O of nucleic acids

All calculations and equations used for determining the atom percent excess ^{18}O (APE ^{18}O) values of microbial nucleic acids (16SrRNA and rDNA) were also previously published in Papp et al. (2018a) and in

Hungate et al. (2015). Some of these equations are provided in this section as well for clarity. Specifically, we converted WAD shifts of thaumarchaeal nucleic acids into atom percent excess ^{18}O (APE ^{18}O) values, which show the nucleic acids' isotopic enrichment above natural abundance of the ^{18}O isotope (0.2 atom%). We used equation (5) from Papp et al. (2018a) (equation (3) below) to convert WAD of thaumarchaeal RNA (16S rRNA or *amoA* mRNA) into atom percent ^{18}O values:

$$WAD = 0.0744 * \text{atom percent } ^{18}\text{O of RNA} + 1.7803 \quad (3)$$

This equation was obtained by using the known molecular weight of the rRNA molecule at different ^{18}O isotopic contents (natural abundance and 100% ^{18}O content), verified by empirical measurements of the density of RNA at natural abundance ^{18}O . We also used equation (6) from Papp et al. (2018a) (equation (4) below) to convert WAD of thaumarchaeal DNA (16S rDNA or *amoA* gene) into atom percent ^{18}O values:

$$WAD = 0.0644 * \text{atom percent } ^{18}\text{O of DNA} + 1.6946. \dots \dots \quad (4)$$

This equation was generated with results from a culture experiment of growing *E. coli* (strain HB101) at 5 different ^{18}O enrichment levels (natural abundance, 5, 25, 50 and 70% atom fraction ^{18}O). DNA was extracted from these cultures which was centrifuged to determine its density, and analyzed by IRMS to determine its ^{18}O signature. The equation is fully described and explained in Hungate et al. (2015). To calculate the atom percent excess ^{18}O (APE ^{18}O), the isotopic values of nucleic acids extracted from non-labeled incubations were subtracted from the isotopic values of nucleic acids extracted from the labeled treatment. We obtained three replicate APE ^{18}O values for each thaumarchaeal taxon at each time point.

2.6. rRNA to rDNA ratio calculations

Ratios of rRNA to rDNA were calculated for Thaumarchaea by dividing the relative abundance of RNA in each non-fractionated sample by the relative abundance of rDNA in the same sample. This was done for each replicate at each time point, yielding a total of 9 ratios per treatment. The replicate ratios were then averaged. There were two reasons for averaging the ratios: (1) there was no statistically significant difference over time ($p = 0.404$), and (2) we did not intend to investigate how the ratios changed over time.

2.7. Model of nucleic acid turnover

Details about turnover of microbial rRNA are provided in (Papp et al., 2018b). Briefly, Excel's Solver was used to determine the turnover rate of thaumarchaeal 16S rRNA and *amoA* mRNA that yielded the minimum sum of squared deviations from the observed data. We assumed that 50% of oxygen atoms in the thaumarchaeal nucleic acids originated from water and 50% from organic substrates (Chaney et al., 1972) and used findings from Ostle et al. (2003) and Yuan and Shen (1975) to obtain the starting turnover rates (20% per day, and 25% per hour respectively). After calculating the expected atom percent ^{18}O values of rRNA and mRNA at each of the reported rates (20% per day, and 25% per hour), we compared them to our observed values (on day 1, 4 and 8) and used Excel's Solver function to find the optimal turnover rate for each RNA type.

2.8. Statistical analyses

Statistical analyses were performed in SPSS (IBM SPSS Statistics 24), with statistical significance set to $\alpha = 0.05$. Normality was tested with the Shapiro-Wilk test, homogeneity of variances was tested with Levene's test, and sphericity was tested with Mauchly's test. Additionally, we used a two-way mixed ANOVA to test for a significant interaction

between $APE^{18}O$ of the different nucleic acid types (i.e. 16S rRNA, 16S rDNA, *amoA* mRNA, and *amoA* gene) and time. We also used a Spearman's rank order correlation test to assess the strength of correlations between (I) $APE^{18}O$ of 16S rDNA and *amoA* gene, (II) $APE^{18}O$ of *amoA* mRNA and 16S rRNA, (III) $APE^{18}O$ of 16S rRNA and 16S rDNA, and (IV) $APE^{18}O$ of 16S rDNA and 16S rRNA or *amoA* mRNA. Regressions were carried out using a model II regression analysis performed in R to test whether the slope of each correlation was significantly different from 1. We acknowledge that taxa within each replicate are not independent, and therefore, from the standpoint of inference from experimental design, the taxa are pseudoreplicates (taxa within replicate samples). However, the goal of our analyses was not to make inferences about differences between samples, or between groups of samples. The level of pseudoreplication therefore does not affect our conclusions. Combining all taxa within replicates would furthermore result in a significant loss of biological meaning caused by a considerable and particularly interesting variation among the individual taxa. Populations of different taxa are precisely the fundamental groups within fundamental biological and evolutionary units, where we would expect to see with great resolution a decoupling of DNA and mRNA synthesis - if it were to occur. Therefore, regressions in this manuscript show individual taxa within each replicate to illustrate general trends in nucleic acid synthesis with the highest resolution. The regression slopes were bootstrapped, which does not violate assumptions of the approach.

2.9. Accession numbers

All sequences have been deposited in the NCBI Sequence Read Archive (SRA) under accession numbers SAMN07960499 to SAMN07960874, SAMN07965143 to SAMN07965605, and SAMN07968111 to SAMN07968486. Data can directly be accessed at <https://www.ncbi.nlm.nih.gov/Traces/study/?acc=SRP123236>.

3. Results

3.1. Incorporation of ^{18}O into nucleic acids

The amount of ^{18}O incorporated into thaumarchaeal nucleic acids, expressed as atom percent excess ($APE^{18}O$), significantly differed over time and between 16S rRNA, the 16S rDNA, *amoA* mRNA, and the *amoA* gene ($p < 0.005$). *amoA* mRNA contained more ^{18}O than thaumarchaeal 16S rRNA ($p < 0.005$) and, with the exception of day 4, the *amoA* gene (Fig. 1, Figure S2, Supplemental material S2). As expected, the 16S rDNA and *amoA* genes had similar ^{18}O enrichment on each day ($p > 0.1$). The ^{18}O isotopic composition of 16S rRNA and *amoA* mRNA increased over time ($p < 0.05$, Fig. 1). The ^{18}O signature of




Fig. 1. Atom percent excess ($APE^{18}O$) of thaumarchaeal 16S rRNA (open diamond), *amoA* mRNA (filled diamond), 16S rDNA (open circle), and *amoA* gene (filled circle) at three incubation time points. Error bars show standard deviation of the mean of taxa ($n = 4$ taxa for 16S rRNA and *amoA* mRNA, and $n = 5$ taxa for 16S rDNA and *amoA* gene).




Fig. 2. Correlation between the $APE^{18}O$ of thaumarchaeal *amoA* gene and 16S rDNA (all individual time points are shown). Symbols represent individual thaumarchaeal taxa in replicate soil samples. Filled symbols show positive $APE^{18}O$ values, and open symbols show negative values, which were not included in the analysis. The green line represents a 1:1 line where the $APE^{18}O$ of both genes are equal. (For interpretation of the references to color in this figure legend, the reader is referred to the Web version of this article.)

thaumarchaeal 16S rDNA and *amoA* genes also increased over time with a slope significantly greater than zero for both genes (slope = 0.014 for 16S rDNA, and slope = 0.012 for *amoA* gene, Fig. 1, Supplemental material S2).

3.2. Correlations between isotopic signature of thaumarchaeal nucleic acids

Incorporation of ^{18}O into thaumarchaeal *amoA* genes was strongly, and positively, correlated to incorporation of ^{18}O into 16S rDNA ($\rho(37) = 0.953$, $p < 0.005$, Fig. 2). The equation describing the overall relationship between the two variables showed that the rates of incorporation of ^{18}O into both genes were similar (Fig. 2, Supplemental material S2). Correlation between the isotopic signature of thaumarchaeal *amoA* mRNA and 16S rRNA was also strong and positive ($\rho(34) = 0.860$, $p < 0.005$). The relationship, depicted in Fig. 3, showed that rate of assimilation of ^{18}O isotopes into *amoA* mRNA was greater than into 16S rRNA (Fig. 3, Supplemental material S2). Additionally, the rate of *amoA* mRNA turnover was higher (6.78% per day) compared to the rate of 16S rRNA turnover (4.61% per day). Lastly, incorporation of ^{18}O into 16S rDNA of Thaumarchaeota was correlated with incorporation of ^{18}O into their 16S rRNA (Figure S1A) and *amoA* mRNA (Figure S1B). Model II regression analysis showed that the slopes of both regressions were significantly smaller than one (slope = 0.40 for 16S rRNA vs 16S rDNA, and slope = 0.45 for *amoA* mRNA vs 16S




Fig. 3. Correlation between the $APE^{18}O$ of thaumarchaeal *amoA* mRNA and 16S rRNA using all individual incubation time points. Symbols show individual taxa in replicate soil samples. Positive $APE^{18}O$ values are represented by black symbols, and values of \sim zero by open symbols, the latter were not used in the analyses. The green line represents a 1:1 ratio, where the isotopic content of mRNA and rRNA is equal. (For interpretation of the references to color in this figure legend, the reader is referred to the Web version of this article.)




Fig. 4. Ratios of the relative abundance of rRNA over relative abundance of rDNA for major microbial groups in soil (A) and correlation between $\text{APE }^{18}\text{O}$ of 16S rDNA and 16S rRNA (B). The rRNA to rDNA ratios (A) represent means of three replicates and three time points ($n = 9$), error bars show standard deviation of the mean. Significant differences are indicated by *. The correlation (B) shows taxa at deepest taxonomic resolution, day 1 is represented by triangles, day 4 by circles and day 8 by squares; black symbols represent thaumarchaeal taxa, gray symbols represent bacterial taxa. Symbols are bootstrapped means; error bars are not shown for clarity.

rDNA; $p < 0.005$ for both, Supplemental material S2). The isotopic content of nucleic acids was generally lower on day one and increased with incubation time, although there was substantial variation among the replicates at each time point.

3.3. rRNA to rDNA ratios

As a phylum, Thaumarchaeota constituted ~5% of the rDNA and ~16% of the 16S rRNA sequencing libraries (Figure S3), which yielded a relatively high rRNA to rDNA ratio of 3.47. This ratio was significantly higher ($p < 0.005$) than rRNA to rDNA ratios for other bacterial phyla (with the exception of Armatimonadetes, which had an exceptionally high rRNA to rDNA ratio of 7.83). Most bacterial phyla had ratios below 1, with the exception of Proteobacteria (1.08) and Planctomycetes (1.24) (Fig. 4A, Supplemental material S2). Across all bacterial taxa, the correlation between the ^{18}O content of 16S rDNA and 16S rRNA was strong, positive, and significant ($\rho(345) = 0.859$, $p < 0.005$) (Fig. 4B, Supplemental material S2). Relative to bacterial phyla, thaumarchaeal taxa generally incorporated less ^{18}O into their nucleic acids (Fig. 4B, black symbols). The maximum $\text{APE }^{18}\text{O}$ was ~15% for thaumarchaeal 16S rRNA and ~19% for their 16S rDNA after 8 days of incubation with H_2^{18}O , while the maximum $\text{APE }^{18}\text{O}$ for bacterial 16S rRNA was ~43% and ~35% for their 16S rDNA at that time.

4. Discussion

4.1. ^{18}O isotopic content of thaumarchaeal nucleic acids

The natural abundance isotopic composition of newly synthesized nucleic acids within a cell, whether DNA, mRNA, or rRNA, should be similar because they all originate from interchangeable nucleotide pools. Ribonucleotide reductase catalyzes the conversion of ribonucleotides to deoxyribonucleotides (Herrick and Sclavi, 2007; Reichard, 1993), so therefore it is likely that both newly made ribonucleotides and deoxyribonucleotides have a similar ^{18}O content since ^{18}O water was added in very high amounts and fractionation processes could be ignored. Even if fractionation led to differences in natural abundance ^{18}O composition between types of nucleic acids, a large change in natural abundance (e.g., 50%) corresponds to a very small difference in tracer concentrations (± 0.01 atom % ^{18}O). The observed significant differences in the ^{18}O content of the bulk DNA, mRNA, or rRNA, following the addition of ^{18}O -water can thus only be explained by the fraction of the nucleotide pool composed of older, non-labeled molecules. Additionally, the ^{18}O content of a nucleic acid extract is also affected by the amount of newly synthesized molecules, the turnover rate, the synthesis strategy (*de-novo* vs recycling of nucleotides, discussed further), and the nucleic acid pool size.

^{18}O incorporation into *amoA* mRNA was greater than into the *amoA* gene at the beginning and at the end of our experiment, likely because mRNA synthesis rates were greater than DNA replication rates. This is consistent with the observations of Dumont et al. (2011) throughout a 4-day incubation with ^{13}C -labeled methane, whereby the isotopic signature of *pmoA* mRNA of methanotrophic communities was greater than the ^{13}C isotopic signature of *pmoA* genes. Since growth is not a prerequisite for metabolic activity (Blazewicz et al., 2013), greater ^{18}O content in microbial mRNA relative to the corresponding gene was expected.

Surprisingly, the overall isotopic enrichment of thaumarchaeal *amoA* mRNA was relatively low (~20% after 8 days of incubation with H_2^{18}O , turnover rate 6.78% per day). This suggested that turnover rate of mRNA may be slower in soil than in pure cultures on media with ideal growth conditions (such as 37°C temperature, Hambraeus et al., 2003) and high cellular activity. In these cultures, studies have measured short half-lives of bacterial mRNAs, ranging anywhere from ~30 s to ~30 min (Belasco and Higgins, 1988; Hambraeus et al., 2003; Rauhut and Klug, 1999; Wong and Chang, 1986). At such a fast turnover rate, we hypothesized that many oxygen atoms in *amoA* transcripts would be replaced with ^{18}O . We did not expect the mRNA to be 100% labeled with ^{18}O because microorganisms assimilate oxygen into ribonucleic acids from both water and other organic or inorganic compounds. In the case of Thaumarchaeota, oxygen is likely to come from water and potentially from fixed carbohydrates as they mostly use inorganic HCO_3^- as a carbon source (Hatzenpichler, 2012; Könneke et al., 2014). Oxygen in the fixed carbohydrates may partially come from water since water is used to produce Acetyl-CoA, a precursor for gluconeogenesis (Berg et al., 2010). Thus, newly synthesized nucleic acids of the Thaumarchaeota could potentially be more labeled with ^{18}O than nucleic acids of heterotrophs, but further research is needed to elucidate the rate of ^{18}O incorporation into newly fixed carbohydrates among autotrophs. Additionally, many ribonucleotides within a cell are likely recycled, and therefore new mRNA sequences will still contain many ^{16}O atoms, which likely contributed to the low ^{18}O content of thaumarchaeal *amoA* mRNA.

4.2. Chemolithoautotrophic metabolism

The thaumarchaeal *amoA* gene encodes a subunit of the ammonia monooxygenase enzyme (AMO) (Francis et al., 2005), which, in Thaumarchaeota, is responsible for energy generation (Hatzenpichler, 2012). AMO catalyzes the oxidation of ammonia to hydroxylamine,

which is the first and rate-limiting step of nitrification (Jia and Conrad, 2009; Leininger et al., 2006; Mertens et al., 2009; Nicol and Schleper, 2006). Preserving *amoA* transcripts for longer periods of time could increase the number of translated AMOs per *amoA* template and directly save energy associated with transcription. Ammonia oxidizing bacteria (AOB) are already known for their high *in vivo* stability of *amo* mRNA, proteins, and ribosomes (Koops et al., 2003; Bock and Wagner, 2006). Our study suggests that soil ammonia oxidizing archaea (AOA) exhibit a similar physiology. Incomplete mRNA labeling was also observed in German agricultural soils (Pratscher et al., 2011). In their study, up to 50% of the *amoA* transcripts had densities below 1.80 g/ml even after 12 weeks of incubation with $^{13}\text{CO}_2$ and $100 \mu\text{g N g}^{-1}$ dry weight soil. This indicated that the transcripts were not fully labeled with the heavy isotope (Pratscher et al., 2011). The detection of ^{18}O -labeled *amoA* mRNA in our study suggested that soil Thaumarchaea were active in nitrogen cycling (Gubry-Rangin et al., 2010; Huang et al., 2012; Stopnišek et al., 2010; Yao et al., 2011), but relating ^{18}O content of *amoA* mRNAs to ammonia oxidation rates was beyond the scope of this study.

4.3. Presence of old rRNA

We found that Thaumarchaea had higher ^{18}O content in their *amoA* mRNA than 16S rRNA, indicating that old thaumarchaeal rRNA was present in the cells (Deutscher, 2006, 2003; Jain, 2002). We expected the ^{18}O content of newly made mRNA and rRNA to be similar because both are transcribed by the same RNA polymerase (Ebright, 2000; Darst, 2001) and because, within a cell, ribonucleotides used for synthesis of mRNA and rRNA derive from the same nucleotide pool. Thus, if all RNA molecules had very high turnover rates and were recently synthesized, the *APE* ^{18}O of *amoA* and 16S ribosomal RNA should be very similar, which was not what we observed. The difference in *APE* ^{18}O of the two transcripts was most likely due to the presence of a greater amount of older non-labeled rRNA than mRNA. This suggests that mRNA turns over faster than rRNA. The latter, more stable, can thus be reused for maintenance and reduce cell's overall energy expenditure. Ribosome storage has been previously documented by, for example, Wada (1998) who observed formation of 100S dimers, which was proposed to be a mechanism for excess ribosome storage during the stationary phase of growth. Large amounts of stored ribosomes have also been observed in dormant cyanobacteria (Sukenik et al., 2012). Some thaumarchaeal cells may have been in stationary phase and have contained stored ribosomes, thus contributing to overall low isotope content measurements. Yet we observed that the nucleic acids of all taxa were enriched in ^{18}O . Such nucleic acid labeling indicated that at least some individuals within a population were metabolically active and growing.

4.4. Genome replication

As expected, the 16S rDNA and *amoA* genes had very similar isotopic contents. Both genes reside on a single chromosome within a thaumarchaeal cell, and as the cell divided, both became similarly enriched. Our finding of a strong correlation between the *APE* ^{18}O of the 16S rDNA and *amoA* genes with a slope of 1 confirmed that genome replication was uniform, resulting in all genes being labeled with isotopic tracers at the same rate. It also demonstrated that qSIP can measure the rate of genome replication, thus providing a unique opportunity to measure microbial growth rates *in situ*.

Our experiment simulated a rain event. Studies have shown that microbial respiration and growth rates increase after rewetting (Bloem et al., 1992; Iovieno and Bååth, 2008; Pesaro et al., 2004), but the total microbial biomass can decrease (Fierer and Schimel, 2002; Gordon et al., 2008). In H_2^{18}O stable isotope studies, no major changes in the absolute size of bacterial or archaeal populations were found after rewetting because growth was counter-balanced by death (Angel and

Conrad, 2013; Blazewicz et al., 2014; Koch et al., 2018). Water addition may have stimulated metabolic activity, including growth, of taxa that might otherwise have stayed dormant in our soil. Water addition increased the soil moisture content from 4% to 20% approximately. Nevertheless, since even desert ecosystems receive precipitation, our experiment remains representative of naturally occurring events and provides valuable information about microbial dynamics following these events.

4.5. Metabolic activity

RNA to DNA ratios are often used as an indicator of microbial metabolic activity. The assumption is that the higher the ratio the higher the activity level because more ribosomes are needed to sustain rapid metabolism. We were thus interested to see how well the ratios compared to qSIP. Through qSIP, we found that Thaumarchaeota incorporated low amounts of ^{18}O into their rRNA, suggesting that synthesis of new rRNA molecules was rather low compared to bacterial taxa. However, we measured high rRNA to rDNA ratios of the soil Thaumarchaeota, which is traditionally interpreted as an indication of high metabolic activity (Roszak and Colwell, 1987; DeAngelis et al., 2010; Baldrian et al., 2012; Dlott et al., 2015). This was in disagreement with result from qSIP but can be explained by the thaumarchaeal capability to preserve and use older ribosomes. The Thaumarchaeota were relatively abundant in the rRNA library but the level of ^{18}O incorporation into their rRNA was low. These two findings suggested that a substantial fraction of their rRNA was synthesized prior to H_2^{18}O addition and was thus at least 8 days old. Our results suggest that ribosomal RNA can be relatively long lived in the environment, whether used by metabolically active bacteria and archaea or perhaps intact and preserved but no longer part of living cells.

Declarations of interest

None.

Acknowledgements

This research was supported by award 1142096 from the National Science Foundation, division of solar programs, NSF DEB-1241094, the Department of Energy's Biological Systems Science Division, Program in Genomic Science (DE-SC0010579, and DE-SC0016207), and by the IGERT Fellowship. Additionally, the authors would like to thank Paul Dijkstra, Matthew Bowker, Rebecca Mau, Michaela Hayer, Ben Koch, and Lela Andrews from Northern Arizona University and Joshua Sackett from Desert Research Institute for their help and suggestions. The authors declare no conflict of interest.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.soilbio.2018.12.016>.

References

- Adair, K.L., Schwartz, E., 2008. Evidence that ammonia-oxidizing archaea are more abundant than ammonia-oxidizing bacteria in semiarid soils of northern Arizona, USA. *Microbial Ecology* 56, 420–426.
- Angel, R., Conrad, R., 2013. Elucidating the microbial resuscitation cascade in biological soil crusts following a simulated rain event. *Environmental Microbiology* 15, 2799–2815.
- Baker, K.E., Condon, C., 2004. Under the Tucson sun: a meeting in the desert on mRNA decay. *RNA* 10, 1680–1691.
- Baldrian, P., Kolařík, M., Štursová, M., Kopecký, J., Valášková, V., Větrovský, T., Žifčáková, L., Šnajdr, J., Rídl, J., Vlček, Č., Voříšková, J., 2012. Active and total microbial communities in forest soil are largely different and highly stratified during decomposition. *The ISME Journal* 6, 248–258.
- Belasco, J.G., Higgins, C.F., 1988. Mechanisms of mRNA decay in bacteria: a perspective. *Gene* 72, 15–23.

- Berg, I.A., Kockelkorn, D., Ramos-Vera, W.H., Say, R.F., Zarzycki, J., Hügler, M., Alber, B.E., Fuchs, G., 2010. Autotrophic carbon fixation in archaea. *Nature Reviews Microbiology* 8, 447–460.
- Blazewicz, S.J., Barnard, R.L., Daly, R.A., Firestone, M.K., 2013. Evaluating rRNA as an indicator of microbial activity in environmental communities: limitations and uses. *The ISME Journal* 7, 2061–2068.
- Blazewicz, S.J., Schwartz, E., Firestone, M.K., 2014. Growth and death of bacteria and fungi underlie rainfall-induced carbon dioxide pulses from seasonally dried soil. *Ecology* 95, 1162–1172.
- Bloem, J., de Ruyter, P.C., Koopman, G.J., Lebbink, G., Brussaard, L., 1992. Microbial numbers and activity in dried and rewetted arable soil under integrated and conventional management. *Soil Biology and Biochemistry* 24, 655–665.
- Bock, E., Wagner, M., 2006. Oxidation of inorganic nitrogen compounds as an energy source. *Prokaryotes* 2, 457–495.
- Brenner, S., Jacob, F., Meselson, M., 1961. An unstable intermediate carrying information from genes to ribosomes for protein synthesis. *Nature* 190, 576–581.
- Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer, N., Peña, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley, S.T., Knights, D., Koenig, J.E., Ley, R.E., Lozupone, C.A., McDonald, D., Muegge, B.D., Pirrung, M., Reeder, J., Sevinsky, J.R., Turnbaugh, P.J., Walters, W.A., Widmann, J., Yatsunenko, T., Zaneveld, J., Knight, R., 2010. QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* 7, 335–336.
- Chaney, S.G., Duffy, J.J., Boyer, P.D., 1972. Patterns of oxygen interchange between water, substrates, and phosphate compounds of *Escherichia coli* and *Bacillus subtilis*. *Journal of Biological Chemistry* 247, 2145–2150.
- Chen, X.P., Zhu, Y.G., Xia, Y., Shen, J.P., He, J.Z., 2008. Ammonia-oxidizing archaea: important players in paddy rhizosphere soil? *Environmental Microbiology* 10, 1978–1987.
- Darst, S.A., 2001. Bacterial RNA polymerase. *Current Opinion in Structural Biology* 11, 155–162.
- DeAngelis, M.M., Wang, D.G., Hawkins, T.L., 1995. Solid-phase reversible immobilization for the isolation of PCR products. *Nucleic Acids Research* 23, 4742–4743.
- DeAngelis, K.M., Silver, W.L., Thompson, A.W., Firestone, M.K., 2010. Microbial communities acclimate to recurring changes in soil redox potential status. *Environmental Microbiology* 12, 3137–3149.
- Deutscher, M.P., 2003. Degradation of stable RNA in bacteria. *Journal of Biological Chemistry* 278, 45041–45044.
- Deutscher, M.P., 2006. Degradation of RNA in bacteria: comparison of mRNA and stable RNA. *Nucleic Acids Research* 34, 659–666.
- Dlott, G., Maul, J.E., Buyer, J., Yarwood, S., 2015. Microbial rRNA:rRNA gene ratios may be unexpectedly low due to extracellular DNA preservation in soils. *Journal of Microbiological Methods* 115, 112–120.
- Dumont, M.G., Pommerenke, B., Casper, P., Conrad, R., 2011. DNA-, rRNA- and mRNA-based stable isotope probing of aerobic methanotrophs in lake sediment. *Environmental Microbiology* 13, 1153–1167.
- Ebright, R.H., 2000. RNA polymerase: structural similarities between bacterial RNA polymerase and eukaryotic RNA polymerase II. *Journal of Molecular Biology* 304, 687–698.
- Fierer, N., Schimel, J.P., 2002. Effects of drying-rewetting frequency on soil carbon and nitrogen transformations. *Soil Biology and Biochemistry* 34, 777–787.
- Francis, C.A., Roberts, K.J., Beman, J.M., Santoro, A.E., Oakley, B.B., 2005. Ubiquity and diversity of ammonia-oxidizing archaea in water columns and sediments of the ocean. *Proceedings of the National Academy of Sciences of the United States of America* 102, 14683–14688.
- Gordon, H., Haygarth, P.M., Bardgett, R.D., 2008. Drying and rewetting effects on soil microbial community composition and nutrient leaching. *Soil Biology and Biochemistry* 40, 302–311.
- Gubry-Rangin, C., Nicol, G.W., Prosser, J.I., 2010. Archaea rather than bacteria control nitrification in two agricultural acidic soils. *FEMS Microbiology Ecology* 74, 566–574.
- Hambraeus, G., Von Wachenfeldt, C., Hederstedt, L., 2003. Genome-wide survey of mRNA half-lives in *Bacillus subtilis* identifies extremely stable mRNAs. *Molecular Genetics and Genomics* 269, 706–714.
- Hatzenpichler, R., 2012. Diversity, physiology, and niche differentiation of ammonia-oxidizing archaea. *Applied and Environmental Microbiology* 78, 7501–7510.
- Herrick, J., Scavi, B., 2007. Ribonucleotide reductase and the regulation of DNA replication: an old story and an ancient heritage. *Molecular Microbiology* 63, 22–34.
- Hsu, D., Shih, L.M., Zee, Y.C., 1994. Degradation of rRNA in *Salmonella* strains: a novel mechanism to regulate the concentrations of rRNA and ribosomes. *Journal of Bacteriology* 176, 4761–4765.
- Huang, R., Wu, Y., Zhang, J., Zhong, W., Jia, Z., Cai, Z., 2012. Nitrification activity and putative ammonia-oxidizing archaea in acidic red soils. *Journal of Soils and Sediments* 12, 420–428.
- Hungate, B.A., Mau, R.L., Schwartz, E., Caporaso, J.G., Dijkstra, P., van Gestel, N., 2015. Quantitative Microbial Ecology through Stable Isotope Probing 81, 7570–7581.
- IBM SPSS Statistics for Windows, Version 24.0. IBM Corp., Armonk, NY.
- Iovieno, P., Bååth, E., 2008. Effect of drying and rewetting on bacterial growth rates in soil. *FEMS Microbiology Ecology* 65, 400–407.
- Jain, C., 2002. Degradation of mRNA in *Escherichia coli*. *IUBMB Life* 54, 315–321.
- Jia, Z., Conrad, R., 2009. Bacteria rather than Archaea dominate microbial ammonia oxidation in an agricultural soil. *Environmental Microbiology* 11, 1658–1671.
- Kemnitz, D., Kolb, S., Conrad, R., 2007. High abundance of Crenarchaeota in a temperate acidic forest soil. *FEMS Microbiology Ecology* 60, 442–448.
- Kirchman, D.L., Elifantz, H., Dittel, A.I., Malmstrom, R.R., Cottrell, M.T., 2007. Standing stocks and activity of archaea and bacteria in the western Arctic ocean. *Limnology & Oceanography* 52, 495–507.
- Koch, B.J., McHugh, T.A., Hayer, M., Schwartz, E., Blazewicz, S.J., Dijkstra, P., van Gestel, N., Marks, J.C., Mau, R.L., Morrissey, E.M., Pett-Ridge, J., Hungate, B.A., 2018. Estimating taxon-specific population dynamics in diverse microbial communities. *Ecosphere* 9, 1–15.
- Könneke, M., Bernhard, A.E., de la Torre, J.R., Walker, C.B., Waterbury, J.B., Stahl, D.A., 2005. Isolation of an autotrophic ammonia-oxidizing marine archaeon. *Nature* 437, 543–546.
- Könneke, M., Schubert, D.M., Brown, P.C., Hugler, M., Standfest, S., Schwander, T., von Borzyskowski, L.S., Erb, T.J., Stahl, D.A., Berg, I.A., 2014. Ammonia-oxidizing archaea use the most energy-efficient aerobic pathway for CO₂ fixation. *Proceedings of the National Academy of Sciences* 111, 8239–8244.
- Koops, H.P., Purkhold, U., Pommerening-Röser, A., Timmermann, G., Wagner, M., 2003. The lithoautotrophic ammonia-oxidizing bacteria. In: Dworkin, D., Falkow, S., Rosenberg, E., Schleifer, K.H., Stackebrandt, E. (Eds.), *The Prokaryotes: an Evolving Electronic Resource for the Microbiological Community*, third ed. Springer-Verlag, NY, USA release 3.13.
- Krohn, A., 2016. akutils-v12: Facilitating Analyses of Microbial Communities through QIIME. pp. 5281 Zenodo10.
- Kushner, S., 2002. mRNA decay in *Escherichia coli* comes of age. *Journal of Bacteriology* 184, 4658–4665.
- Lee, K., Zhan, X., Gao, J., Qiu, J., Feng, Y., Meganathan, R., Cohen, S.N., Georgiou, G., 2003. RraA: a protein inhibitor of RNase E activity that globally modulates RNA abundance in *E. coli*. *Cell* 114, 623–634.
- Leininger, S., Urlich, T., Schloter, M., Schwark, L., Qi, J., Nicol, G.W., Prosser, J.I., Schuster, S.C., Schleper, C., 2006. Archaea predominate among ammonia-oxidizing prokaryotes in soils. *Nature* 442, 806–809.
- Lennon, J.T., Jones, S.E., 2011. Microbial seed banks: the ecological and evolutionary implications of dormancy. *Nature Reviews Microbiology* 9, 119–130.
- Malik, A.A., Dannert, H., Griffiths, R.I., Thomson, B.C., Gleixner, G., 2015. Rhizosphere bacterial carbon turnover is higher in nucleic acids than membrane lipids: implications for understanding soil carbon cycling. *Frontiers in Microbiology* 6, 1–9.
- Marstorp, H., Witter, E., 1999. Extractable dsDNA and product formation as measures of microbial growth in soil upon substrate addition. *Soil Biology and Biochemistry* 31, 1443–1453.
- Mertens, J., Broos, K., Wakelin, S.A., Kowalchuk, G.A., Springael, D., Smolders, E., 2009. Bacteria, not archaea, restore nitrification in a zinc-contaminated soil. *The ISME Journal* 3, 916–923.
- Meselson, M., Nomura, M., Brenner, S., Davern, C., Schlessinger, D., 1964. Conservation of ribosomes during bacterial growth. *Journal of Molecular Biology* 9, 696–711.
- Morrissey, E.M., McHugh, T.A., Preteska, L., Hayer, M., Dijkstra, P., Hungate, B.A., Schwartz, E., 2015. Dynamics of extracellular DNA decomposition and bacterial community composition in soil. *Soil Biology and Biochemistry* 86, 42–49.
- Nicol, G.W., Schleper, C., 2006. Ammonia-oxidizing Crenarchaeota: important players in the nitrogen cycle? *Trends in Microbiology* 14, 207–212.
- Ostle, N., Whiteley, A.S., Bailey, M.J., Sleep, D., Ineson, P., Manfield, M., 2003. Active microbial RNA turnover in a grassland soil estimated using a ¹³C₂O spike. *Soil Biology and Biochemistry* 35, 877–885.
- Papp, K., Hungate, B.A., Schwartz, E., 2018a. Microbial rRNA synthesis and growth compared through quantitative stable isotope probing with H₂¹⁸O. *Applied and Environmental Microbiology* 84, e02441-17.
- Papp, K., Mau, R.L., Hayer, M., Koch, B.J., Hungate, B.A., Schwartz, E., 2018b. Quantitative stable isotope probing with H₂¹⁸O reveals that most bacterial taxa in soil synthesize new ribosomal RNA. *The ISME Journal* 18–20.
- Pesaro, M., Nicollier, G., Zeyer, J., Widmer, F., 2004. Impact of soil drying-rewetting stress on microbial communities and activities and on degradation of two crop protection products. *Applied and Environmental Microbiology* 70, 2577–2587.
- Piir, K., Paier, A., Liiv, A., Tenson, T., Mai Väli, U., 2011. Ribosome degradation in growing bacteria. *EMBO Reports* 12, 458–462.
- Pratscher, J., Dumont, M.G., Conrad, R., 2011. Ammonia oxidation coupled to CO₂ fixation by archaea and bacteria in an agricultural soil. *Proceedings of the National Academy of Sciences of the United States of America* 108, 4170–4175.
- R Core Team, 2014. R: a Language and Environment for Statistical Computing (R Version 3.2.1). (Vienna, Austria).
- Rauhut, R., Klug, G., 1999. mRNA degradation in bacteria. *FEMS Microbiology Reviews* 23, 353–370.
- Reichard, P., 1993. From RNA to DNA, why so many ribonucleotide reductases? *Science* 260, 1773–1777.
- Roszak, D.B., Colwell, R.R., 1987. Survival strategies of bacteria in the natural environment. *Microbiological Reviews* 51, 365–379.
- Ruimy, R., Breittmayer, V., Biovin, V., Christen, R., 1994. Assessment of the state of activity of individual bacterial cells by hybridization with a ribosomal RNA targeted fluorescently labelled oligonucleotide probe. *FEMS Microbiology Ecology* 15, 207–213.
- Schleper, C., Jurgens, G., Jonuscheit, M., 2005. Genomic studies of uncultivated archaea. *Nature Reviews Microbiology* 3, 479–488.
- Shen, J.P., Zhang, L.M., Zhu, Y.G., Zhang, J.B., He, J.Z., 2008. Abundance and composition of ammonia-oxidizing bacteria and ammonia-oxidizing archaea communities of an alkaline sandy loam. *Environmental Microbiology* 10, 1601–1611.
- Soltanpour, P.N., Schwab, A.P., 1977. A new soil test for simultaneous extraction of macro- and micro-nutrients in alkaline soils. *Communications in Soil Science and Plant Analysis* 8, 195–207.
- Steege, D.A., 2000. Emerging features of mRNA decay in bacteria. *RNA* 6, 1079–1090.
- Stopnišek, N., Gubry-Rangin, C., Höfferle, Š., Nicol, G.W., Mandič-Mulec, I., Prosser, J.I., 2010. Thaumarchaeal ammonia oxidation in an acidic forest peat soil is not influenced by ammonium amendment. *Applied and Environmental Microbiology* 76, 7626–7634.

- Sukenik, A., Kaplan-Levy, R.N., Welch, J.M., Post, A.F., 2012. Massive multiplication of genome and ribosomes in dormant cells (akinetes) of *Aphanizomenon ovalisporum* (Cyanobacteria). *The ISME Journal* 6, 670–679.
- Teira, E., Lebaron, P., van Aken, H., Herndl, G.J., 2006. Distribution and activity of bacteria and archaea in the deep water masses of the North Atlantic. 51, 2131–2144.
- Treusch, A.H., Leininger, S., Kietzin, A., Schuster, S.C., Klenk, H.P., Schleper, C., 2005. Novel genes for nitrite reductase and Amo-related proteins indicate a role of uncultivated mesophilic crenarchaeota in nitrogen cycling. *Environmental Microbiology* 7, 1985–1995.
- van Bodegom, P., 2007. Microbial maintenance: a critical review on its quantification. *Microbial Ecology* 53, 513–523.
- Venter, J.C., Remington, K., Heidelberg, J.F., Halpern, A.L., Rusch, D., Eisen, J.A., Wu, D., Paulsen, I., Nelson, K.E., Nelson, W., Fouts, D.E., Levy, S., Knap, A.H., Lomas, M.W., Nealson, K., White, O., Peterson, J., Hoffman, J., Parsons, R., Baden-Tillson, H., Pfannkoch, C., Rogers, Y.H., Smith, H.O., 2004. Environmental genome shotgun sequencing of the Sargasso Sea. *Science* 304, 66–74.
- Wada, A., 1998. Growth phase coupled modulation of *Escherichia coli* ribosomes. *Genes to Cells* 3, 203–208.
- Wong, H.C., Chang, S., 1986. Identification of a positive retroregulator that stabilizes mRNAs in bacteria. *Proceedings of the National Academy of Sciences of the United States of America* 83, 3233–3237.
- Yakimov, M.M., La Cono, V., Denaro, R., 2009. A first insight into the occurrence and expression of functional *amoA* and *accA* genes of autotrophic and ammonia-oxidizing bathypelagic Crenarchaeota of Tyrrhenian Sea. *Deep-Sea Research Part II Topical Studies in Oceanography* 56, 748–754.
- Yao, H., Gao, Y., Nicol, G.W., Campbell, C.D., Prosser, J.I., Zhang, L., Han, W., Singh, B.K., 2011. Links between ammonia oxidizer community structure, abundance, and nitrification potential in acidic soils. *Applied and Environmental Microbiology* 77, 4618–4625.
- Yuan, D., Shen, V., 1975. Stability of ribosomal and transfer ribonucleic acid in *Escherichia coli* B/r after treatment with ethylenedinitrotetraacetic acid and rifampicin. *Journal of Bacteriology* 122, 425–435.
- Zundel, M.A., Basturea, G.N., Deutscher, M.P., 2009. Initiation of ribosome degradation during starvation in *Escherichia coli*. *RNA* 15, 977–983.