



Journal of Health Disparities Research and Practice
Volume 9, Special Edition 1, Summer 2016, pp. 128-129
© 2011 Center for Health Disparities Research
School of Community Health Sciences
University of Nevada, Las Vegas

Localization of Fulicin-like Immunoreactivity in the Central Nervous System and Periphery of *Biomphalaria glabrata*, an Intermediate Host for Schistosomiasis

Anthony Hernandez-Vasquez
M.W. Miller, University of Puerto Rico
S. Rolon-Martinez, University of Puerto Rico
L.O. Vaasjo, University of Puerto Rico
M.B. Rodriguez, University of Puerto Rico
Coordinating Center: University of Nevada Las Vegas

ABSTRACT

An estimate of about ten percent of the population worldwide live at risk of contracting the parasitic disease schistosomiasis, or “snail fever”. The digenetic trematode worm species *Schistosoma mansoni* that is responsible for causing the most common form of intestinal schistosomiasis requires the freshwater snail *Biomphalaria glabrata* to serve as its primary intermediate host, where it multiplies and develops into its cercarial form that is infectious to humans. Parasitic castration and parasitic gigantism are among some of the profound behavioral changes that the infection of *B. glabrata* by *S. mansoni* is known to cause. For this reason, a neural transcriptomics approach was undertaken to determine precursor prohormones that could encode neuropeptides implicated in *Biomphalaria* reproductive and feeding behaviors. A transcript (1616 nucleotides) was found to encode a putative precursor polypeptide (316 aminoacids) that could give way to the neuropeptide fulicin (Phe–D-Asn-Glu-Phe-Val-NH₂; Ohta et al. 1991; Yasuma Kamatani et al. 1995) and five additional related peptides. For this investigation, affinity purified polyclonal antibodies (rabbit) were developed against the anticipated fulicin neuropeptide. Fulicin-like immunoreactivity was observed throughout the central nervous system (CNS) with distinct neurons and clusters on the ventral and dorsal surfaces, as well as in peripheral tissues. Fulicin-like cells of both large and small diameter were present on the dorsal and ventral surfaces of the buccal ganglion. In addition dispersed clusters of small diameter cells were observed in the cerebral and pedal ganglia. However, in the right pleural ganglion no fulicin-like neurons were present, although it was rich in immunoreactive fibers. Within the left parietal and visceral ganglia, clusters of large prominent cells appeared to give rise to axons projecting to the anal and intestinal nerves. Additionally, peripheral tissue of *B. glabrata*, specifically regions of the mantle, lip and tentacle were rich in fulicin-like immunoreactive fibers and cell bodies. These results suggest that fulicin and other peptides

129 Localization of Fulicin-like Immunoreactivity in the Central Nervous System and Periphery of *Biomphalaria glabrata*
Hernandez-Vasquez et al.

derived from the fulicin precursor could regulate behaviors related to food intake, reproduction, and growth that are altered during the course of infection in this host-parasite system.

Keywords: Biomphalaria Glabrata, Schistosomiasis, Snail Fever

ACKNOWLEDGEMENTS

The STEP-UP HS program is supported by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health, Grant number: 1R25DK098067-01.