

Journal of Health Disparities Research and Practice

Volume 12, Issue 4

2018

Article 14

2019 STEP-UP SPECIAL ISSUE

Assessing Indications of Riskiness in Adolescents

Niral Patel*

Roland Alexander Blackwood, MD, PhD[†]

*

[†]University of Michigan

Copyright ©2018 by the authors. *Journal of Health Disparities Research and Practice* is produced by The Berkeley Electronic Press (bepress). <https://digitalscholarship.unlv.edu/jhdrp>

Assessing Indications of Riskiness in Adolescents*

Niral Patel and Roland Alexander Blackwood, MD, PhD

Abstract

Adolescents are often likely to engage in perilous behaviors during their transitional years from youth to adulthood. Nearly 75% of the primary causes of death in the adolescent population are of preventable causes. In order to address the harmful issues facing this young population today, researchers at the University of Michigan developed the Rapid Assessment for Adolescent Preventive Services (RAAPS), a 21-question risk screening examination that identifies the behaviors contributing the most to adolescent morbidity, mortality, and social problems.

Over the course of this study, beginning in 2015, researchers utilized the RAAPS to gather data from University of Michigan Health System (UMHS) health clinics around Southeast Michigan. Clinic patients between the ages of 11 and 21 were administered the questionnaire as part of their primary care visit. The simple yes/no structure of each question allowed for a straightforward entry of data that could be analyzed to compare different factors affecting adolescents. Data from the assessment was entered into the Statistical Package for the Social Sciences (SPSS) statistical software for an in-depth analysis by researchers through the use of the Chi-squared test for categorical variables.

Results regarding the data obtained from the assessment are still pending at the time of abstract submission; however, it is predicted that significant indications of harmful behavior will be linked to factors such as sexual orientation and/or median household income-level. These findings may suggest a new direction in which physicians can address issues for adolescents who identify with certain characteristics.

KEYWORDS: Adolescent risk assessment; risk behavior; University of Michigan Health System; questionnaire

*The STEP-UP HS program is supported by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health, Grant number: R25DK113659.



Journal of Health Disparities Research and Practice
Volume 12, STEP-UP Special Issue, Summer 2019, pp. 25
© 2011 Center for Health Disparities Research
School of Public Health
University of Nevada, Las Vegas

Assessing Indications of Riskiness in Adolescents

Niral Patel

Roland Alexander Blackwood, MD, PhD, University of Michigan

Coordinating Center: University of California, Los Angeles

ABSTRACT

Adolescents are often likely to engage in perilous behaviors during their transitional years from youth to adulthood. Nearly 75% of the primary causes of death in the adolescent population are of preventable causes. In order to address the harmful issues facing this young population today, researchers at the University of Michigan developed the Rapid Assessment for Adolescent Preventive Services (RAAPS), a 21-question risk screening examination that identifies the behaviors contributing the most to adolescent morbidity, mortality, and social problems.

Over the course of this study, beginning in 2015, researchers utilized the RAAPS to gather data from University of Michigan Health System (UMHS) health clinics around Southeast Michigan. Clinic patients between the ages of 11 and 21 were administered the questionnaire as part of their primary care visit. The simple yes/no structure of each question allowed for a straightforward entry of data that could be analyzed to compare different factors affecting adolescents. Data from the assessment was entered into the Statistical Package for the Social Sciences (SPSS) statistical software for an in-depth analysis by researchers through the use of the Chi-squared test for categorical variables.

Results regarding the data obtained from the assessment are still pending at the time of abstract submission; however, it is predicted that significant indications of harmful behavior will be linked to factors such as sexual orientation and/or median household income-level. These findings may suggest a new direction in which physicians can address issues for adolescents who identify with certain characteristics.

Keywords: Adolescent risk assessment, risk behavior, University of Michigan Health System, questionnaire

ACKNOWLEDGEMENTS

The STEP-UP HS program is supported by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health, Grant number: R25DK113659.

Journal of Health Disparities Research and Practice Volume 12, STEP-UP Special Issue,
Summer 2019

<http://digitalscholarship.unlv.edu/jhdrp/>

Follow on Facebook: Health.Disparities.Journal

Follow on Twitter: @jhdrp