



Bioanalytical Assay of Antimicrobial Polymers Binding to Bacterial Cells

Journal of Health Disparities Research and Practice

Volume 12
Issue 4 2019 *STEP-UP Special Issue*

Article 46

© Center for Health Disparities Research, School of Public Health, University of Nevada, Las Vegas

2018

Bioanalytical Assay of Antimicrobial Polymers Binding to Bacterial Cells

Natalia Roberts

Charles V. Rice, PhD , *Department of Chemistry and Biochemistry, University of Oklahoma*

Melissa Foxley, MS , *University of Oklahoma*

See next page for additional authors

Follow this and additional works at: <https://digitalscholarship.unlv.edu/jhdrp>

 Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Roberts, Natalia; Rice, PhD, Charles V.; Foxley, MS, Melissa; Strange, Stoffel; Xiao, Min; and Wright, Summer (2018) "Bioanalytical Assay of Antimicrobial Polymers Binding to Bacterial Cells," *Journal of Health Disparities Research and Practice*: Vol. 12: Iss. 4, Article 46.

Available at: <https://digitalscholarship.unlv.edu/jhdrp/vol12/iss4/46>

This Article is protected by copyright and/or related rights. It has been brought to you by Digital Scholarship@UNLV with permission from the rights-holder(s). You are free to use this Article in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/or on the work itself.

This Article has been accepted for inclusion in Journal of Health Disparities Research and Practice by an authorized administrator of Digital Scholarship@UNLV. For more information, please contact digitalscholarship@unlv.edu.

Bioanalytical Assay of Antimicrobial Polymers Binding to Bacterial Cells

Abstract

Branched polyethylenimine (BPEI) has an antimicrobial effect on bacteria. The killing mechanism of BPEI centers on its cationic properties. The mechanism of action against Gram-positive bacteria is less understood but recent reports erroneously suggest that membrane depolarization occurs. To the contrary, data from our laboratory suggests that BPEI binds to the anionic sites provided by the biopolymer wall teichoic acid (WTA). To test the validity of this hypothesis, we measure the amount BPEI binding to whole, intact, bacterial cells of *Bacillus subtilis*. Comparative measurements are made with *Bacillus subtilis* bacteria that contain WTA and *Bacillus subtilis* genetic mutants that lack WTA.

Using equilibrium dialysis, *Bacillus subtilis* bacteria were exposed to different solution concentrations of BPEI. Removal of small aliquots from solution and subsequent assay with the ninhydrin test were used to measure the amount of BPEI remaining in solution and the amount of BPEI bound to the bacterial cell walls. These data were used to obtain the amount of bound vs. unbound BPEI and determine the equilibrium constant. These data influence the understanding of BPEI antimicrobial properties and impacts the development of antibiotics to treat human disease.

Keywords

Branched polyethylenimine (BPEI); *Bacillus subtilis*; wall teichoic acid (WTA); equilibrium dialysis

Cover Page Footnote

The STEP-UP HS program is supported by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health, Grant number: 1R25DK098067-01.

Authors

Natalia Roberts; Charles V. Rice, PhD; Melissa Foxley, MS; Stoffel Strange; Min Xiao; and Summer Wright



Journal of Health Disparities Research and Practice
Volume 12, STEP-UP Special Issue, Summer 2019, pp. 67
© 2011 Center for Health Disparities Research
School of Public Health
University of Nevada, Las Vegas

Bioanalytical Assay of Antimicrobial Polymers Binding to Bacterial Cells

Natalia Roberts

Charles V. Rice, PhD, Department of Chemistry and Biochemistry, University of Oklahoma

Melissa Foxley, MS, University of Oklahoma

Stoffel Strange, University of Oklahoma

Min Xiao, University of Oklahoma

Summer Wright, University of Oklahoma

Coordinating Center: University of Nevada, Las Vegas

ABSTRACT

Branched polyethylenimine (BPEI) has an antimicrobial effect on bacteria. The killing mechanism of BPEI centers on its cationic properties. The mechanism of action against Gram-positive bacteria is less understood but recent reports erroneously suggest that membrane depolarization occurs. To the contrary, data from our laboratory suggests that BPEI binds to the anionic sites provided by the biopolymer wall teichoic acid (WTA). To test the validity of this hypothesis, we measure the amount BPEI binding to whole, intact, bacterial cells of *Bacillus subtilis*. Comparative measurements are made with *Bacillus subtilis* bacteria that contain WTA and *Bacillus subtilis* genetic mutants that lack WTA.

Using equilibrium dialysis, *Bacillus subtilis* bacteria were exposed to different solution concentrations of BPEI. Removal of small aliquots from solution and subsequent assay with the ninhydrin test were used to measure the amount of BPEI remaining in solution and the amount of BPEI bound to the bacterial cell walls. These data were used to obtain the amount of bound vs. unbound BPEI and determine the equilibrium constant. These data influence the understanding of BPEI antimicrobial properties and impacts the development of antibiotics to treat human disease.

Keywords: Branched polyethylenimine (BPEI), *Bacillus subtilis*, wall teichoic acid (WTA), equilibrium dialysis

ACKNOWLEDGEMENTS

The STEP-UP HS program is supported by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health, Grant number: 1R25DK098067-01.

Journal of Health Disparities Research and Practice Volume 12, STEP-UP Special Issue,
Summer 2019

<http://digitalscholarship.unlv.edu/jhdp/>

Follow on Facebook: Health.Disparities.Journal

Follow on Twitter: @jhdp