
An Article Submitted to

*Journal of Health Disparities Research
and Practice*

Manuscript 1989

Analysis of Knee Motion to Prevent
and Treat the Increasing Incidence of
Premature Knee OA

Author #1*

Author #2[†]

Author #3[‡]

*

†

‡

Analysis of Knee Motion to Prevent and Treat the Increasing Incidence of Premature Knee OA

Author #1, Author #2, and Author #3

Abstract

Purpose/Background: Osteoarthritis (OA) is a costly, debilitating musculoskeletal disease. A disparity in OA incidence in young individuals who don body borne load while physically active, such as service members, reportedly exists. Incidence of premature OA in service members is greater than twice the general population and steadily increasing at the knee. Moreover, there is disparity in knee OA incidence for female and African-American service members, who are 15%-25% more likely to develop the disease. A service member's physical activity directly contributes to premature knee OA development. Specific knee locomotion biomechanics indicative of joint instability, including large and/or abrupt knee adduction motions, may increase their likelihood of OA development. Yet, no objective measure of knee instability currently exists; nor is it understood how body borne load leads to altered, and potentially hazardous knee locomotion biomechanics that increase incidence of premature knee OA among service members. This study sought to determine whether knee motions posited to be indicative of joint instability progressively increase with body-borne load and duration of load carriage.

Materials & Methods: To date, six recreationally active adults had knee biomechanics quantified while walking 1.3 m/s for 60 minutes with a different, randomly ordered body borne load: (unloaded: 0 kg, 15 kg and 30 kg). During the load carriage task, dominant limb knee biomechanics were quantified from three walk trials ($1.3 \text{ m/s} \pm 5\%$) recorded at minute 0, and every 15 minutes thereafter (i.e., 5, 15, 30, etc.). For each trial, the peak, range, and jerk cost of knee adduction angle was quantified using data from the video-based motion capture system and accelerometer-based IMU sensors. To determine whether body-borne load or the duration of load carriage increases knee joint instability, each dependent knee adduction measure was submitted to a RM ANOVA to test the main effect and interaction between time (minutes 0, 15, 30, . . . 60) and load (0, 15 and 30 kg). To compare the video-based and accelerometer-based data, each dependent knee adduction measure was submitted to equivalence testing, or two one-sided t-tests of the difference between these measures, in the same participant. Alpha was $p < 0.05$.

Results: Preliminary analysis found neither body borne load, nor time had a significant effect on peak, range, or jerk cost of knee adduction ($p > 0.05$) (Table 1 and Figure 1). No significant difference was evident between the video-based and accelerometer-based derived knee adduction measures ($p > 0.05$).

Discussion/Conclusion: Walking with body borne load may increase knee OA risk for service members. During locomotion, current participants exhibited a non-significant increased in range and jerk cost of knee adduction, and OA risk with each addition of body borne load. Specifically, participants exhibited a 11% and 22% increase in range and jerk cost of knee adduction with the 30 kg load. In contradiction to our hypothesis, participants did not further increase knee adduction throughout the duration of locomotion, despite the 33% increase in jerk cost after 45 minutes of load carriage. With additional testing, we expect a statistically significant increase range and jerk cost of knee adduction with body borne load, but only an increase in jerk cost throughout the duration of load carriage.

KEYWORDS: Knee Motion; Knee. Knee OA



Journal of Health Disparities Research and Practice
Volume 12, MW CTR-IN Special Issue, Summer 2019, pp. 7-9
© 2011 Center for Health Disparities Research
School of Public Health
University of Nevada, Las Vegas

Analysis of Knee Motion to Prevent and Treat the Increasing Incidence of Premature Knee OA

Tyler Brown, Boise State University

Micah Drew, Boise State University

Samantha Krammer, Boise State University

Corresponding Author: Tyler Brown, tynbrown@boisestate.edu

ABSTRACT

Purpose/Background: Osteoarthritis (OA) is a costly, debilitating musculoskeletal disease. A disparity in OA incidence in young individuals who don't body borne load while physically active, such as service members, reportedly exists. Incidence of premature OA in service members is greater than twice the general population and steadily increasing at the knee. Moreover, there is disparity in knee OA incidence for female and African-American service members, who are 15%-25% more likely to develop the disease. A service member's physical activity directly contributes to premature knee OA development. Specific knee locomotion biomechanics indicative of joint instability, including large and/or abrupt knee adduction motions, may increase their likelihood of OA development. Yet, no objective measure of knee instability currently exists; nor is it understood how body borne load leads to altered, and potentially hazardous knee locomotion biomechanics that increase incidence of premature knee OA among service members. This study sought to determine whether knee motions posited to be indicative of joint instability progressively increase with body-borne load and duration of load carriage.

Materials & Methods: To date, six recreationally active adults had knee biomechanics quantified while walking 1.3 m/s for 60 minutes with a different, randomly ordered body borne load: (unloaded: 0 kg, 15 kg and 30 kg). During the load carriage task, dominant limb knee biomechanics were quantified from three walk trials ($1.3 \text{ m/s} \pm 5\%$) recorded at minute 0, and every 15 minutes thereafter (i.e., 5, 15, 30, etc.). For each trial, the peak, range, and jerk cost of knee adduction angle was quantified using data from the video-based motion capture system and accelerometer-based IMU sensors. To determine whether body-borne load or the duration of load carriage increases knee joint instability, each dependent knee adduction measure was submitted to a RM ANOVA to test the main effect and interaction between time (minutes 0, 15, 30...60) and load (0,15 and 30 kg). To compare the video-based and accelerometer-based data, each dependent

Journal of Health Disparities Research and Practice Volume 12, MW-CTRIN Special Issue,
Summer 2019

<http://digitalscholarship.unlv.edu/jhdrp/>

Follow on Facebook: Health.Disparities.Journal

Follow on Twitter: @jhdrp

8 Analysis of Knee Motion to Prevention and Treat the Increasing Incidence of Premature Knee OA

Brown, Drew, and Krammer

knee adduction measure was submitted to equivalence testing, or two one-sided t-tests of the difference between these measures, in the same participant. Alpha was $p < 0.05$.

Results: Preliminary analysis found neither body borne load, nor time had a significant effect on peak, range, or jerk cost of knee adduction ($p > 0.05$) (Table 1 and Figure 1). No significant difference was evident between the video-based and accelerometer-based derived knee adduction measures ($p > 0.05$).

Discussion/Conclusion: Walking with body borne load may increase knee OA risk for service members. During locomotion, current participants exhibited a non-significant increased in range and jerk cost of knee adduction, and OA risk with each addition of body borne load. Specifically, participants exhibited a 11% and 22% increase in range and jerk cost of knee adduction with the 30 kg load. In contradiction to our hypothesis, participants did not further increase knee adduction throughout the duration of locomotion, despite the 33% increase in jerk cost after 45 minutes of load carriage. With additional testing, we expect a statistically significant increase range and jerk cost of knee adduction with body borne load, but only an increase in jerk cost throughout the duration of load carriage.

Table 1. Mean peak, range, and jerk cost of knee adduction exhibited during stance during locomotion with each body borne load.

Time	Peak (deg)			Range (deg)			Jerk Cost (deg ² /sec ⁵)		
	0 kg	15 kg	30 kg	0 kg	15 kg	30 kg	0 kg	15 kg	30 kg
0	3.62	3.21	3.21	4.57	3.32	4.54	36.35	38.59	44.38
15	3.55	4.00	3.16	3.67	3.48	4.35	36.38	42.14	46.04
30	3.73	4.10	3.31	3.69	3.84	4.64	29.24	39.70	63.52
45	4.20	3.95	3.11	4.00	3.69	4.25	79.63	41.29	39.46
60	3.73	3.75	3.19	4.12	3.48	4.59	52.38	39.46	66.89

9 Analysis of Knee Motion to Prevention and Treat the Increasing Incidence of Premature Knee OA

Brown, Drew, and Kramer




Figure 1. Depicts knee adduction joint angle (A) and angular jerk cost (B) during the stance phase (heel strike to toe off) of locomotion with each body-borne loads (0, 15 and 30 kg).