

---

An Article Submitted to

*Journal of Health Disparities Research  
and Practice*

Manuscript 1992

---

Augmenting Exposure Therapy for  
Social Anxiety with tDCS

Author #1\*

Author #2<sup>†</sup>

Author #3<sup>‡</sup>

Author #4\*\*

\*

†

‡

\*\*

# Augmenting Exposure Therapy for Social Anxiety with tDCS

Author #1, Author #2, Author #3, and Author #4

## Abstract

**Purpose/Background:** Exposure therapy is one of the most potent techniques available to treat social anxiety. However, studies suggest that exposure therapy only produces full remission in 20-50% of patients. Furthermore, laboratory conditioning and extinction studies suggest that fear responses toward individuals who differ from one's own ethnicity/race may be more resistant to extinction. Because activation of the medial prefrontal cortex has been associated with facilitating fear reduction during exposure therapy, we expect that targeting activation of this region with a stimulation technique called transcranial direct current stimulation (tDCS) may improve outcomes from exposure therapy for social anxiety. The present study will therefore test the hypotheses that (1) fear responding at baseline will be greater toward an audience that does not match (vs matches) the participant's own ethnicity, (2) pairing exposure therapy with active (vs sham) tDCS will facilitate alleviation of social anxiety symptoms, and (3) pairing exposure therapy with active (vs sham) tDCS facilitates extinction of fear response toward individuals who differ from the participant's own ethnicity.

**Materials & Methods:** We are recruiting Latino and non-Latino/Caucasian undergraduates with a fear of public speaking, the most commonly feared situation among individuals with social anxiety. Participants (N = 128) will receive either active/anodal (n = 64) or sham (n = 64) tDCS stimulation targeting the mPFC during an exposure therapy session delivered through virtual reality (VR). During exposure therapy, participants will complete six, 3-minute public speaking trials, alternating in a randomized order between audiences that are 75% matched to the participant's ethnicity and 75% unmatched to the participant's ethnicity. At one-month follow up, participants will complete two behavioral avoidance tests (BATs) parallel to therapy procedures, with one ethnic-matched trial and one ethnic-unmatched trial. Fear response during each BAT will be assessed behaviorally (duration of speech), physiologically (heart rate variability and electrodermal response), and subjectively (peak fear rating, on a 0 to 100 scale). At baseline and one-month follow-up, participants will also complete a battery of social anxiety questionnaires.

**Results:** We will present methods and preliminary findings from the study. Results will include a preliminary examination of whether fear responding is greater toward individuals who differ from (vs match) the participant's own ethnicity, whether pairing exposure therapy with active (vs sham) tDCS facilitates alleviation of social anxiety symptoms overall, and whether pairing exposure therapy with active (vs sham) tDCS facilitates alleviation of social anxiety responding toward individuals who differ from (vs match) the participant's own ethnicity.

Discussion/Conclusion: Findings point to key strategies to improve outcomes from exposure therapy for social anxiety, and could also have implications for improving response to exposure-based therapies for other anxiety disorders. Furthermore, if tDCS facilitates reductions in fear response toward ethnic/racial out-groups, minority/Latino individuals may experience better generalization of treatment effects for daily-life scenarios (in which they are surrounded by outgroup members), and ethnic/racial majority individuals will be better able to contribute to an inclusive social environment.

**KEYWORDS:** Exposure Therapy; Social Anxiety; tDCS



**Journal of Health Disparities Research and Practice**  
**Volume 12, MW CTR-IN Special Issue, Summer 2019, pp. 13-14**  
© 2011 Center for Health Disparities Research  
School of Public Health  
University of Nevada, Las Vegas

## **Augmenting Exposure Therapy for Social Anxiety with tDCS**

Cynthia L. Lancaster, University of Nevada, Reno  
Samantha J. Meckes, University of Nevada, Reno  
Mollie A. McDonald, University of Nevada, Reno  
Marian E. Berryhill, University of Nevada, Reno

*Corresponding Author:* Cynthia L. Lancaster, [cynthialancaster@unr.edu](mailto:cynthialancaster@unr.edu)

### **ABSTRACT**

**Purpose/Background:** Exposure therapy is one of the most potent techniques available to treat social anxiety. However, studies suggest that exposure therapy only produces full remission in 20-50% of patients. Furthermore, laboratory conditioning and extinction studies suggest that fear responses toward individuals who differ from one's own ethnicity/race may be more resistant to extinction. Because activation of the medial prefrontal cortex has been associated with facilitating fear reduction during exposure therapy, we expect that targeting activation of this region with a stimulation technique called transcranial direct current stimulation (tDCS) may improve outcomes from exposure therapy for social anxiety. The present study will therefore test the hypotheses that (1) fear responding at baseline will be greater toward an audience that does not match (vs matches) the participant's own ethnicity, (2) pairing exposure therapy with active (vs sham) tDCS will facilitate alleviation of social anxiety symptoms, and (3) pairing exposure therapy with active (vs sham) tDCS facilitates extinction of fear response toward individuals who differ from the participant's own ethnicity.

**Materials & Methods:** We are recruiting Latino and non-Latino/Caucasian undergraduates with a fear of public speaking, the most commonly feared situation among individuals with social anxiety. Participants (N = 128) will receive either active/anodal (n = 64) or sham (n = 64) tDCS stimulation targeting the mPFC during an exposure therapy session delivered through virtual reality (VR). During exposure therapy, participants will complete six, 3-minute public speaking trials, alternating in a randomized order between audiences that are 75% matched to the participant's ethnicity and 75% unmatched to the participant's ethnicity. At one-month follow up, participants will complete two behavioral avoidance tests (BATs) parallel to therapy procedures, with one ethnic-matched trial and one ethnic-unmatched trial. Fear response during each BAT will be assessed behaviorally (duration of speech), physiologically (heart rate variability and electrodermal response), and subjectively (peak fear rating, on a 0 to 100 scale). At baseline and one-month follow-up, participants will also complete a battery of social anxiety questionnaires.

**Results:** We will present methods and preliminary findings from the study. Results will

Journal of Health Disparities Research and Practice Volume 12, MW-CTRIN Special Issue,  
Summer 2019

<http://digitalscholarship.unlv.edu/jhdrp/>

Follow on Facebook: [Health.Disparities.Journal](https://www.facebook.com/Health.Disparities.Journal)

Follow on Twitter: [@jhdrp](https://twitter.com/jhdrp)

14 Augmenting Exposure Therapy for Social Anxiety with tDCS  
Lancaster, Meckes, McDonald, and Berryhill

include a preliminary examination of whether fear responding is greater toward individuals who differ from (vs match) the participant's own ethnicity, whether pairing exposure therapy with active (vs sham) tDCS facilitates alleviation of social anxiety symptoms overall, and whether pairing exposure therapy with active (vs sham) tDCS facilitates alleviation of social anxiety responding toward individuals who differ from (vs match) the participant's own ethnicity.

Discussion/Conclusion: Findings point to key strategies to improve outcomes from exposure therapy for social anxiety, and could also have implications for improving response to exposure-based therapies for other anxiety disorders. Furthermore, if tDCS facilitates reductions in fear response toward ethnic/racial out-groups, minority/Latino individuals may experience better generalization of treatment effects for daily-life scenarios (in which they are surrounded by outgroup members), and ethnic/racial majority individuals will be better able to contribute to an inclusive social environment.