

Aug 3rd, 9:00 AM - 12:00 PM

High pressure infrared studies of HMX

Jennifer Wojno
University of Louisville

Michael G. Pravica
University of Nevada, Las Vegas, pravica@physics.unlv.edu

Martin Galley
University of Nevada, Las Vegas

Follow this and additional works at: https://digitalscholarship.unlv.edu/cs_urop

 Part of the [Biological and Chemical Physics Commons](#)

Repository Citation

Wojno, Jennifer; Pravica, Michael G.; and Galley, Martin, "High pressure infrared studies of HMX" (2010). *Undergraduate Research Opportunities Program (UROP)*. 16.
https://digitalscholarship.unlv.edu/cs_urop/2010/aug3/16

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Scholarship@UNLV with permission from the rights-holder(s). You are free to use this Event in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/or on the work itself.

This Event has been accepted for inclusion in Undergraduate Research Opportunities Program (UROP) by an authorized administrator of Digital Scholarship@UNLV. For more information, please contact digitalscholarship@unlv.edu.

High pressure infrared studies of HMX

Jennifer Wojno¹, Michael Pravica², Martin Galley²

1. Department of Physics & Astronomy, 102 Natural Science Building, University of Louisville, Louisville, KY 40292

2. University of Nevada, Las Vegas, Department of Physics & Astronomy, Box 454002, 4505 Maryland Parkway, Las Vegas, NV 89154-4002

Abstract

We are studying the effects of pressure on HMX (Octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazocine) using infrared spectroscopy. The sample is put under pressure using a diamond anvil cell at ambient temperature, data is collected at the National Synchrotron Light Source in Brookhaven National Laboratory. In analyzing this data, we hope to learn more about the molecular vibrations as the molecule bends and deforms under pressure. Such understanding could aid in determining new safety standards or more efficient ways of using HMX. In future studies, we intend to include the aspect of temperature variation in addition to pressure, with the goal of describing the molecule in a phase diagram.

Background

High pressure physics is used to study how materials change under pressure. One of the most common methods of obtaining high pressures in the gigapascal (GPa) range is the use of the diamond anvil cell (DAC.) A figure of a typical DAC is provided below.

HMX is used in a variety of applications, although it is almost exclusively used in military applications as a secondary explosive. As it is an insensitive explosive, it will not react under most normal conditions, even though it is one of the most powerful explosives manufactured today. Because of this, HMX is very easy to study, even when it is subjected to high pressure.

Ruby is used in most high pressure applications as a way of monitoring pressure. It is used as a well established method of monitoring pressure, as ruby fluorescence peaks are very sensitive to pressure changes.




Figure 1. Illustration of a diamond anvil cell (earth.northwestern.edu)






Figure 2. Molecular structure of HMX



Graph 1. Infrared spectrum HMX compressed from 0.45 GPa to 31.18 GPa



Graph 2. Decompression of HMX, from high pressure (bottom) to low pressure (top)

Experiment

We loaded HMX into a symmetric diamond anvil cell (DAC) at Brookhaven National Laboratory. We used KBr (Potassium Bromide) as a quasi-hydrostatic pressure transmitting medium, which is invisible in the infrared spectrum. Ruby powder was used to determine the current pressure of the sample. Half of the sample was pure KBr to be used for background subtraction, the second half was a thin layer of HMX with KBr on it. A Bruker Hyperion and Vertex 80V spectrometer was used to collect the infrared spectra of the HMX sample.



Figure 3. Symmetric DAC (top)



Figure 4. Symmetric DAC (side)


Data was collected at approximately 1.5 GPa intervals, up to a maximum of 30 GPa from ambient pressure. Ruby fluorescence was monitored using a 514 nm diode laser.

The collected data was then analyzed using OriginLab8 and CrystalSleuth. Each of the peaks was then matched to a database to determine which chemical bonds or vibrations it represents. This analysis is consolidated in a table to the right.

Analysis

Analysis of HMX major peaks at 0.45 GPa

Peak Number	Theoretical (cm ⁻¹)	Experimental (cm ⁻¹)	Bond Type
1	600	603	Skeleton motion (whole ring vibration), ring vibration
2	628	629	CH ₂ rk, NO ₂ rk/sc
3	660	660	N-N str, NO ₂ sc, CNC sc, CH ₂ rk
4	758	761	NO ₂ wg, N-N wg, CNC sc
5	771	773	NO ₂ wg, N-N rk
6	831	831	CNC str, NO ₂ str
7	857	860	CNC str, NO ₂ str
8	880	871	N-N str, NO ₂ sc, CNC str, CH ₂ rk
9	927	947	N-N str, CH ₂ rk, CNC str
10	1066	1089	CH ₂ tw, CH ₂ rk, N-N str (all bonds)
11	1136	1147	CH ₂ rk (all bonds), N-N str (all bonds)
12	1221	1205	CH wg, N-N str, CNC asym str, CNC str, NO ₂ str
13	1250	1269	CH wg, CNC asym str, NO ₂ str
14	1296	1290	CH ₂ wg, CH wg, NO ₂ str, N-N str
15	1315	1326	CH ₂ wg, NO ₂ str, CH ₂ sc, N-N str, CNC asym str
16	1341	1350	CH ₂ tw, CNC asym str
17	1363/1374	1396	CH ₂ sc, CH ₂ wg
18	1414	1433	CH ₂ sc
19	1422	1467	CH ₂ sc
20	1668	1556	NO ₂ asym str
21	2990	2998	CH ₂ str
22	3042/3053	3039	CH ₂ asym str



Graph 3. Infrared spectrum at 0.45 GPa

In analyzing the data obtained from each pressure, we are able to discern how certain peaks move or change with pressure, and how this may represent changes in the molecular structure. The above table compiles information for each of the major peaks and their corresponding bond types. Such tables were constructed for each pressure increase.

Each of the peaks are labeled according to their number on the graph to the left. These peaks change or disappear with increasing pressure, as shown in the waterfall plot, however in decompression, changes are reversed.

From this analysis we can determine no irreversible phase change as upon decompression the sample's spectrum returned to its ambient state.

References

Reusch, William. "Infrared Spectroscopy." *Infrared Spectroscopy*. William Reusch, 10 Aug 2007. Web. 15 Jun 2010. <<http://www2.chemistry.msu.edu/faculty/reusch/VirtTxtJml/Spectrpy/InfraRed/infrared.htm>>.

Chickos, James, and David Garin. "Infrared Spectroscopy." *INTERACTIVE INSTRUCTIONAL TEXT*. UM-St. Louis Chemistry Department, 08 Dec 1997. Web. 28 Jul 2010. <<http://www.umsl.edu/~orglab/documents/IR/IR2.html>>.

Support from the REU program of the National Science Foundation under grant DMR-1005247 is gratefully acknowledged.

