



**Journal of Health Disparities Research and Practice**  
**Volume 9, Special Edition 1, Summer 2016, pp. 6-7**  
© 2011 Center for Health Disparities Research  
School of Community Health Sciences  
University of Nevada, Las Vegas

## **Induction of DUSP9 in Xenografts from Human Breast Cancer Cell Lines increases Mammary Cancer Stem Cells**

Albert Barrios

Meher Parveen, PhD, Charles R. Drew University of Medicine and Science

Easter Thames, Charles R. Drew University of Medicine and Science

Melanie Baker, Charles R. Drew University of Medicine and Science

Shelha Pervin, PhD, Charles R. Drew University of Medicine and Science

**Coordinating Center:** Charles R. Drew University of Medicine and Science

### **ABSTRACT**

Breast cancer remains a complex disease that kills 40,000 women every year. Initiation and progression of breast cancer is influenced by heterogeneous groups of cells, including mammary cancer stem cells (MCSCs). Progression of this dreadful disease is driven by many signaling pathways among which MAPK pathway is highly prominent. Since targeting prominent kinases in MAPK pathway has been unsuccessful to control breast cancer, it is important to examine the phosphatases that regulate the activity of these kinases.

Using xenograft model from breast cancer cell lines, our lab has found that during the initial stages of xenograft development (week 1-4, 100-200mg weight), ERK1/2 remains inactive. However, during the later stages of tumor development (week 5-15, 300-700mg weight), we found phospho-ERK1/2, which is the active form of ERK1/2 remains highly up regulated. Our lab has also found that presence of inactive ERK1/2 during the initial stages of tumor development was independent of pMEK1/2, which is the upstream kinase that activates ERK1/2. We have also found that a specific dual specific phosphatase (DUSP9) was induced in the xenografts from breast cancer cell lines during initial stages of their development.

I will be examining the significance of DUSP9 induction in the early stages of xenograft development. There are reports that DUSP9-mediated pERK1/2 inactivation has been found to increase mouse embryonic stem cell content. I have been examining the expression levels of MCSCS (Aldehyde dehydrogenase 1(ALDH1)/ OCT3/4 / CD44/SOX2) in various stages of xenograft development and correlating them with DUSP9 expression and pERK1/2. I am examining the expression levels of MCSCs by Western blot analysis as well as qPCR. I have also been treating breast cancer cell line HMLE<sup>HRASV12</sup> with MEK inhibitors, *in vitro*, for inactivating ERK to analyze the levels of mammary cancer stem cells (ALDH1)/ OCT3/4 /

## 7 Induction of DUSP9 in Xenografts from Human Breast Cancer Cell Lines increases Mammary Cancer Stem Cells

Barrios et al.

CD44/SOX2. I am also examining the levels of DUSP9 as well as the levels of ERK/pERK1/2 /AKT /pAKT in cells treated with MEK1/2 inhibitors *in vitro*. I intend to determine whether inhibition of pERK1/2 could influence embryonic stem cell content as analyzed by the expression of markers Oct <sup>3</sup>/<sub>4</sub> and Sox2.

**Keywords:** Breast Cancer, Mammary Cancer Stem Cells, Xenograft

### **ACKNOWLEDGEMENTS**

The STEP-UP HS program is supported by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health, Grant Number: 5R25DK078384-09.