



**Journal of Health Disparities Research and Practice**  
**Volume 9, Special Edition 1, Summer 2016, pp. 76-77**  
© 2011 Center for Health Disparities Research  
School of Community Health Sciences  
University of Nevada, Las Vegas

## **Effects of Salinity on Plant Development of *in vitro* Sweet Potato (*Ipomoea batatas*) Explants and Growth of Pakchoi (*Brassica rapa* var. *chinensis*) Seedlings**

Tristan Paulino

Mari Marutani, PhD, University of Guam

**Coordinating Center:** University of Hawaii John A. Burns School of Medicine

### **ABSTRACT**

Climate change has affected the Pacific Islands with increased sea levels, and the contamination of freshwater with saltwater, which has become a serious problem for plant development. Bioassays were performed to examine the effects of salinity on root and shoot development of *in-vitro* sweet potato (*Ipomoea batatas*), and effects on the seedling growth of pakchoi (*Brassica rapa* var. *chinensis*).

Sterilized stems with one or two nodes of sweet potato cv. Liberty were placed in *in-vitro* Murashige & Skoog solid media with NaCl concentrations of 0%, 0.5%, 1.0% and 1.5%. Newly emerged roots and shoots were recorded on the 18th days after initiation of the experiment. For the second bioassay, two pakchoi cultivars, 'Speedy' and 'Bino,' were tested for seedling development in 0%, 0.5%, 1.0%, and 1.5% NaCl solutions. The fresh weights of seedlings were recorded on the 15th day. Simple statistics and ANOVA were performed to compare plant growth.

Sweet potato explants generated more roots (14.8) in control (0%) than those in 0.5% (6.9), 1.0% (5.4), and 1.5% (5.2) ( $n=12$ ,  $p<0.05$ ). The number of shoots was greater with control (5.4) and 0.5% (5.2) than those in 1.0% (3.0) and 1.5% (3.8). Both pakchoi cultivars showed seedling growth only in 0% and 0.5% NaCl solutions. No growth was observed in 1.0% and 1.5% solutions. Seedling fresh weight of 'Bino' was greater than 'Speedy' ( $n=5$ ,  $p<0.05$ ). Tolerance level of pakchoi varied between two cultivars. Increasing salinity had a negative impact on plant growth.

**Key Words:** Sweet Potato, Pakchoi, Salinity, Bioassay, In-Vitro Culture

77 Effects of Salinity on Plant Development of *in vitro* Sweet Potato (*Ipomoea batatas*)  
Explants and Growth of Pakchoi (*Brassica rapa* var. *chinensis*) Seedlings  
Paulino and Marutani

#### **ACKNOWLEDGEMENTS**

The STEP-UP HS program is supported by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health, Grant number: R25DK078386.