



Effects of Salinity on Plant Development of in vitro Sweet Potato (*Ipomoea batatas*)  
Explants and Growth of Pakchoi (*Brassica rapa* var. *chinensis*) Seedlings

## Journal of Health Disparities Research and Practice

Volume 9  
Issue 5 *Special Issue - NIDDK STEP UP*

Article 51

© Center for Health Disparities Research, School of Public Health, University of Nevada, Las Vegas

2016

# Effects of Salinity on Plant Development of in vitro Sweet Potato (*Ipomoea batatas*) Explants and Growth of Pakchoi (*Brassica rapa* var. *chinensis*) Seedlings

Tristan Paulino

Mari Marutani, PhD, *University of Guam*

Follow this and additional works at: <https://digitalscholarship.unlv.edu/jhdrp>



Part of the [Bilingual, Multilingual, and Multicultural Education Commons](#), [Community College Leadership](#)

[Commons](#), [Higher Education Commons](#), [Immune System Diseases Commons](#), [Public Health Commons](#), [Translational Medical Research Commons](#), and the [Virus Diseases Commons](#)

### Recommended Citation

Paulino, Tristan and Marutani, PhD, Mari (2016) "Effects of Salinity on Plant Development of in vitro Sweet Potato (*Ipomoea batatas*) Explants and Growth of Pakchoi (*Brassica rapa* var. *chinensis*) Seedlings,"

*Journal of Health Disparities Research and Practice*. Vol. 9: Iss. 5, Article 51.

Available at: <https://digitalscholarship.unlv.edu/jhdrp/vol9/iss5/51>

This Article is protected by copyright and/or related rights. It has been brought to you by Digital Scholarship@UNLV with permission from the rights-holder(s). You are free to use this Article in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/or on the work itself.

This Article has been accepted for inclusion in Journal of Health Disparities Research and Practice by an authorized administrator of Digital Scholarship@UNLV. For more information, please contact [digitalscholarship@unlv.edu](mailto:digitalscholarship@unlv.edu).

---

# Effects of Salinity on Plant Development of in vitro Sweet Potato (*Ipomoea batatas*) Explants and Growth of Pakchoi (*Brassica rapa* var. *chinensis*) Seedlings

## Abstract

Climate change has affected the Pacific Islands with increased sea levels, and the contamination of freshwater with saltwater, which has become a serious problem for plant development. Bioassays were performed to examine the effects of salinity on root and shoot development of in-vitro sweet potato (*Ipomoea batatas*), and effects on the seedling growth of pakchoi (*Brassica rapa* var. *chinensis*).

Sterilized stems with one or two nodes of sweet potato cv. Liberty were placed in in-vitro Murashige & Skoog solid media with NaCl concentrations of 0%, 0.5%, 1.0% and 1.5%. Newly emerged roots and shoots were recorded on the 18th days after initiation of the experiment. For the second bioassay, two pakchoi cultivars, 'Speedy' and 'Bino,' were tested for seedling development in 0%, 0.5%, 1.0%, and 1.5% NaCl solutions. The fresh weights of seedlings were recorded on the 15th day. Simple statistics and ANOVA were performed to compare plant growth.

Sweet potato explants generated more roots (14.8) in control (0%) than those in 0.5% (6.9), 1.0% (5.4), and 1.5% (5.2) ( $n=12$ ,  $p<0.05$ ). The number of shoots was greater with control (5.4) and 0.5% (5.2) than those in 1.0% (3.0) and 1.5% (3.8). Both pakchoi cultivars showed seedling growth only in 0% and 0.5% NaCl solutions. No growth was observed in 1.0% and 1.5% solutions. Seedling fresh weight of 'Bino' was greater than 'Speedy' ( $n=5$ ,  $p<0.05$ ). Tolerance level of pakchoi varied between two cultivars. Increasing salinity had a negative impact on plant growth.

## Keywords

Sweet Potato; Pakchoi; Salinity; Bioassay; In-Vitro Culture



**Journal of Health Disparities Research and Practice**  
**Volume 9, Special Edition 1, Summer 2016, pp. 76-77**  
© 2011 Center for Health Disparities Research  
School of Community Health Sciences  
University of Nevada, Las Vegas

## **Effects of Salinity on Plant Development of *in vitro* Sweet Potato (*Ipomoea batatas*) Explants and Growth of Pakchoi (*Brassica rapa* var. *chinensis*) Seedlings**

Tristan Paulino

Mari Marutani, PhD, University of Guam

**Coordinating Center:** University of Hawaii John A. Burns School of Medicine

### **ABSTRACT**

Climate change has affected the Pacific Islands with increased sea levels, and the contamination of freshwater with saltwater, which has become a serious problem for plant development. Bioassays were performed to examine the effects of salinity on root and shoot development of in-vitro sweet potato (*Ipomoea batatas*), and effects on the seedling growth of pakchoi (*Brassica rapa* var. *chinensis*).

Sterilized stems with one or two nodes of sweet potato cv. Liberty were placed in in-vitro Murashige & Skoog solid media with NaCl concentrations of 0%, 0.5%, 1.0% and 1.5%. Newly emerged roots and shoots were recorded on the 18th days after initiation of the experiment. For the second bioassay, two pakchoi cultivars, 'Speedy' and 'Bino,' were tested for seedling development in 0%, 0.5%, 1.0%, and 1.5% NaCl solutions. The fresh weights of seedlings were recorded on the 15th day. Simple statistics and ANOVA were performed to compare plant growth.

Sweet potato explants generated more roots (14.8) in control (0%) than those in 0.5% (6.9), 1.0% (5.4), and 1.5% (5.2) ( $n=12$ ,  $p<0.05$ ). The number of shoots was greater with control (5.4) and 0.5% (5.2) than those in 1.0% (3.0) and 1.5% (3.8). Both pakchoi cultivars showed seedling growth only in 0% and 0.5% NaCl solutions. No growth was observed in 1.0% and 1.5% solutions. Seedling fresh weight of 'Bino' was greater than 'Speedy' ( $n=5$ ,  $p<0.05$ ). Tolerance level of pakchoi varied between two cultivars. Increasing salinity had a negative impact on plant growth.

**Key Words:** Sweet Potato, Pakchoi, Salinity, Bioassay, In-Vitro Culture

77 Effects of Salinity on Plant Development of *in vitro* Sweet Potato (*Ipomoea batatas*)  
Explants and Growth of Pakchoi (*Brassica rapa* var. *chinensis*) Seedlings  
Paulino and Marutani

#### **ACKNOWLEDGEMENTS**

The STEP-UP HS program is supported by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health, Grant number: R25DK078386.