



Lipid Metabolism

Journal of Health Disparities Research and Practice

Volume 9
Issue 5 *Special Issue - NIDDK STEP UP*

Article 73

© Center for Health Disparities Research, School of Public Health, University of Nevada, Las Vegas

2016

Lipid Metabolism

Paige Moore

Lawrence Chan, DSc, MBBS, *Baylor College of Medicine*

Elaine Chen, *Baylor University*

Follow this and additional works at: <https://digitalscholarship.unlv.edu/jhdp>



Part of the [Bilingual, Multilingual, and Multicultural Education Commons](#), [Community College Leadership Commons](#), [Higher Education Commons](#), [Immune System Diseases Commons](#), [Public Health Commons](#), [Translational Medical Research Commons](#), and the [Virus Diseases Commons](#)

Recommended Citation

Moore, Paige; Chan, DSc, MBBS, Lawrence; and Chen, Elaine (2016) "Lipid Metabolism," *Journal of Health Disparities Research and Practice*: Vol. 9: Iss. 5, Article 73.

Available at: <https://digitalscholarship.unlv.edu/jhdp/vol9/iss5/73>

This Article is protected by copyright and/or related rights. It has been brought to you by Digital Scholarship@UNLV with permission from the rights-holder(s). You are free to use this Article in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/or on the work itself.

This Article has been accepted for inclusion in Journal of Health Disparities Research and Practice by an authorized administrator of Digital Scholarship@UNLV. For more information, please contact digitalscholarship@unlv.edu.

Lipid Metabolism

Abstract

Perilipin 2, also known as Adipose differentiation-related protein or PLIN2, is a lipid droplet-binding protein present in almost every tissue. The absence of PLIN2 upregulates hepatic very low-density lipoprotein secretion, relieves hepatosteatosis, and improves whole body insulin resistance in mice. Despite of the importance in mediating lipid metabolism, the regulation of PLIN2 itself remains largely unknown. Previous reports have shown that X-box binding protein 1 (XBP1) is an important regulator of lipogenesis. XBP1 is a transcription factor that recognizes and binds to a consensus sequence, 5'-TGACGTGG-3'. Interestingly, when we looked through the promoter region of mouse *Plin2* gene, we found that the consensus sequence is present in the *Plin2* promoter. Therefore, we hypothesize that XBP1 might directly bind to *Plin2* promoter and regulate the *Plin2* expression.

To test our hypothesis, we will perform the luciferase assay to examine whether the *Plin2* promoter activity is regulated by XBP1. We first designed forward and reverse PCR primers, which include BglII and BamHI restriction enzyme sites respectively, to amplify the *Plin2* promoter region (from -1100 to +40). We performed PCR and cloned the *Plin2* promoter to a TA vector. The TA vector was then sequenced to exclude any point mutations. After sequencing, we sub cloned the *Plin2* promoter into a vector containing a luciferase reporter. In the future, we will transfect 293T, a human embryonic kidney cell line, with the *Plin2* promoter-luciferase vector we generated. We will compare the *Plin2* promoter activity by measuring the luminescence in the presence or absence of XBP1.

Keywords

Plin2; XBP1; Consensus Sequence; Luciferase Assay



Journal of Health Disparities Research and Practice
Volume 9, Special Edition 1, Summer 2016, pp. 107

© 2011 Center for Health Disparities Research

School of Community Health Sciences

University of Nevada, Las Vegas

Lipid Metabolism

Paige Moore

Lawrence Chan, DSc, MBBS, Baylor College of Medicine

Elaine Chen, Baylor University

Coordinating Center: Stanford University

ABSTRACT

Perilipin 2, also known as Adipose differentiation-related protein or PLIN2, is a lipid droplet-binding protein present in almost every tissue. The absence of PLIN2 upregulates hepatic very low-density lipoprotein secretion, relieves hepatosteatosis, and improves whole body insulin resistance in mice. Despite of the importance in mediating lipid metabolism, the regulation of PLIN2 itself remains largely unknown. Previous reports have shown that X-box binding protein 1 (XBP1) is an important regulator of lipogenesis. XBP1 is a transcription factor that recognizes and binds to a consensus sequence, 5'-TGACGTGG-3'. Interestingly, when we looked through the promoter region of mouse *Plin2* gene, we found that the consensus sequence is present in the *Plin2* promoter. Therefore, we hypothesize that XBP1 might directly bind to *Plin2* promoter and regulate the *Plin2* expression.

To test our hypothesis, we will perform the luciferase assay to examine whether the *Plin2* promoter activity is regulated by XBP1. We first designed forward and reverse PCR primers, which include BglII and BamHI restriction enzyme sites respectively, to amplify the *Plin2* promoter region (from -1100 to +40). We performed PCR and cloned the *Plin2* promoter to a TA vector. The TA vector was then sequenced to exclude any point mutations. After sequencing, we sub cloned the *Plin2* promoter into a vector containing a luciferase reporter. In the future, we will transfect 293T, a human embryonic kidney cell line, with the *Plin2* promoter-luciferase vector we generated. We will compare the *Plin2* promoter activity by measuring the luminescence in the presence or absence of XBP1.

Key words: Plin2, XBP1, Consensus Sequence, Luciferase Assay

ACKNOWLEDGEMENTS

The STEP-UP HS program is supported by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health, Grant number: R25DK078382.