

12-2010

## Reconstructing species responses to past climatic changes using niche modeling and genetic data

Tereza Jezkova

*University of Nevada, Las Vegas*

Follow this and additional works at: <https://digitalscholarship.unlv.edu/thesesdissertations>



Part of the [Desert Ecology Commons](#), [Evolution Commons](#), [Genetics Commons](#), [Terrestrial and Aquatic Ecology Commons](#), and the [Zoology Commons](#)

---

### Repository Citation

Jezkova, Tereza, "Reconstructing species responses to past climatic changes using niche modeling and genetic data" (2010). *UNLV Theses, Dissertations, Professional Papers, and Capstones*. 750.  
<http://dx.doi.org/10.34917/2026082>

This Dissertation is protected by copyright and/or related rights. It has been brought to you by Digital Scholarship@UNLV with permission from the rights-holder(s). You are free to use this Dissertation in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/or on the work itself.

This Dissertation has been accepted for inclusion in UNLV Theses, Dissertations, Professional Papers, and Capstones by an authorized administrator of Digital Scholarship@UNLV. For more information, please contact [digitalscholarship@unlv.edu](mailto:digitalscholarship@unlv.edu).

RECONSTRUCTING SPECIES RESPONSES TO PAST CLIMATIC CHANGES USING NICHE  
MODELING AND GENETIC DATA

by

Tereza Jezkova

Master of Science  
Charles University, Prague, Czech Republic  
2002

A dissertation submitted in partial fulfillment of  
the requirements for the

**Doctor of Philosophy in Biological Sciences**  
**School of Life Sciences**  
**College of Science**

**Graduate College**  
**University of Nevada, Las Vegas**  
**December 2010**

Copyright by Tereza Jezkova 2011  
All Rights Reserved



**THE GRADUATE COLLEGE**

We recommend the dissertation prepared under our supervision by

**Tereza Jezkova**

entitled

**Reconstructing Species Responses to Past Climatic Changes Using  
Niche Modeling and Genetic Data**

be accepted in partial fulfillment of the requirements for the degree of

**Doctor of Philosophy in Biological Sciences**

School of Life Sciences

Brett Riddle, Committee Chair

John Klicka, Committee Member

Daniel Thompson, Committee Member

John Hafner, Committee Member

Matthew Lachniet, Graduate Faculty Representative

Ronald Smith, Ph. D., Vice President for Research and Graduate Studies  
and Dean of the Graduate College

**December 2010**

## ABSTRACT

### **Reconstructing Species Responses to Past Climatic Changes Through Niche Modeling and Genetic Data**

by

Tereza Jezkova

Dr. Brett R. Riddle, Examination Committee Chair  
Professor of Biology  
University of Nevada, Las Vegas

Glacial – interglacial cycles have a pronounced impact on species distributions and genetic structure. Many species shift their distributions to lower latitudes and altitudes during the colder glacial periods and expand northwards and up the elevation during warmer interglacial periods. Some species however are capable of adapting to changing environment which allows them to persist in place despite climatic changes. I explored how climatic changes after the last glacial maximum (LGM) effected two species inhabiting the deserts of western North America: one mammal (Chisel-toothed Kangaroo Rat, *Dipodomys microps*) and one reptile (Desert Horned Lizard, *Phrynosoma platyrhinos*). I used a methodology of transferal modeling which is commonly used to predict species responses to future climatic changes. I approximated the species current and LGM distribution by modeling their current climatic niches, which I then projected onto the climatic conditions of the LGM. The accuracy of the transferal models, however, is dependent on several conceptual and algorithmic assumptions. Therefore, I compared the models with the phylogeographic structure of each species as phylogeographic signals imprinted in species genomes can inform us about species past geographic and demographic processes. The transferal models predicted that the

northern parts of the species current ranges were unsuitable during the LGM and that both species could have persisted only within the more southern deserts where climatic conditions remained suitable. The phylogeographic analyses, however, suggested that *D. microps* did not experience large scale distributional changes in response to the warming climate after the LGM as suggested by the models and instead persisted in place throughout most of its current range. *Phrynosoma platyrhinos* expanded its range northwards after the LGM but was able to expand further than indicated by models, into colder and wetter areas than those experienced during the LGM. My results indicate that the two species responded to the warming climate after the LGM in an idiosyncratic fashion and that the transferal models did not correctly predict the species response to the climate change. These results motivated me to explore in the last chapter several high-priority challenges in transferal modeling through theoretical background and sets of experiments. I demonstrated how these challenges can affect resulting models and, when possible, offered suggestions on how uncertainties might be diminished.

## ACKNOWLEDGEMENTS

I am grateful to my advisor, Dr. Brett Riddle, who gave me the opportunity to join his research group and introduced me to the field of biogeography. I am also grateful to my other committee members, Dr. John Klicka, Dr. Daniel Thompson, Dr John Hafner, and Dr. Matthew Lachniet for their guidance throughout my graduate years. I thank Dr. Jef Jaeger who taught me a great deal of laboratory techniques and writing skills, and I am thankful for his tremendous help, support, and friendship. I also thank Dr. Javier Rodriguez, who introduced me to the world of publishing and jump started my career by letting me join several of his research projects. I thank Viktoria Hemmings for being an amazing friend and collaborator, Matthew Graham and Brian Smith for inspiring mental torture throughout the years, Mallory Eckstut, Derek Houston, Adam Leland, and Robert Bryson for valuable discussions and peer reviews, and Dr. Lois Alexander for remarkable help in the laboratory in the early stages of my graduate program. I thank my undergraduate students Anna San, Calais Fossier, Esther Rahardjo, and Kathleen Reich for their help in the laboratory and numerous students, both graduate and undergraduate, for help in the field. I thank Keala Kiko, Scott Miller, Pat Hunt, Nicholle Booker, and Eric Lee for administrative help and support.

I also appreciate the support and encouragements of my family and friends. I am especially thankful for all the love and support of my parents Jiri Jezek and Jana Jezkova and my sister Adela Jezkova. I express my deepest gratitude to my husband Aaron Ambos who spend the vast majority of his vacation days within the past five years in the harsh desert, chasing lizards and rodents, and who provided me with intellectual,

mental, and financial support throughout my graduate years. Last but not least, I would like to thank my beloved dog Akira, who loved the field work and tolerated the rest.

This research was supported by Doctoral Dissertation Improvement Grant DEB-0808565 from the National Science Foundation (NSF), University of Nevada, Las Vegas (UNLV) President's Research Award, EPSCoR ACES Grant SFFA UCCSN 02-123, grants from the Graduate and Professional Student Association at UNLV and from the American Society of Mammalogists, and a Major Research Instrumentation Grant DBI-0421519 to UNLV provided by the NSF.



## TABLE OF CONTENTS

|  |     |
|--|-----|
| ABSTRACT.....  | iii |
| ACKNOWLEDGEMENTS.....  | v   |
| LIST OF TABLES.....  | ix  |
| LIST OF FIGURES.....   | x   |
| CHAPTER 1 INTRODUCTION .....   | 1   |
| CHAPTER 2 NICHE SHIFT IN SITU IN RESPONSE TO THE WARMING CLIMATE AT THE<br>END OF THE LAST GLACIAL MAXIMUM: INFERENCE FROM GENETIC DATA<br>AND NICHE ASSESSMENTS IN THE CHISEL-TOOTHED KANGAROO RAT<br>( <i>DIPODOMYS MICROPS</i> )..... | 3   |
| Abstract .....   | 3   |
| Introduction.....  | 4   |
| Methods .....  | 9   |
| Results .....  | 18  |
| Discussion .....   | 22  |
| Tables.....  | 30  |
| Figures .....  | 35  |
| Bibliography.....  | 48  |
| CHAPTER 3 CLIMATIC PROMOTION OF NICHE SHIFTS: A CASE STUDY OF THE DESERT<br>HORNED LIZARD ( <i>PHRYNOSOMA PLATYRHINOS</i> ).....   | 56  |
| Abstract .....   | 56  |
| Introduction.....  | 57  |
| Methods .....  | 60  |
| Results .....  | 69  |
| Discussion .....   | 75  |
| Tables.....  | 85  |
| Figures .....  | 88  |
| Bibliography.....  | 99  |
| CHAPTER 4 SOME MORE CHALLENGES FOR ECOLOGICAL NICHE MODELS: EXPLORING<br>THE CHALLENGES OF TRANSFERAL MODELING IN BIOGEOGRAPHY .....   | 107 |
| Abstract .....   | 107 |
| Introduction.....  | 108 |
| Methods .....  | 110 |
| Challenge 1 .....  | 111 |

|  |     |
|--|-----|
| Challenge 2 .....  | 114 |
| Challenge 3 .....  | 116 |
| Challenge 4 .....  | 121 |
| Challenge 5 .....  | 125 |
| Concluding remarks.....                                      | 130 |
| Tables.....  | 132 |
| Figures .....  | 134 |
| Bibliography.....  | 152 |
| APPENDIX 1    SAMPLES OF <i>DIPODOMYS MICROPS</i> .....      | 160 |
| APPENDIX 2    SAMPLES OF <i>PHRYNOSOMA PLATYRHINOS</i> ..... | 166 |
| APPENDIX 3    GLOSSARY.....                                  | 170 |
| VITA .....   | 173 |

## LIST OF TABLES

|           |   |     |
|-----------|---|-----|
| Table 2.1 | General sampling localities for <i>Dipodomys microps</i> .....  | 30  |
| Table 2.2 | Genetic indices for the full dataset and localities with sample size greater than or equal to 10 .....                                | 34  |
| Table 3.1 | Bayes factors for different partitionings of the ND4+cytB+tRNA sequences of <i>Phrynosoma platyrhinos</i> .....                       | 85  |
| Table 3.2 | Differentiation of the climatic niche based on Tukey tests between the Persistent and expanding groups of <i>P. platyrhinos</i> ..... | 86  |
| Table 3.3 | Niche overlap measure using the index <i>D</i> between evaluated <i>Phrynosoma</i> taxa.....  | 87  |
| Table 4.1 | Range of values of 19 bioclimatic variables for the current and last glacial maximum climatic conditions .....                        | 134 |

## LIST OF FIGURES

|            |  |     |
|------------|--|-----|
| Figure 2.1 | Sampling localities of <i>Dipodomys microps</i> .....  | 39  |
| Figure 2.2 | The genetic consequences based on the two proposed hypotheses .....  | 40  |
| Figure 2.3 | Climatic niche models for <i>D. microps</i> for current climatic conditions and climatic conditions of the last glacial maximum.....   | 41  |
| Figure 2.4 | Median-joining network of control region sequences for <i>D. microps</i> .....   | 42  |
| Figure 2.5 | Interpolated genetic indices across landscape for <i>D. microps</i> .....  | 43  |
| Figure 2.6 | Scatterplots for genetic indices, sample sizes, and latitude .....   | 44  |
| Figure 2.7 | Bayesian skyline plots and mismatch distribution of control region sequences for the entire dataset.....   | 45  |
| Figure 2.8 | Mismatch distributions for 12 sampling localities with sample size greater than or equal to 10 .....   | 46  |
| Figure 3.1 | General sample sites of <i>Phrynosoma platyrhinos</i> species complex .....  | 91  |
| Figure 3.2 | Maximum likelihood tree for combined ND4, tRNA and cytB sequences for 27 unique haplotypes of <i>P. platyrhinos</i> species complex.....   | 92  |
| Figure 3.3 | Median-joining network of concatenated ND4 and tRNA sequences of <i>P. platyrhinos</i> .....   | 93  |
| Figure 3.4 | Distribution of major nested clades identified from the median-joining network.....  | 94  |
| Figure 3.5 | Bayesian skyline plot derived from concatenated ND4 and tRNA sequences of <i>P. platyrhinos</i> .....  | 95  |
| Figure 3.6 | Scatter plot for the first two principal components for the persistent and expanding groups of <i>P. platyrhinos</i> .....   | 96  |
| Figure 3.7 | Bayesian skyline plots and mismatch distribution of control region sequences for the entire dataset.....   | 97  |
| Figure 3.8 | Climatic niche models for the persistent and expanding groups of <i>P. platyrhinos</i> .....   | 98  |
| Figure 4.1 | Difference between the CCSM and MIROC models for the western region of North America.....  | 138 |
| Figure 4.2 | Ecological niche models for <i>Perognathus parvus</i> created using 19 bioclimatic variables and with and without elevation.....   | 140 |
| Figure 4.3 | Histograms of suitable elevation extracted from the ecological niche models of <i>Perognathus parvus</i> .....   | 142 |
| Figure 4.4 | Schematic picture showing different approaches when projecting a model outside the calibration range (extrapolation). .....  | 143 |
| Figure 4.5 | The clamping values for ENMs of <i>Masticophis taeniatus</i> using different extents of variables.....   | 144 |
| Figure 4.6 | A set of response curves of 19 bioclimatic variables for <i>Uta stansburiana</i> produced by Maxent.....   | 145 |
| Figure 4.7 | Ecological niche models and associated clamping values for <i>Uta stansburiana</i> for the climatic conditions of the last glacial maximum using two different sets of variables ..... | 146 |

|             |   |     |
|-------------|---|-----|
| Figure 4.8  | Representation of a niche shift in a hypothetical species from time T1 to time T2 .....   | 147 |
| Figure 4.9  | Principal components scores derived from bioclimatic niches occupied by four species of rodents during the last glacial maximum and at present time ..... | 148 |
| Figure 4.10 | Ecological niche models for four species of rodents using the LGM reconstructions and their evaluation using fossil records.....                          | 149 |
| Figure 4.11 | Two scenarios of a hypothetical species history leading to the same genetic signal .....  | 151 |

## CHAPTER 1

### INTRODUCTION

In the face of ongoing and projected future warming trends, a lot of research has recently focused on predicting how species respond to climatic and environmental changes. These predictions are usually accomplished using a methodology called transferal modeling. Transferal modeling uses information on environmental requirements delimiting species distributions to reconstruct species suitable habitat (referred to as ecological niche) which is then projected on different climate change scenarios. One of the assumptions of the transferal modeling methodology is 'niche conservatism', a property of a species to maintain the same ecological niche through time. Under niche conservatism, the species will track its ecological niche across space as climate changes, resulting in range shifts, expansions, and contractions. If the species ecological niche does not remain conserved and shifts through time instead, the species might not respond to climatic changes in a predictable fashion. Niche shift versus niche conservatism can be either studied by direct observations of species responses to ongoing and simulated climatic changes, or by reconstructing species responses to past climatic changes.

In the second and third chapter, I explore, whether the niches of two species (one mammal and one reptile) remained conserved throughout the climatic changes of the latest glacial – interglacial cycle, in particular between the last glacial maximum (LGM) and present time. These two time periods are ideal for studying species responses to climate change as they represent two extremes of global climate: the LGM is one of the

coldest periods of the late Pleistocene while the current climate represents warm climate of an interglacial period. I use transferal modeling to reconstruct the species ranges under the assumption that their niches remained conserved between the LGM and present time. I then use DNA sequence data derived from samples spanning the species range to infer the species past geographic and demographic processes. For example, populations that persisted in place for quite some time will exhibit a different genetic signal than populations that expanded into the region relatively recently, in response to a major climatic event. If the species niches indeed remained conserved, the transferal models should infer the same geographic processes (i.e. range expansion, contraction, or shift) as the genetic data. If the species experienced niche shifts, however, the results from transferal models and genetic data will be incongruent.

In the last chapter, I review several other challenges and uncertainties associated with transferal modeling that I encountered during my research and that can negatively affect resulting models. Thus, my research will help us to better understand species responses to past climatic changes and ultimately also help us to improve our predictions of species responses to future changes.

## CHAPTER 2

# NICHE SHIFT IN SITU IN RESPONSE TO THE WARMING CLIMATE AT THE END OF THE LAST GLACIAL MAXIMUM: INFERENCE FROM GENETIC DATA AND NICHE ASSESSMENTS IN THE CHISEL-TOOTHED KANGAROO RAT (*DIPODOMYS MICROPS*)

### Abstract

During Pleistocene glacial-interglacial cycles, species are often assumed to have shifted ranges via tracking the climatic niche to which they have been adapted, rather than remaining in place and adapting to a changing environment. However, species that exploit diverse types of habitats might instead be capable of shifting their niches as climates change, which would allow them to persist in place through time. I evaluate whether this kind of response occurred in the Chisel-toothed Kangaroo Rat (*Dipodomys microps*), which currently utilizes two distinct habitats: the low-elevational saltbush (*Atriplex*) and mid-elevational blackbrush (*Coleogyne*) communities in the deserts of western North America. I modeled how the species range would have changed between the last glacial maximum (LGM) and present time if the climatic niche of the species remained identical (conserved) between the two time periods. I then used mitochondrial DNA (mtDNA) assessments of demographic parameters to evaluate whether *D. microps* exhibits signals of distributional stability or of recent geographic changes. The climatic models imply that if the species inhabited the same climatic niche during the LGM as it does today, the range of *D. microps* would have changed significantly: it would have persisted primarily within the southern, warm Mojave Desert and expanded northwards into the cold deserts of the Great Basin and Columbia Plateau



only after the LGM. Conversely, the mtDNA assessment revealed signals of population persistence within the current distribution of the species throughout at least the latest glacial-interglacial cycle. My results suggest that *D. microps* did not track its climatic niche during late Pleistocene oscillations, but rather met the challenge of a changing environment by shifting its niche sufficiently to retain large portions of its current distribution. I speculate that this kind of response to fluctuating climate was possible because of 'niche-drifting', an alteration of the species' realized niche due to plasticity in various (physiological, morphological, behavioral) characters.

## Introduction

When climate changes, populations may go extinct, move to locations that remain suitable, or stay in place and adapt to novel environmental conditions (Barnosky *et al.* 2003; Parmesan 2006; Parmesan *et al.* 1999; Pounds *et al.* 1999; Vrba 1992). Local extinction often is encountered when climate change generates environmental conditions that exceed the existing tolerance limits of the organism if its ecological niche remains conserved (identical) through time (Warren *et al.* 2008; Wiens & Graham 2005). During the late Pleistocene, the desert regions of the western United States (Fig. 2.1) experienced pronounced and repeated climate change (Spaulding 1990a; Thompson 1990). During the latest glacial period, lower temperatures and higher precipitation caused downward (latitudinal or elevational) shifts in plant assemblages, resulting in a general reduction of size and continuity of desert habitats (Thompson & Anderson 2000; Thompson & Mead 1982). This shift was most extreme during the last glacial maximum

(LGM), about 21,000 years before present (Harrison 2000). During this time, many kinds of desert organisms that are currently widespread throughout the western deserts were extirpated from the northern regions (i.e. the Great Basin Desert and Columbia Plateau) during the LGM and persisted within southern refugia (located within the Mojave or Sonoran deserts) where environmental conditions remained suitable (e.g. Jones 1995; Mulcahy 2008). These taxa expanded into the northern regions only when desert habitats were re-established as climate warmed after the LGM (Britten & Rust 1996; Hockett 2000; Jones 1995).

Some populations, however, persisted without substantial range shifts during major environmental changes, perhaps because of an ability to adapt (i.e. shift their niches) to novel conditions. For example, several species of woodrats (*Neotoma* spp.) were documented to have responded to cooling climate during glacial periods by increasing their body mass (Bergmann's Rule) instead of shifting their distributions to lower latitudes or altitudes as would be predicted if their climatic niches remained conserved (Smith & Betancourt 2003). Consequently, one could ask, why are some taxa or populations capable of adapting to novel conditions (Smith & Betancourt 2003) while others respond to an environmental change by tracking the conditions to which they are already adapted (Parmesan *et al.* 1999; Pounds *et al.* 1999)?

Vrba (1992) proposed that taxa capable of utilizing alternative environments are most likely to persist through environmental changes. In general terms, a species would be able to shift their niche and maintain populations under changing climates if (1) it exhibits plasticity in relevant physiological, phenological, morphological, or behavioral

traits that allows its realized niche to expand or shift – a process I call ‘niche drifting’; or (2) it’s fundamental niche is modified via evolutionary change, which I call ‘niche evolution’. Understanding which taxa or populations are capable of niche shifts (both niche drifting and niche evolution) is critical to predicting their likely responses to ongoing and future climate change.

The Chisel-toothed Kangaroo Rat (*Dipodomys microps*) inhabits four desert regions of western North America (Mojave Desert, Colorado Plateau, Great Basin Desert, and Columbia Plateau; 1.1) and exhibits two distinct ecotypes that differ in certain ecological, behavioral, and physiological traits (Csuti 1979). Throughout most of its range, *D. microps* occupies low-elevation saltbush habitat (dominated by several species of *Atriplex* - *A. confertifolia*, *A. canescens*, *A. polycarpa*, or *A. spinifera*; herein referred to as *Atriplex* habitat) where it is mostly folivorous, with 80% of its diet consisting of *Atriplex* (Csuti 1979; Kenagy 1973). In the southern parts of its range, within the Mojave Desert, the species also occupies mid-elevational blackbrush habitat (*Coleogyne ramosissima*; herein referred to as *Coleogyne* habitat), where it is generally granivorous and its diet comprises a variety of different plants (Csuti 1979). Accordingly, I hypothesize that species such as *D. microps* that exploit two very distinct habitats might be capable of niche shifts that allow them to persist in large portions of their ranges during major episodes of climate change.

I test the null hypothesis (Fig. 2.2A) that the climatic niche of *D. microps* remained conserved through time and that this species responded to the changing climate of the late Pleistocene by tracking the climatic niche to which it has been adapted. In

particular, this hypothesis predicts that *D. microps* persisted in glacial refugia within the southern deserts during the LGM and expanded northward after the LGM, in concert with many other desert taxa (Britten & Rust 1996; Hockett 2000; Jones 1995; Mulcahy 2008). Alternatively, given its rather broad habitat preferences, the 'niche shift' hypothesis predicts that *D. microps* might have been able to exploit conditions that were colder and wetter than present and persist throughout its range during the LGM (Fig. 2.2B).

To test my hypotheses, I use climatic niche modeling to estimate the distributional extent of the climatic niche of *D. microps* during the LGM under the assumption that the climatic niche between the LGM and present time remained conserved. From these models, I identify general areas that the species occupies today but would have been unsuitable during the LGM. I contrast these models with assessments of genetic data represented by mitochondrial DNA sequences sampled across the current range of *D. microps*. Mitochondrial DNA exhibits a relatively high mutation rate, which in combination with its small effective population size (four times smaller than nuclear DNA), makes it an appropriate genetic marker for tracking recent population histories including demographic and geographic changes (Avice 2000; Moore 1995; Zink & Barrowclough 2008). I contrast the genetic assessments with the climatic niche models in order to evaluate whether populations inferred by models as being nonexistent during the LGM truly exhibit a genetic signal of a recent expansion and whether those modeled as being geographically persistent exhibit a genetic signal of population stability.

‘Leading edge’ and ‘phalanx’ colonization are two general models for population expansion that were developed based on a set of predictions for the sort of phylogeographic signature that should be exhibited by a species that expanded its range (Hewitt, 1996). Based on the ‘leading edge’ model, newly invaded areas are colonized by a subset of individuals (and therefore a subset of genotypes) from the colonization front and this front inhibits establishment by later migrants. This type of expansion results in a decrease of genetic (i.e. nucleotide and haplotype, see Methods) diversity within newly established populations and a decrease in genetic variation among populations in the direction of the expansion (Fig. 2.2A). Leading edge colonization is typical for rapid, large-scale geographic expansions exhibited by numerous taxa after the last glacial maximum (Austerlitz *et al.* 1997; Excoffier *et al.* 2009; Hewitt 2000; Hewitt 1996; Kerdelhue *et al.* 2009). Some taxa, however, exhibit genetic signals of the ‘phalanx’ colonization (Hewitt 1996). Under this model, new populations at the expansion front are formed from a more even mixture of ancestral populations, resulting in only marginal decrease in local genetic diversity (Fig. 2.2A). While genetic diversity within populations remains higher under this model, genetic variation among expanding populations will be low as in the ‘leading edge’ model, because the expanding populations will consist of similar groups of genotypes (Fig. 2.2A). The ‘phalanx’ model of colonization has been documented in populations with high population densities, high gene flow, wide colonization front, and during slower expansions (Hewitt 1996; Schmitt *et al.* 2005). In both types of expansion models, stable populations exhibit higher genetic diversity within populations due to the accumulation of mutations

through time and usually higher genetic variation among populations due to geographic structuring (Hewitt 2000; Hewitt 1996). Species experiencing a recent range expansion also should exhibit a genetic signal of a recent increase in overall effective population size (Drummond *et al.* 2005).

If populations of *D. microps* experienced a niche shift and retained much of their current geographic range throughout the latest glacial-interglacial cycle, the genetic signal of stable populations (high genetic diversity within and variation among) would be detected throughout the species range (Fig. 2.2B). I employ several phylogeographic and population genetic methods to test the two competing hypotheses outlined above (Fig. 2.2A, B) and infer the recent geographic and population history of *D. microps*.

## Methods

### Taxon sampling

I acquired tissue samples from 364 individuals of *D. microps* from 83 unique localities; localities closer than 10 km without an obvious physical barrier were pooled together for a total of 67 general localities (Fig. 2.1; Table 2.1; Appendix 1). Of these 364 individuals, 340 were sampled specifically for this study. Most of the captured animals were ear-clipped and released ( $N=315$ ), but some ( $N=25$ ) were euthanized following methods approved by the American Society of Mammalogists (Gannon *et al.* 2007) and the Institutional Animal Care and Use Committee Protocol R 0709-244, and were deposited in the mammal collection at the New Mexico Museum of Natural History (NMMNH). Fourteen tissue samples were requested from the Museum of Texas

Technical University and 10 samples were requested from the Museum of Vertebrate Zoology, University of CA (Appendix 1). These 10 samples (six of which were represented by skin clips) came from the two most southern localities at Joshua Tree National Park (locality 9) and Yucca Valley (locality 11) where *D. microps* might be currently extinct (Drost & Hart 2008).

As different analyses require different sampling strategies, I used the following subsets and modifications of the original dataset. For climatic niche models, I used all unique sampling localities resulting in 83 presence records for *D. microps*. For genetic analyses, I used three different datasets. For calculating overall genetic indices and for phylogeographic analyses that do not require large sample size per a locality, I used all 364 individuals combined (Network, Bayesian skyline plot, overall mismatch distribution, and diversity indices) or with each assigned to one of the 67 general localities (pairwise genetic distances). I refer to this as “full dataset”. For population genetic analyses that require multiple individuals per locality (mismatch distributions, diversity indices for individual localities, and the  $R_2$  test) I excluded all localities with sample size smaller than 10 which resulted in 12 sampling localities (165 samples), referred to as “10+ dataset”. Additionally, the nucleotide and haplotype diversity, and frequency of private haplotypes used in the landscape interpolation assessment were calculated for localities with sample size equal or larger than five, resulting in 29 locations and referred to as “5+ dataset”.

### Climatic niche models

I modeled the climatic niche of *D. microps* in order to approximate the species' current distribution and distribution during the LGM under the assumptions that (1) climate is an important factor driving the species distribution and (2) that the climatic niche of *D. microps* remained conserved between the LGM and present time. The climatic niches were reconstructed using the methodology of ecological niche modeling where environmental data are extracted from occurrence records (represented by geographic coordinates) and habitat suitability is evaluated across the landscape using program-specific algorithms (Elith *et al.* 2006). The current models were then projected on the climatic reconstructions of the LGM.

For occurrence records I used unique sampling localities and the current climate was represented by bioclimatic variables from the WorldClim dataset v. 1.4 (<http://www.worldclim.org/>; Hijmans *et al.* 2005). The bioclimatic variables are derived from monthly temperature and precipitation data, and represent biologically meaningful aspects of local climate (Jezkova *et al.* 2009; Waltari *et al.* 2007). From the original 19 variables, I excluded those that were highly correlated and used a total of 14 variables (for detailed methodology, see Jezkova *et al.*, in review). For environmental layers representing the climatic conditions of the LGM, I used ocean-atmosphere simulations (Harrison 2000) available through the Paleoclimatic Modelling Intercomparison Project (Braconnot *et al.* 2007). These reconstructions of the LGM climate are based on simulated changes in concentration of greenhouse gases, ice sheet coverage, insulation, and topography (caused by lowering sea levels). Of more than 15



available climatic models, I used the following two: Community Climate System Model v. 3 (CCSM; Otto-Bliesner *et al.* 2006) with a resolution of 1°, and the Model for Interdisciplinary Research on Climate v. 3.2 (MIROC; Hasumi & Emori 2004) with an original spatial resolution of 1.4° x 0.5° (Braconnot *et al.* 2007). The original climatic variables used in these models have been downscaled to the spatial resolution of 2.5 minutes under the assumption that changes in climate are relatively stable over space (high spatial autocorrelation) and were converted to bioclimatic variables (Peterson & Nyari 2008).

Climatic niche models were built in the software package MAXENT v. 3.2.1 (Phillips *et al.* 2006), a program that calculates relative probabilities of the species' presence in the defined geographic space, with high probabilities indicating suitable environmental conditions for the species (Phillips *et al.* 2004). I used the default parameters in MAXENT (500 maximum iterations, convergence threshold of 0.00001, and regularization multiplier of 1, 10,000 background points) with the application of random seed and logistic probabilities for the output (Phillips & Dudik 2008). I masked my models to the four regions where *D. microps* occurs (Mojave Desert, Great Basin, Columbian Plateau, and Colorado Plateau; Olson *et al.* 2001) because reducing the climatic variation being modeled to that which exists within a geographically realistic area improves model accuracy and reduces problems with extrapolation (Pearson *et al.* 2002; Randin *et al.* 2006; Thuiller *et al.* 2004). I ran 50 replicates for each model and an average model was converted to presence-absence maps using a minimum training presence threshold. I

used the receiver operating characteristic for its area under the curve (AUC) value to evaluate the model performance (Fielding & Bell 1997; Raes & ter Steege 2007).

### Laboratory methods

I isolated total genomic DNA from preserved heart, kidney, or ear tissue following the protocol for the DNeasy Extraction Kit (Qiagen Inc.). I amplified ca. 1000 base pairs (bp) of the mitochondrial control region using genus-specific primer L15926DIOR (GTATAAAAATTACTCAGGTCTTGT) and an universal primer H651 (Kocher *et al.* 1989). Amplifications were accomplished in 12.5 µL reactions using Takara Ex Taq Polymerase Premix (Takara Mirus Bio, Inc.) followed by purification using ExoSap-IT (USB Corp.). Thermal cycling was accomplished at a 56°C annealing temperature in 40 cycles. I conducted double-stranded cycle sequencing using fluorescence-based chemistry (BigDye Terminator v. 3.1 Cycle Sequencing Kit) and the genus-specific primers L15926DIOR and H651Lpen (TAACTGCAGAAGGCTAGGAC). The electrophoresis was conducted on an ABI Prism 3130 automated sequencer (Applied Biosystems, Inc.). The sequences were aligned using SEQUENCHER v. 4.6 (Gene Codes Corp.) and verified manually. The DNA from the six museum skins of *D. microps* was isolated following the protocol for the DNeasy Extraction Kit with the following modifications: 40 µl per sample of proteinase K (double the regular value), 20 µl dithiothreitol (DTT) per sample added prior to incubation, and an extended incubation period (three days). I amplified and sequenced the targeted region using the thermal profiles described above with five overlapping genus-specific primer pairs about 250 bp each (the primer sequences can be provided upon request). Amplifications were accomplished in 25 µl reactions using

GoldTaq (Applied Biosystems, Inc.) followed by purification using ExoSap-IT (USB Corp.). The final DNA segment for all samples included 902 bp of mtDNA control region sequence.

### Genetic analyses

For the full dataset I calculated the number of variable sites, parsimony informative sites and a net uncorrected  $p$ -distance in MEGA v. 4 (Tamura *et al.* 2007), and haplotype diversity ( $H_d$ ) and nucleotide diversity ( $\pi_d$ ) in ARLEQUIN v. 3.11 (Excoffier *et al.* 2005). Preliminary phylogenetic analyses of the mitochondrial DNA (mtDNA) dataset showed that *D. microps* exhibits shallow structuring. Accordingly, I used a median-joining network (Bandelt *et al.* 1999) to reconstruct relationships among the mtDNA sequences (referred to as haplotypes) as networks represent an ideal phylogeographic method when levels of genetic divergence are low, multifurcations occur, and ancestral haplotypes are still present in the populations (Crandall & Templeton 1996). Network analysis was conducted for the full dataset in the program NETWORK v. 4.5.1.6 (Bandelt *et al.* 1999) with transversions weighted twice as high as transitions (as recommended in the NETWORK manual), and with the maximum parsimony (MP) option employed to remove excessive links from the network (Polzin & Daneshmand 2003).

I calculated four genetic indices that can elucidate species geographic and demographic processes, such as a potential recent northward expansion of *D. microps* from southern refugia. Two indices reflect genetic variation (genetic distances among populations and frequency of private haplotypes within populations) and two indices represent genetic diversity (nucleotide and haplotypes diversity; Fig. 2.2). The genetic

distances are represented by mismatch distances between sequences from neighboring sampling sites calculated in the program ALLELES IN SPACE (Miller 2005) for the full dataset (67 locations). These pairwise genetic distances were assigned to mid-points between sampling sites using the Delaunay triangulation-based connectivity network (Miller *et al.* 2006). I used residual genetic distances derived from the linear regression of genetic versus geographical distance to account for correlation between these two distances (Manni *et al.* 2004; Miller *et al.* 2006). The residual genetic distances were imported into ARCGIS 9.2 (ESRI, Redlands, CA) and interpolated across uniformly-spaced 2.5 minute grids (~5 km) using the inverse distance weighted interpolation procedure (Watson & Philip 1985) in the Spatial Analyst extension. I restricted the interpolations to the geographic range of *D. microps*, approximated using the climatic niche model for current climatic conditions.

I calculated nucleotide diversity in ARLEQUIN as an average number of nucleotide differences per site among sequences in each sampling locality and haplotype diversity as a probability that two randomly chosen haplotypes in a population are different, regardless of genetic distance between them. These two indices of genetic diversity can reveal slightly different signals of population history (see Discussion). The private haplotype frequency was calculated by hand as a frequency of DNA sequences (haplotypes) in a population that are unique (do not occur in any other population). Recently expanded populations should exhibit low frequency of private haplotypes as the same haplotypes expand across large geographic areas and consequently occur in multiple populations (Fig. 2.2A). I calculated nucleotide and haplotype diversity and

frequency of private haplotypes for the 5+ dataset, and confirmed that sample size and the diversity indices are not positively correlated by conducting a correlation analysis in SYSTAT v. 12 (Hilbe 2008). The three diversity indices were imported into ArcGIS and interpolated across landscape as described above for genetic distances.

I analyzed past demographic changes of *D. microps* in order to detect any significant changes in population sizes through time. For example, if the species experienced a range expansion after the LGM, I would detect a significant increase in the overall effective population size dating to sometime after the LGM. I used the Bayesian skyline plot (BSP) coalescent model (Drummond *et al.* 2005) implemented in the program BEAST v. 1.5.3 (Drummond & Rambaut 2007) that generates plots of estimated posterior distribution of the effective population size through time (Drummond *et al.* 2005). Following assessment in MRMODELTEST, I selected the substitution model GTR+I+ $\Gamma$ , along with a strict molecular clock (after assessing clock-like behavior), five skyline groups, and a 1-year generation time (Gummer *et al.* 1997; Thomas *et al.* 1990). Since mutation rates for *D. microps* control region are unavailable from the literature, I applied a broad range of rates (1.5 and 6 % /lineage/million years) that have been used for control region sequences of other small mammals (Galbreath *et al.* 2009). I conducted several independent Markov Chain Monte Carlo (MCMC) runs of 80 million generations with sampling every 8000 generations and a burn-in of 10%. For final analysis three final MCMC runs were combined using LOGCOMBINER (distributed with BEAST). I checked convergence (effective sample sizes > 200) and visualized the median

and 95% highest posterior density intervals using TRACER v. 1.5 (Rambaut & Drummond 2007).

I further tested for demographic changes using mismatch distribution analyses under the sudden expansion model (Rogers & Harpending 1992; Schneider & Excoffier 1999) in ARLEQUIN for all sequences combined (n=364) and for the 10+ dataset. Mismatch distribution analysis calculates the number of pairwise differences among sequences and identifies populations with star-like phylogeny, a structure typical for populations that experienced a sudden expansion (Excoffier 2002; Excoffier *et al.* 2009). The star-like phylogeny will appear as unimodal distribution of pairwise differences with the peak corresponding to a lower number of pairwise differences in younger expansions and higher number in older expansions (Excoffier 2002). This analysis results in a multimodal graph if a population is in demographic equilibrium, subdivided or in decline (Rogers & Harpending 1992; Slatkin & Hudson 1991). A significant sum of squared deviation ( $SSD$ ;  $P < 0.05$ ) and Harpending's raggedness index ( $r$ ;  $P < 0.05$ ) indicate the rejection of the null hypothesis of sudden expansion. If the sudden expansion model was not rejected, I calculated population expansion parameters  $\tau$  (time since expansion),  $\theta_0$  (population size prior to expansion), and  $\theta_1$  (final population size). The parameter  $\tau$  was used to calculate time in years since expansion (Rogers & Harpending 1992) using the same mutation rates and generation time as above.

To further evaluate whether the species experienced an expansion I used Ramos-Onsins and Rozas'  $R_2$  test (Ramos-Onsins & Rozas 2002) in DnaSP v. 5.1 (Librado & Rozas 2009) for the full dataset and the 10+ dataset (Table 2.2). The  $R_2$  test is based on the

differences between the number of singleton mutations (mutations that occur in only one individual) and the average number of nucleotide differences (Ramos-Onsins & Rozas 2002) and significantly low ( $P \leq 0.02$ )  $R_2$  values indicate population growth.

## Results

### Ecological niche models

The model of current conditions (Fig. 2.3A) captures well the known distribution of *D. microps* but with some over-predictions of occurrence within the eastern part of the Colorado Plateau and on the Snake River Plain in southern Idaho. These areas, however, might currently be unoccupied by *D. microps* due to non climate-related factors such as insufficient time for colonization and establishment of more widespread populations (these two areas exhibit signals of a recent expansion, see below), recent habitat destruction (especially in the Snake River Plain) or possibly competition with other species (e.g. congeneric Ord's kangaroo rat, *D. ordii*). The paleo-models (Fig. 2.3B, C) indicate absence of a suitable climatic niche in the Great Basin and on the Columbia Plateau during the LGM. A suitable climatic niche is predicted to have been available within the Mojave Desert, but it is modeled as being fragmented and more restricted in distribution compared to present.

The mean AUC value for the current ecological niche model was only moderately high (0.82); AUC values range from 0.5 for a random prediction to 1 for perfect prediction (Winker *et al.* 2007). As pointed out by Lobo *et al.* (2008) and Peterson *et al.* (2008), the AUC values are greatly dependent on the number and variation of the

background points that are available. Because I masked my models to four ecoregions, the number of background points was limited. I believe that the moderately high AUC values in the masked models are likely the result of the methodological limitations, as the AUC values were high ( $<0.98$ ) when I ran the same models with no mask.

### Genetic analyses

Of the 902 bp of control region, 205 characters are variable and 160 are parsimony informative. The overall haplotype diversity ( $0.9957 \pm 0.0008$ ) and nucleotide diversity ( $0.0242 \pm 0.0118$ ) are high, and the net pairwise uncorrected  $p$ -distance is 2.4%. The median-joining network (Fig. 2.4) collapsed the 364 sequences into 243 haplotypes. The number of mutations between haplotypes ranges from 1 to 14 and the number of sequences belonging to any given haplotype ranges from 1 to 10. Most haplotypes are restricted to a single locality and of the 13 haplotypes shared among localities (Fig. 2.4), three are shared between *Atriplex* and *Coleogyne* localities. The median-joining network shows an overall star shape, with missing (extinct or unsampled) or low frequency haplotypes within the central parts of the network and with a burst of haplotypes separated from the center by up to 40 mutational steps. Nucleotide diversity within populations ranges from 0.00056 to 0.02222 (Fig. 2.5A) and there is no significant correlation between sample size and nucleotide diversity ( $r = -0.223$ ,  $p = 0.244$ ; Fig. 2.6). The nucleotide diversity does not decrease in the northward direction, as would be expected under the leading edge colonization model ( $r = -0.148$ ,  $p = 0.443$ ; Fig. 2.2, Fig. 2.5A, Fig. 2.6). Nucleotide diversity is relatively high throughout the entire species range except the edge populations in the Colorado Plateau (northern Arizona), southern



Mojave Desert (southern CA), and northwestern Columbia Plateau (northeastern CA, southern OR and southwestern ID) regions. Haplotype diversity (Fig. 2.5B) ranges from 0.46 to 1.00 (where 1.00 represents populations with no redundant haplotypes) and is not significantly correlated with sample size ( $r = -0.260$ ,  $p = 0.174$ ; Fig. 2.6) or latitude ( $r = 0.229$ ,  $p = 0.232$ ; Fig. 2.6). Haplotype diversity is highest within the central parts of the species range and lowest within northwestern, southeastern, and southern edge localities (Fig. 2.5B).

The residual genetic distances range from -0.0209 to +0.0127 (Fig. 2.5C). The negative values indicate that the genetic distance between populations is lower than the average, given the geographic distances between them. The genetic distances among populations are relatively high throughout the entire range of *D. microps*, indicating genetic structuring and limited gene flow, inconsistent with recent northward expansion (Fig. 2.2). Low genetic distances are found within the peripheral areas and generally correspond with areas of low nucleotide and haplotype diversity. Sampling localities within the area covered with the pluvial Lake Bonneville during the LGM show intermediate genetic distances. The private haplotype frequency ranges from 0.41 to 1.00 (frequency of 1.00 represents populations where none of the haplotypes occurs in any other population), is high throughout most of the range (Fig. 2.5D), and is not correlated with sample size ( $r = -0.273$ ,  $p = 0.455$ ; Fig. 2.6) or latitude ( $r = 0.142$ ,  $p = 1.000$ ; Fig. 2.6).

The Bayesian skyline plots (BSPs; Fig. 2.7A upper plot with a mutation rate 1.5%/lineage/million years (Mys) and lower plot with 6%/lineage/Mys) indicate an

increase in genetic diversity, then a period of relatively stable genetic diversity, followed by a recent decline. This signal is consistent (but see Discussion for possible caveats) with an increase in effective population sizes during the middle to late Pleistocene (ca. 750 Kya, upper plot; ca. 200 Kya, lower plot), followed by constant population sizes during the middle and late Pleistocene (starting ca. 500 Kya, upper plot; ca. 125 Kya, lower plot) and declines during the late Pleistocene or early Holocene (ca. 20 Kya, upper plot; ca. 5 Kya, lower plot).

The mismatch distribution (Fig. 2.7B) for the full dataset shows a unimodal distribution of the pairwise sequence differences consistent with a rapid population expansion. This expansion, however, was likely not recent as the median of the distribution corresponds to 27 pairwise differences. The small sum of squared deviation and raggedness index are not significantly different from a model of rapid population expansion ( $SSD=0.003$ ,  $P>0.05$ ;  $r=0.001$ ,  $P>0.05$ ; Table 2.2). This expansion was further supported by the values of  $\theta$ , as  $\theta_1 > \theta_0$  ( $\theta_1 = 105.045$ , 95% CI = 75.418-71515.045;  $\theta_0 = 0.004$ , 95% CI = 0.00-4.642). The beginning of population expansion based on the value of  $\tau$  ( $\tau = 25.127$ , 95% CI = 18.389-28.549) and the two mutation rates applied roughly correspond to 928,570 (95% CI = 679,564- 1,055,026) and 232,140 (95% CI=169,891-263,756) years ago (early-middle Pleistocene). Out of the 12 evaluated populations (Table 2.2; Fig. 2.8), five (localities 8, 15, 50, 61, 63) show a unimodal mismatch distribution. Localities 8, 15, and 61 have likely experienced a relatively recent population expansion while localities 50 and 63 probably experienced population growth long before the LGM. Recent population expansion is not supported by the  $R_2$

test for the full dataset and for none of the localities with the exception of locality 63 (Table 2.2). The sensitivity of the  $R_2$  test decreases with the time since the expansion and this decrease is even more rapid when sample size is small (Ramos-Onsins & Rozas 2002) which could explain why expansion was not detected by this test.

## Discussion

### Discordance between climatic niche models, genetic data, and the fossil record

In this study, I evaluated whether *D. microps* shifted its range in response to climate changes of the last glacial-interglacial transition, or alternatively, whether the species retained most of its range by adapting to the changing environment. The ecological niche models, built under the assumption that the climatic niche of *D. microps* has remained conserved between the LGM and present day, indicate that the climatic niche within the Great Basin and Columbia Plateau would have been unsuitable to the species during the LGM and that the suitable climatic niche within the Mojave Desert would have been reduced and fragmented. These results imply that if the climatic niche of the species was conserved, *D. microps* would have persisted within the Mojave Desert during the LGM and experienced a substantial expansion northwards after the LGM. My climatic models are in agreement with reconstructions of the LGM environment from packrat midden and pollen records (Spaulding 1990b; Spaulding *et al.* 1983; Thompson 1990), which indicate that during the LGM, the low elevation *Atriplex* habitats within the Great Basin and Columbia Plateau that currently represent the prime habitat for *D. microps* were covered with large pluvial lakes (Grayson 1993; Reheis 1999) or replaced

by an assemblage of plants currently found in higher elevations, such as *Artemisia* (sage), *Chrysothamnus* (rabbitbrush), and various woodland species (e.g. Utah juniper, *Juniperus osteosperma*; Single-leaf Pinyon, *Pinus monophylla*) (Grayson 1993; Spaulding 1990a; Spaulding 1990b; Spaulding *et al.* 1983; Thompson 1990). *Atriplex*-dominated communities were not established until the late Pleistocene or even the beginning of Holocene, when the climate warmed, plant communities shifted upwards and northwards, and the large pluvial lakes desiccated (Spaulding 1990b; Thompson 1990). Such vegetation shifts imply that environmental conditions during the LGM within the northern parts of the current range of *D. microps* were quite different from those that the species experiences today. The response of *Coleogyne* to the latest climate changes is less well understood (Summers *et al.* 2009) but it is believed that *Coleogyne* could have persisted mostly *in situ* (Wells 1983) and therefore could have maintained suitable niche for *D. microps* within portions of the Mojave Desert.

Despite these environmental changes, none of the genetic analyses supported post-LGM range expansion from the Mojave Desert into the Great Basin and Colombia Plateau. I did not detect the decrease in nucleotide or haplotype diversity in a northward direction (Fig. 2.5A, B; Fig. 2.6) that would be predicted under the ‘leading edge’ model of post-LGM range expansion (Fig. 2.2). High nucleotide and haplotypes diversity is usually a result of population stability as mutations are accumulated over time, but can also be maintained under the ‘phalanx’ model of population expansion (Fig. 2.2). Relatively high nucleotide diversity can also be observed within expanding areas when previously isolated (and therefore quite divergent) populations come to

secondary contact, such as after expansion from multiple refugia (Hewitt 1999). In such cases, however, I would observe lower levels of haplotype diversity within the areas of secondary contact in comparison to the stable areas, because of the reduced number of individual haplotypes coming from each of the colonization fronts.

I detected high genetic distances among populations (Fig. 2.5C) and high frequency of private haplotypes within populations (Fig. 2.5D), suggesting geographic structuring typical for long term population persistence with limited gene flow (Kerdelhue *et al.* 2009). Such signal contradicts the 'leading edge' as well as the 'phalanx' models of population expansion, where certain genotypes expand across large areas in the direction of the expansion, promoting gene flow, and resulting in decrease of genetic distances among populations and frequency of private haplotypes (Excoffier *et al.* 2009; Kerdelhue *et al.* 2009). Low genetic distances were only detected within the extreme edges in the northern, southern, and southeastern parts of the species range (localities 2, 8, 9, 15, 61), in localities that also exhibit low haplotype and nucleotide diversity. These areas could indeed be recently colonized, or might represent pockets of isolated populations that exhibit a signal of a population bottleneck (Avice 2000). More detailed sampling and analyses would be necessary to reveal the history of these edge populations.

The demographic analyses (BSP and mismatch distributions) revealed constant genetic diversity through at least the last several thousand years (Fig. 2.7A and B), which implies no significant post-LGM increase in effective population sizes that would be expected if the species experienced a range expansion. The estimate of effective

population size is confounded with genetic structuring which means that highly structured populations with low gene flow will exhibit larger genetic diversity (and therefore seemingly higher effective population size) than panmictic populations (Drummond *et al.* 2005). Because *D. microps* exhibits pronounced geographic structuring (as apparent from the patterns of genetic diversity; Fig. 2.5C), the assumption of panmixia was violated in both demographic analyses. Accordingly, I also performed mismatch distributions for the 12 sampling locations that had adequate sample size. Samples were collected for each of these localities from a small geographic area (less than 10 km) with no obvious barriers and should approximate a panmictic population. The resulting graphs (Fig. 2.8) further support my conclusions that most populations throughout the species range did not experience a recent expansion. In summary, all genetic analyses support the hypothesis that *D. microps* has not experienced a post-LGM northward expansion, but rather was able to retain most of its range despite pronounced climatic and environmental changes of the late Pleistocene.

There is no fossil record for *D. microps* dating back to the LGM that would corroborate my interpretation of the genetic data that the species persisted throughout the northern parts of its current range. However, an abundant late Pleistocene and Holocene fossil record spanning the last 11,300 years is available from the Homestead Cave in Northeastern Utah. This fossil record documented fluctuations in numbers of *D. microps* coinciding with climate oscillations throughout the Holocene (Grayson 2000). At this locality, the species fossil record was quite rare at the end of the Pleistocene and beginning of the Holocene, with a subsequent increase and peak in relative abundance

during the middle Holocene, presumably reflecting the warming climate that promoted the increase of *Atriplex* in the area. As the warm middle Holocene ended, fossils of *D. microps* declined in relative abundance again (Grayson 2000). My BSP (Fig. 2.7) could indeed be consistent with a decrease in effective population sizes during the late Holocene but my general conclusions do not support the increase in abundance during early and middle Holocene. I speculate that apparent low numbers of *D. microps* recovered from the oldest strata might result from the geographic position of the cave. Homestead Cave is situated in the middle of the pluvial Lake Bonneville (near my locality 66) and therefore might not have been colonized until after the water level receded. *Dipodomys microps* might therefore have been only locally but not regionally rare within the Great Basin at the beginning of the Holocene.

#### Niche drift or niche evolution?

Empirical and experimental studies show that species often respond individualistically to environmental change (Colautti *et al.* 2010; Graham *et al.* 1996; Parmesan 2006 and citations within; Rowe *et al.* 2010). Many taxa expand and retract their ranges in response to an oscillating climate, as implied from the fossil record (Grayson 2000; Hockett 2000) as well as from direct observations of range shifts caused by recent human-induced climate changes (Moritz *et al.* 2008; Parmesan *et al.* 1999). Some taxa have showed plastic responses in morphological, phenological, or physiological characters that allow them to meet the challenge of changing environment *in situ* and thus retain large portions of their current geographic distributions (Gibbs & Breisch 2001; Huppopp & Huppopp 2003; Menzel 2000; Post *et al.* 1999; Smith &

Betancourt 2003). Such niche responses resulting from an alteration of the realized niche (Jackson & Overpeck 2000; Rodder & Lotters 2009) do not require evolutionary adaptation. As such, here I defined a change in realized niche space as a 'niche drifting' in order to differentiate it from 'niche evolution', a change in the fundamental niche of a species. Niche evolution incorporates a shifting or broadening of the niche to meet new environmental conditions through evolutionary adaptation (Davis & Shaw 2001; Davis *et al.* 2005; Reale *et al.* 2003; Urban *et al.* 2007). Above, I demonstrate that the climatic niche of *D. microps* has not been conserved between the LGM and present time but I have not yet addressed the possible mechanisms that may have allowed the species to adapt to changing climate. Thus I ask the question, has the niche of *D. microps* drifted or evolved?

In a previous study, Csuti (1979) demonstrated that *D. microps* currently exhibits a generalized genotype that allows exploitation of the *Atriplex* and *Coleogyne* habitats. In the lab, animals from one habitat were able to adjust behaviorally and physiologically to exploit food resources from the other habitat. Despite this general plasticity, however, the descendants of the *Atriplex* population were still more efficient in shaving the *Atriplex* leaves than the descendants of the *Coleogyne* population, which suggests a certain level of evolutionary adaptation. The idea of a generalized genotype and therefore 'niche drift' seems to be a favored mechanism behind the adaptations to the developing *Atriplex*-dominated community after the last glacial period. Frequent glacial-interglacial fluctuations of the Pleistocene likely favored the retention of a generalized genotype and prevented fixation of a specialized genotype for any particular habitat.



Without the unpredictability conferred by climatic fluctuations, it is possible that differential selective pressure in *Atriplex* and *Coleogyne* habitats would induce genetic differences among populations and promote diversification and specialization.

Furthermore, adaptation to the *Atriplex* community is likely not a novel, but rather a reoccurring event within cycles of oscillating climate. Indeed, Kenagy (1973) suggested that it was adaptation to *Atriplex* that allowed *D. microps* to diverge from granivorous congeners in the Pliocene or beginning of the Pleistocene. This adaptation is apparent from the unique, chisel-shaped lower incisors of *D. microps* that are ideal for removing hypersaline surface tissues from *Atriplex* leaves (Hayssen 1991; Kenagy 1973). This morphology is conserved even in *Coleogyne* populations where *Atriplex* consumption is low or where *Atriplex* is not consumed at all. But why would a species specialize to a plant community typical of only short interglacial periods? Although *Atriplex*-dominated communities are likely prevalent only during interglacials (Spaulding *et al.* 1983), *Atriplex* itself was likely present throughout the entire Pleistocene. Indeed, *A. confertifolia*, was recorded in non-analogous plant communities within the Mojave Desert and the Great Basin during the last glacial period, mixed with plants currently found in higher altitudes (Spaulding *et al.* 1983). It is currently not clear whether *Atriplex* represented a dominant food source during glacial periods or whether *D. microps* utilized a wider variety of plant material as it currently does in the *Coleogyne* habitat. It is possible, however, that the permanent presence of *Atriplex* facilitated persistence of *D. microps* throughout much of its current range during the profound climate transition following the LGM.

## Conclusions

Pleistocene glacial-interglacial cycles have been documented to commonly induce shifts in species ranges through tracking the changing climates, rather than remaining in place and adapting to the changing environment (Parmesan 2006). In my study I show that *D. microps* was able to shift its niche and retain much of its current geographic range during at least the latest Pleistocene glacial-interglacial cycle. Such a niche shift in response to fluctuating climate and changing environment might have been possible because of the preservation of a generalized genotype and could have been enhanced by permanent availability of *Atriplex*, its dominant food source. Identification of factors that determine whether and to what extent taxa are able to adapt to a changing environment is crucial in our attempt to predict biotic responses to past, current, and future environmental changes.

**Table 2.1.** General sampling localities for *Dipodomys microps* depicted in Figure 1.1. *N* represents the sample size for each location. The geographic coordinates represent an approximate location; the exact coordinates for each sample are listed in Appendix 1.

| Locality<br>number | State | County         | Locality           | Lat   | Long    | <i>N</i> |
|--------------------|-------|----------------|--------------------|-------|---------|----------|
| 1                  | AZ    | Coconino       | House Rock Valley  | 36.49 | -111.96 | 11       |
| 2                  | AZ    | Mohave         | Wolf's Hole        | 36.89 | -113.56 | 15       |
| 3                  | CA    | Inyo           | Haiwee             | 36.15 | -117.98 | 7        |
| 4                  | CA    | Inyo           | Owens Valley       | 36.54 | -117.93 | 1        |
| 5                  | CA    | Inyo           | Saline Valley      | 36.84 | -117.91 | 1        |
| 6                  | CA    | Inyo           | Deep Spring Valley | 37.30 | -118.06 | 4        |
| 7                  | CA    | Kern           | Koehl Lake         | 35.29 | -117.88 | 12       |
| 8                  | CA    | Modoc          | Eagleville         | 41.16 | -119.99 | 13       |
| 9                  | CA    | Riverside      | Joshua Tree NP     | 33.99 | -116.23 | 8        |
| 10                 | CA    | San Bernardino | Boron              | 35.15 | -117.57 | 1        |
| 11                 | CA    | San Bernardino | Yucca Valley       | 34.15 | -116.47 | 2        |
| 12                 | CA    | San Bernardino | Superior Lake      | 35.23 | -117.03 | 2        |
| 13                 | CA    | San Bernardino | Clark Mountain     | 35.56 | -115.64 | 14       |
| 14                 | ID    | Owyhee         | Bruneau Canyon     | 42.76 | -115.76 | 4        |
| 15                 | ID    | Owyhee         | Murphy             | 43.19 | -116.64 | 10       |
| 16                 | NV    | Churchill      | Fallon             | 39.38 | -118.88 | 2        |

---

|    |    |           |                       |        |         |    |
|----|----|-----------|-----------------------|--------|---------|----|
| 17 | NV | Churchill | Hot Springs Mountains | 39.76  | -118.87 | 3  |
| 18 | NV | Clark     | Cottonwood Valley     | 36.00  | -115.45 | 2  |
| 19 | NV | Clark     | Kyle Canyon           | 36.26  | -115.48 | 14 |
| 20 | NV | Elko      | Tecoma                | 41.32  | -114.08 | 9  |
| 21 | NV | Esmeralda | Fish Lake Valley      | 37.78  | -118.10 | 1  |
| 22 | NV | Esmeralda | Tonopah               | 37.91  | -117.22 | 6  |
| 23 | NV | Esmeralda | White River Valley    | 38.41  | -115.12 | 3  |
| 24 | NV | Eureka    | Beowawe               | 40.52  | -116.51 | 4  |
| 25 | NV | Humboldt  | Battle Mountain       | 40.48  | -117.06 | 5  |
| 26 | NV | Humboldt  | Sulphur Landing       | 40.84  | -118.76 | 7  |
| 27 | NV | Humboldt  | Golconda              | 40.96  | -117.42 | 2  |
| 28 | NV | Humboldt  | 30 mi S Denio         | 41.65  | -118.44 | 2  |
| 29 | NV | Lander    | Monitor Valley        | 39.23  | -116.70 | 6  |
| 30 | NV | Lander    | Big Smoky Valley      | 39.39  | -116.93 | 1  |
| 31 | NV | Lincoln   | Coyote Spring         | 37.15  | -114.72 | 3  |
| 32 | NV | Lincoln   | Pahranagat NWR        | 37.23  | -115.10 | 8  |
| 33 | NV | Lincoln   | Kane Spring Valley    | 37. 30 | -114.58 | 18 |
| 34 | NV | Lincoln   | Tickaboo Valley       | 37.37  | -115.47 | 4  |
| 35 | NV | Lincoln   | Delmar Valley         | 37.52  | -114.87 | 2  |
| 36 | NV | Lincoln   | Rachel                | 37.71  | -115.81 | 6  |
| 37 | NV | Lincoln   | Dry Lake Valley       | 37.71  | -114.80 | 3  |
| 38 | NV | Lincoln   | Tempiute              | 37.74  | -115.58 | 1  |

---

---

|    |    |            |                       |       |         |    |
|----|----|------------|-----------------------|-------|---------|----|
| 39 | NV | Lincoln    | Dry Lake Valley       | 37.94 | -114.83 | 2  |
| 40 | NV | Lincoln    | Cave Valley           | 38.37 | -114.82 | 3  |
| 41 | NV | Lincoln    | Lake Valley           | 38.45 | -114.61 | 3  |
| 42 | NV | Mineral    | Marietta, Teels Marsh | 38.23 | -118.31 | 1  |
| 43 | NV | Mineral    | Tonapah Junction      | 38.27 | -118.11 | 3  |
| 44 | NV | Mineral    | Smith Valley          | 38.84 | -119.31 | 1  |
| 45 | NV | Nye        | Ash Meadows NWR       | 36.47 | -116.32 | 6  |
| 46 | NV | Nye        | Beatty                | 36.91 | -116.76 | 9  |
| 47 | NV | Nye        | Nevada Test Site      | 37.16 | -115.97 | 5  |
| 48 | NV | Nye        | Hot Creek Valley      | 38.21 | -116.18 | 2  |
| 49 | NV | Nye        | Railroad Valley       | 38.51 | -115.79 | 3  |
| 50 | NV | Nye        | Currant               | 38.79 | -115.53 | 12 |
| 51 | NV | Washoe     | Mullen Pass           | 39.87 | -119.62 | 13 |
| 52 | NV | Washoe     | Flanigan              | 40.19 | -119.85 | 3  |
| 53 | NV | Washoe     | Gerlach               | 40.75 | -119.29 | 1  |
| 54 | NV | White Pine | Snake Valley          | 38.63 | -114.27 | 7  |
| 55 | NV | White Pine | Spring Valley         | 38.94 | -114.42 | 2  |
| 56 | NV | White Pine | Spring Valley         | 39.14 | -114.50 | 7  |
| 57 | NV | White Pine | Steptoe Valley        | 39.23 | -114.84 | 1  |
| 58 | NV | White Pine | Spring Valley         | 39.49 | -114.39 | 1  |
| 59 | OR | Harney     | Fields                | 42.31 | -118.66 | 4  |
| 60 | OR | Harney     | 20 mi S Narrows       | 43.07 | -118.87 | 4  |

---

---

|    |    |            |                      |        |         |    |
|----|----|------------|----------------------|--------|---------|----|
| 61 | OR | Lake       | Alkali Lake          | 43.01  | -120.01 | 10 |
| 62 | UT | Boxelder   | Kelton               | 41.73  | -113.14 | 9  |
| 63 | UT | Juab       | Snake Valley         | 39.46  | -113.96 | 13 |
| 64 | UT | Millard    | White Sage Valley    | 39.10  | -112.85 | 9  |
| 65 | UT | Tooele     | Rush Valley          | 40.08  | -112.26 | 1  |
| 66 | UT | Tooele     | Puddle Valley        | 40. 90 | -112.95 | 4  |
| 67 | UT | Washington | Beaver Dam Mountains | 37.08  | -113.92 | 8  |

---

**Table 2.2.** Genetic indices for the full dataset and localities identified by numbers and sample sizes ( $N$ ). Shown are haplotype diversity ( $H_d$ ), nucleotide diversity ( $\pi_d$ ),  $R_2$  test (indicates significant population growth at  $P \leq 0.02$ ; shown as \*), and the mismatch distribution curves that are evaluated as unimodal, bimodal, or multimodal. The curves are tested for significant departure from the expansion model at  $P \leq 0.05$  (indicated as \*\*) using Sum of squared deviation ( $SSD$ ) and Harpending's raggedness index ( $r$ ).

| Locality     | $N$ | $H_d$ | $\pi_d$ | $R_2$  | $SSD$   | $r$     | Curve      |
|--------------|-----|-------|---------|--------|---------|---------|------------|
| 1            | 11  | 0.49  | 0.0010  | 0.178  | 0.262** | 0.151   | Bimodal    |
| 2            | 15  | 0.65  | 0.0025  | 0.100  | 0.06    | 0.125   | Bimodal    |
| 7            | 12  | 0.76  | 0.0101  | 0.143  | 0.088   | 0.145   | Multimodal |
| 8            | 13  | 0.53  | 0.0010  | 0.147  | 0.009   | 0.093   | Unimodal   |
| 13           | 14  | 0.85  | 0.0220  | 0.189  | 0.070** | 0.079** | Multimodal |
| 15           | 10  | 0.87  | 0.0023  | 0.132  | 0.004   | 0.04    | Unimodal   |
| 19           | 14  | 0.90  | 0.0215  | 0.136  | 0.049** | 0.092** | Multimodal |
| 33           | 18  | 0.87  | 0.0081  | 0.155  | 0.053   | 0.076   | Multimodal |
| 50           | 12  | 0.97  | 0.0183  | 0.139  | 0.014   | 0.031   | Unimodal   |
| 51           | 13  | 0.96  | 0.0192  | 0.133  | 0.023   | 0.038   | Multimodal |
| 61           | 10  | 0.64  | 0.0012  | 0.156  | 0.126   | 0.424   | Unimodal   |
| 63           | 13  | 0.96  | 0.0089  | 0.099* | 0.013   | 0.048   | Unimodal   |
| Full dataset | 364 | 0.99  | 0.0242  | 0.051  | 0.003   | 0.001   | Unimodal   |

**Figure 2.1.** Sampling localities for *Dipodomys microps*. The circle size and shading reflects the sample size, progressing from 1 to 18. Circles with black dots represent populations occupying *Coleogyne* habitats; empty circles represent populations occupying *Atriplex* habitats. The lighter shading in the inset represents the extent of the Mojave Desert, Sonoran Desert, Great Basin, Columbia Plateau, and Colorado Plateau ecoregions adapted from Olson *et al.* (adapted from Olson *et al.* 2001).

**Figure 2.2.** The genetic consequences based on the two proposed hypotheses: (A) The null hypothesis of range shift in *D. microps* in response to the changing climate at the end of the LGM. Stable populations that persisted throughout the LGM (top) harbor a large number of genotypes (represented by an array of different colors) generating high genetic diversity within population and high genetic variation among populations. Under the leading edge model of population expansion (bottom left), random subsets of genotypes are involved in the expansion, resulting in gradual loss of genotypes and therefore decrease in genetic diversity within populations in the direction of the expansion. Only the yellow genotypes are still present at the furthest front of the expansion. The white genotype is novel and has emerged during the expansion. Under the phalanx model (bottom right), more genotypes are involved in the expansion, maintaining moderate to high genetic diversity within the expanding populations. Under both models, genetic variation among expanding populations are low because expanding populations consist of similar groups of genotypes. (B) The alternative hypothesis of niche shift resulting in population persistence within the species range



despite climatic changes. Population persistence during the LGM generates high genetic diversity within population and high genetic variation among populations (top) that are preserved and detected in present populations (bottom). In this generalized illustration, genetic diversity represents both haplotype diversity ( $H_d$ ) and nucleotide diversity ( $\pi_d$ ; Fig. 2.5A,B) and genetic distances represent both mismatch distances and frequency of private haplotypes (Fig. 2.5C,D) but see Discussion for more details about each genetic index.

**Figure 2.3.** Climatic niche models for *Dipodomys microps* approximating the species range based on 14 bioclimatic variables representing current climatic conditions (A), and projected on two reconstructions of climatic conditions at the last glacial maximum (LGM) – CCSM (B) and MIROC (C) under the assumption of a conserved climatic niche. The green shading represents areas identified as suitable using the minimum presence threshold for logistic probabilities. Dark grey shading indicates areas identified as unsuitable. Blue areas correspond to the extent of pluvial lakes during the LGM (Raines *et al.* 1996). White dots represent current occurrence records of *D. microps* used to build the models.

**Figure 2.4.** Median-joining network of control region sequences for 364 samples of *Dipodomys microps*. Circle size reflects the number of individuals exhibiting a haplotype (smallest=1, largest=8). The length of connection lines between haplotypes is proportional to the number of mutational changes, with the shortest connection line

representing one mutational change. In (A), the colored circles represent haplotypes present in the 12 sampling localities with sample size equal or larger than 10. In (B), the dark grey circles represent haplotypes found in individuals from *Coleogyne* populations and white circles represent *Atriplex* populations.

**Figure 2.5.** Interpolated nucleotide diversity (A), haplotype diversity (B) pairwise genetic distances (C), and frequency of private haplotypes (D) across landscape for *Dipodomys microps* using a 2.5 minutes (ca. 5 km) grid size and restricted to the current climatic niche model of the species. The shading gradation progresses from green (lowest), yellow, brown to white (highest). The white circles indicate sampling localities used in each analysis; for nucleotide and haplotype diversity and frequency of private haplotypes, I used sampling localities with a sample size equal or larger than five while for genetic distances, I used all sampling sites.

**Figure 2.6.** Scatterplot between nucleotide diversity and sample size (A), nucleotide diversity and latitude (B), genetic distances and latitude (C), haplotype diversity and sample size (D), haplotype diversity and latitude (E), genetic distance among populations and latitude (E) and frequency of private haplotypes and latitude (F) based on mitochondrial control region sequences of *Dipodomys microps*.

**Figure 2.7.** (A) Bayesian skyline plots derived from control region sequences for 364 samples of *Dipodomys microps* using mutation rates 1.5%/lineage/Mys (upper plot) and 6%/lineage/Mys (lower plot). The x-axis shows the time progressing from right (oldest) to left (present). The y-axis shows an index of effective population size assuming a 1-year generation time. The black line is the median for genetic diversity and the grey area shows the 95% upper and lower highest posterior density limits. The arrows point to the timing of the last glacial maximum (~21 Kya). (B) Mismatch distribution analysis under the sudden expansion model.

**Figure 2.8.** Mismatch distributions for 12 sampling localities with sample size greater than or equal to 10.

Figure 2.1




Figure 2.2




Figure 2.3




Figure 2.4






Figure 2.5






Figure 2.6




Figure 2.7







Figure 2.8





## Bibliography

- Austerlitz F, Jung-Muller B, Godelle B, Gouyon PH (1997) Evolution of coalescence times, genetic diversity and structure during colonization. *Theoretical Population Biology* **51**, 148-164.
- Avice JC (2000) *Phylogeography: The History and Formation of Species* Harvard University Press, Cambridge, MA.
- Bandelt HJ, Forster P, Rohl A (1999) Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution* **16**, 37-48.
- Barnosky AD, Hadly EA, Bell CJ (2003) Mammalian response to global warming on varied temporal scales. *Journal of Mammalogy* **84**, 354-368.
- Braconnot P, Otto-Bliesner B, Harrison S, Joussaume S, Peterchmitt JY, Abe-Ouchi A, Crucifix M, Driesschaert E, Fichefet T, Hewitt CD, Kageyama M, Kitoh A, Laine A, Loutre MF, Marti O, Merkel U, Ramstein G, Valdes P, Weber SL, Yu Y, Zhao Y (2007) Results of PMIP2 coupled simulations of the Mid-Holocene and Last Glacial Maximum - Part 1: experiments and large-scale features. *Climate of the Past* **3**, 261-277.
- Britten HB, Rust RW (1996) Population structure of a sand dune-obligate beetle, *Eusattus muricatus*, and its implications for dune management. *Conservation Biology* **10**, 647-652.
- Colautti RI, Eckert CG, Barrett SCH (2010) Evolutionary constraints on adaptive evolution during range expansion in an invasive plant. *Proceedings of the Royal Society B-Biological Sciences* **277**, 1799-1806.
- Crandall KA, Templeton AR (1996) Applications of intraspecific phylogenetics. In: *New Uses for New Phylogenies* (eds. Harvey PH, Brown AJL, Smith JM, Nee S), pp. 81-99. Oxford University Press, Oxford, U.K.
- Csuti BA (1979) *Patterns of adaptation and variation in the Great Basin kangaroo rat (Dipodomys microps)*. University of California Publications in Zoology 111.
- Davis MB, Shaw RG (2001) Range shifts and adaptive responses to Quaternary climate change. *Science* **292**, 673-679.
- Davis MB, Shaw RG, Etterson JR (2005) Evolutionary responses to changing climate. *Ecology* **86**, 1704-1714.

- Drost CA, Hart J (2008) Mammal inventory of the Mojave Network parks; Death Valley and Joshua Tree National Parks, Lake Mead National Recreation Area, Manzanar National Historic Site, and Mojave National Preserve, p. 74. U.S. Geological Survey Open-File Report 2008-1167.
- Drummond AJ, Rambaut A (2007) BEAST: Bayesian evolutionary analysis by sampling trees. *Bmc Evolutionary Biology* **7**, 214.
- Drummond AJ, Rambaut A, Shapiro B, Pybus OG (2005) Bayesian coalescent inference of past population dynamics from molecular sequences. *Molecular Biology and Evolution* **22**, 1185-1192.
- Elith J, Graham CH, Anderson RP, Dudik M, Ferrier S, Guisan A, Hijmans RJ, Huettmann F, Leathwick JR, Lehmann A, Li J, Lohmann LG, Loiselle BA, Manion G, Moritz C, Nakamura M, Nakazawa Y, Overton JM, Peterson AT, Phillips SJ, Richardson K, Scachetti-Pereira R, Schapire RE, Soberon J, Williams S, Wisz MS, Zimmermann NE (2006) Novel methods improve prediction of species' distributions from occurrence data. *Ecography* **29**, 129-151.
- Excoffier L (2002) Reconstructing the demography of prehistoric human populations from molecular data. *Evolutionary Anthropology* **11**, 166-170.
- Excoffier L, Foll M, Petit RJ (2009) Genetic consequences of range expansions. *Annual Review of Ecology Evolution and Systematics* **40**, 481-501.
- Excoffier L, Laval G, Schneider S (2005) Arlequin (version 3.0): An integrated software package for population genetics data analysis. *Evolutionary Bioinformatics* **1**, 47-50.
- Fielding AH, Bell JF (1997) A review of methods for the assessment of prediction errors in conservation presence/absence models. *Environmental Conservation* **24**, 38-49.
- Galbreath KE, Hafner DJ, Zamudio KR (2009) When cold is better: climate-driven elevation shifts yield complex patterns of diversification and demography in an alpine specialist (American pika, *Ochotona princeps*). *Evolution* **63**, 2848-2863.
- Gannon WL, Sikes RS, Comm ACU (2007) Guidelines of the American Society of Mammalogists for the use of wild mammals in research. *Journal of Mammalogy* **88**, 809-823.
- Gibbs JP, Breisch AR (2001) Climate warming and calling phenology of frogs near Ithaca, New York, 1900-1999. *Conservation Biology* **15**, 1175-1178.

- Graham RW, Lundelius EL, Graham MA, Schroeder EK, Toomey RS, Anderson E, Barnosky AD, Burns JA, Churcher CS, Grayson DK, Guthrie RD, Harington CR, Jefferson GT, Martin LD, McDonald HG, Morlan RE, Semken HA, Webb SD, Werdelin L, Wilson MC (1996) Spatial response of mammals to late quaternary environmental fluctuations. *Science* **272**, 1601-1606.
- Grayson DK (1993) *The desert's past. A natural prehistory of the Great Basin* Washington & London.
- Grayson DK (2000) Mammalian responses to Middle Holocene climatic change in the Great Basin of the western United States. *Journal of Biogeography* **27**, 181-192.
- Gummer DL, Forbes MR, Bender DJ, Barclay RMR (1997) Botfly (Diptera: Oestridae) parasitism of Ord's kangaroo rats (*Dipodomys ordii*) at Suffield National Wildlife Area, Alberta, Canada. *Journal of Parasitology* **83**, 601-604.
- Harrison SP (2000) Palaeoenvironmental data sets and model evaluation in PMIP, 9-25.
- Hasumi H, Emori S (2004) K-1 Coupled Model (MIROC) Description. Center for Climate System Research, University of Tokyo, Tokyo, Japan.
- Hayssen V (1991) *Dipodomys microps*. *Mammalian Species*, 1-9.
- Hewitt G (2000) The genetic legacy of the Quaternary ice ages. *Nature* **405**, 907-913.
- Hewitt GM (1996) Some genetic consequences of ice ages, and their role in divergence and speciation. *Biological Journal of the Linnean Society* **58**, 247-276.
- Hewitt GM (1999) Post-glacial re-colonization of European biota. *Biological Journal of the Linnean Society* **68**, 87-112.
- Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005) Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* **25**, 1965-1978.
- Hilbe JM (2008) Systat 12.2: An overview. *American Statistician* **62**, 177-178.
- Hockett BS (2000) Paleobiogeographic changes at the Pleistocene-Holocene boundary near Pintwater Cave, southern Nevada. *Quaternary Research* **53**, 263-269.
- Huppopp O, Huppopp K (2003) North Atlantic Oscillation and timing of spring migration in birds. *Proceedings of the Royal Society of London Series B-Biological Sciences* **270**, 233-240.

- Jezkova T, Jaeger JR, Marshall ZL, Riddle BR (2009) Pleistocene impacts on the phylogeography of the desert pocket mouse (*Chaetodipus penicillatus*). *Journal of Mammalogy* **90**, 306-320.
- Jones KB (1995) *Phylogeography of the desert horned lizard (Phrynosoma platyrhinos) and the short-horned lizard (Phrynosoma douglassii): patterns of divergence and diversity* Dissertation, University of Nevada Las Vegas.
- Kenagy GJ (1973) Adaptations for leaf eating in Great Basin kangaroo rat, *Dipodomys microps*. *Oecologia* **12**, 383-412.
- Kerdelhue C, Zane L, Simonato M, Salvato P, Rousselet J, Roques A, Battisti A (2009) Quaternary history and contemporary patterns in a currently expanding species. *BMC Evolutionary Biology* **9**, 14.
- Kocher TD, K. TW, A. M, Edwards SV, Pääbo S, Villablanca FX, Wilson AC (1989) Dynamics of Mitochondrial-DNA Evolution in Animals - Amplification and Sequencing with Conserved Primers. *Proceedings of the National Academy of Sciences of the United States of America* **86**, 6196-6200.
- Librado P, Rozas J (2009) DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* **25**, 1451-1452.
- Lobo JM, Jimenez-Valverde A, Real R (2008) AUC: a misleading measure of the performance of predictive distribution models. *Global Ecology and Biogeography* **17**, 145-151.
- Manni F, Guerard E, Heyer E (2004) Geographic patterns of (genetic, morphologic, linguistic) variation: How barriers can be detected by using Monmonier's algorithm. *Human Biology* **76**, 173-190.
- Menzel A (2000) Trends in phenological phases in Europe between 1951 and 1996. *International Journal of Biometeorology* **44**, 76-81.
- Miller MP (2005) Alleles In Space (AIS): Computer software for the joint analysis of interindividual spatial and genetic information. *Journal of Heredity* **96**, 722-724.
- Miller MP, Bellinger MR, Forsman ED, Haig SM (2006) Effects of historical climate change, habitat connectivity, and vicariance on genetic structure and diversity across the range of the red tree vole (*Phenacomys longicaudus*) in the Pacific Northwestern United States. *Molecular Ecology* **15**, 145-159.
- Moore WS (1995) Inferring phylogenies from mtDNA variation - mitochondrial gene trees versus nuclear gene trees. *Evolution* **49**, 718-726.



- Moritz C, Patton JL, Conroy CJ, Parra JL, White GC, Beissinger SR (2008) Impact of a century of climate change on small-mammal communities in Yosemite National Park, USA. *Science* **322**, 261-264.
- Mulcahy DG (2008) Phylogeography and species boundaries of the western North American nightsnake (*Hypsiglena torquata*): Revisiting the subspecies concept. *Molecular Phylogenetics and Evolution* **46**, 1095-1115.
- Olson DM, Dinerstein E, Wikramanayake ED, Burgess ND, Powell GVN, Underwood EC, D'Amico JA, Itoua I, Strand HE, Morrison JC, Loucks CJ, Allnutt TF, Ricketts TH, Kura Y, Lamoreux JF, Wettengel WW, Hedao P, Kassem KR (2001) Terrestrial ecoregions of the worlds: A new map of life on Earth. *Bioscience* **51**, 933-938.
- Otto-Bliesner BL, Brady EC, Clauzet G, Tomas R, Levis S, Kothavala Z (2006) Last Glacial Maximum and Holocene climate in CCSM3. *Journal of Climate* **19**, 2526-2544.
- Parmesan C (2006) Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology Evolution and Systematics* **37**, 637-669.
- Parmesan C, Ryrholm N, Stefanescu C, Hill JK, Thomas CD, Descimon H, Huntley B, Kaila L, Kullberg J, Tammaru T, Tennent WJ, Thomas JA, Warren M (1999) Poleward shifts in geographical ranges of butterfly species associated with regional warming. *Nature* **399**, 579-583.
- Pearson RG, Dawson TP, Berry PM, Harrison PA (2002) SPECIES: A Spatial Evaluation of Climate Impact on the Envelope of Species. *Ecological Modelling* **154**, 289-300.
- Peterson AT, Nyari AS (2008) Ecological niche conservatism and Pleistocene refugia in the thrush-like mourner, *Schiffornis* sp., in the neotropics. *Evolution* **62**, 173-183.
- Peterson AT, Papes M, Soberon J (2008) Rethinking receiver operating characteristic analysis applications in ecological niche modeling. *Ecological Modelling* **213**, 63-72.
- Phillips SJ, Anderson RP, Schapire RE (2006) Maximum entropy modeling of species geographic distributions. *Ecological Modelling* **190**, 231-259.
- Phillips SJ, Dudik M (2008) Modeling of species distributions with Maxent: new extensions and a comprehensive evaluation. *Ecography* **31**, 161-175.
- Phillips SJ, Dudik M, Schapire RE (2004) A maximum entropy approach to species distribution modeling, 655-662.

- Polzin T, Daneshmand SV (2003) On Steiner trees and minimum spanning trees in hypergraphs. *Operations Research Letters* **31**, 12-20.
- Post E, Langvatn R, Forchhammer MC, Stenseth N (1999) Environmental variation shapes sexual dimorphism in red deer. *Proceedings of the National Academy of Sciences of the United States of America* **96**, 4467-4471.
- Pounds JA, Fogden MPL, Campbell JH (1999) Biological response to climate change on a tropical mountain. *Nature* **398**, 611-615.
- Raes N, ter Steege H (2007) A null-model for significance testing of presence-only species distribution models. *Ecography* **30**, 727-736.
- Raines GL, Sawatzky DL, Connors KA (1996) Great Basin geoscience data base. In: *U.S. Geological Survey Digital Data Series DDS-41*.
- Rambaut A, Drummond AJ (2007) Tracer v1.4. <http://beast.bio.ed.ac.uk/Tracer>.
- Ramos-Onsins SE, Rozas J (2002) Statistical properties of new neutrality tests against population growth. *Molecular Biology and Evolution* **19**, 2092-2100.
- Randin CF, Dirnbock T, Dullinger S, Zimmermann NE, Zappa M, Guisan A (2006) Are niche-based species distribution models transferable in space? *Journal of Biogeography* **33**, 1689-1703.
- Reale D, McAdam AG, Boutin S, Berteaux D (2003) Genetic and plastic responses of a northern mammal to climate change. *Proceedings of the Royal Society of London Series B-Biological Sciences* **270**, 591-596.
- Reheis MC (1999) Extent of Pleistocene lakes in the western Great Basin. In: *Geological Survey Miscellaneous Field Studies Map MF-2323*. Geological Survey, Denver, CO.
- Rogers AR, Harpending H (1992) Population growth makes waves in the distribution of pairwise genetic differences. *Molecular Biology and Evolution* **9**, 552-569.
- Rowe R, Finarelli J, Rickart E (2010) Range dynamics of small mammals along an elevational gradient over an 80-year interval. *Global Change Biology* **early view**, 1-14.
- Schmitt T, Rober S, Seitz A (2005) Is the last glaciation the only relevant event for the present genetic population structure of the meadow brown butterfly *Maniola jurtina* (Lepidoptera : Nymphalidae)? *Biological Journal of the Linnean Society* **85**, 419-431.

- Schneider S, Excoffier L (1999) Estimation of past demographic parameters from the distribution of pairwise differences when the mutation rates vary among sites: Application to human mitochondrial DNA. *Genetics* **152**, 1079-1089.
- Slatkin M, Hudson RR (1991) Pairwise comparisons of mitochondrial-DNA sequences in stable and exponentially growing populations. *Genetics* **129**, 555-562.
- Smith FA, Betancourt JL (2003) The effect of Holocene temperature fluctuations on the evolution and ecology of *Neotoma* (woodrats) in Idaho and northwestern Utah. *Quaternary Research* **59**, 160-171.
- Spaulding AW (1990a) Vegetational and climatic development of the Mojave Desert: The last glacial maximum to the Present. In: *Packrat middens: The last 40,000 years of biotic change* (eds. Betancourt JL, Van Devender TR, Martin PS), pp. 200-239. The University of Arizona Press, Tucson.
- Spaulding WG (1990b) Vegetation dynamics during the last deglaciation, southeastern Great Basin, USA. *Quaternary Research* **33**, 188-203.
- Spaulding WG, Leopold EB, Van Devender TR (1983) Late Wisconsin paleoecology of the American southwest. In: *Late Wisconsin of the United States* (ed. Porter SC), pp. 259-293. University of Minnesota Press, Minneapolis.
- Summers HA, Smith BN, Hansen LD (2009) Comparison of respiratory and growth characteristics of two co-occurring shrubs from a cold desert, *Coleogyne ramosissima* (blackbrush) and *Atriplex confertifolia* (shadscale). *Journal of Arid Environments* **73**, 1-6.
- Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: Molecular evolutionary genetics analysis (MEGA) software version 4.0. *Molecular Biology and Evolution* **24**, 1596-1599.
- Thomas WK, Paabo S, Villablanca FX, Wilson AC (1990) Spatial and temporal continuity of kangaroo rat populations shown by sequencing mitochondrial DNA from museum specimens. . *Journal of Molecular Evolution* **31**, 101-112.
- Thompson RS (1990) Late Quaternary vegetation and climate in the Great Basin. In: *Packrat middens; The last 40,000 years of biotic change* (eds. Betancourt JL, Van Devender TR, Martin PS), pp. 200-239. The University of Arizona Press, Tucson.
- Thompson RS, Anderson KH (2000) Biomes of western North America at 18,000, 6000 and 0 C-14 yr BP reconstructed from pollen and packrat midden data. *Journal of Biogeography* **27**, 555-584.

- Thompson RS, Mead JI (1982) Late Quaternary Environments and Biogeography in the Great-Basin. *Quaternary Research* **17**, 39-55.
- Thuiller W, Brotons L, Araujo MB, Lavorel S (2004) Effects of restricting environmental range of data to project current and future species distributions. *Ecography* **27**, 165-172.
- Urban MC, Phillips BL, Skelly DK, Shine R (2007) The cane toad's (*Chaunus* [*Bufo*] *marinus*) increasing ability to invade Australia is revealed by a dynamically updated range model. *Proceedings of the Royal Society B-Biological Sciences* **274**, 1413-1419.
- Vrba ES (1992) Mammals as a key to evolutionary theory. *Journal of Mammalogy* **73**, 1-28.
- Waltari E, Hijmans RJ, Peterson AT, Nyari AS, Perkins SL, Guralnick RP (2007) Locating Pleistocene refugia: comparing phylogeographic and ecological niche model predictions. *PLoS ONE* **2**, e563.
- Warren DL, Glor RE, Turelli M (2008) Environmental niche equivalency versus conservatism: quantitative approaches to niche evolution. *Evolution* **62**, 2868-2883.
- Watson DF, Philip GM (1985) A Refinement of Inverse Distance Weighted Interpolation. *Geo-Processing* **2**, 315-327.
- Wells PV (1983) Paleobiography of montane islands in the Great Basin since the last glaciopluvial. *Ecological Monographs* **53**, 341-382.
- Wiens JJ, Graham CH (2005) Niche conservatism: integrating evolution, ecology, and conservation biology. *Annual Review of Ecology Evolution and Systematics* **36**, 519-539.
- Winker K, Rocque DA, Braile TM, Pruett CL (2007) Vainly beating the air: Species-concept debates need not impede progress in science or conservation. *Ornithological Monographs* **63**, 30-44.
- Zink RM, Barrowclough GF (2008) Mitochondrial DNA under siege in avian phylogeography. *Molecular Ecology* **17**, 2107-2121.

## CHAPTER 3

### CLIMATIC PROMOTION OF NICHE SHIFTS: A CASE STUDY OF THE DESERT HORNED LIZARD (*PHRYNOSOMA PLATYRHINOS*).

#### Abstract

During climate change, species are often assumed to shift geographic distributions by tracking environmental conditions to which they are adapted while maintaining their niches, generally referred to as niche conservatism. To test this assumption, I used mitochondrial DNA and climatic niche assessments to evaluate response to climate changes following the Last Glacial Maximum (LGM) in the desert horned lizard (*Phrynosoma platyrhinos*). The phylogeographic analysis indicated persistence of *P. platyrhinos* through climate changes within the southern Mojave and Sonoran deserts and a recent expansion from these deserts into the Great Basin in western North America, likely following warming climate after the LGM. Assessment of climatic niche similarity revealed that populations within the Great Basin occupy a different climatic niche (colder and wetter) with respect to the southern, persistent populations. Additionally, my assessment inferred that the climatic niche within the southern regions during the LGM did not reach the temperature and precipitation extremes that *P. platyrhinos* experiences within the Great Basin today. I hypothesize that climatic changes at the end of the LGM opened novel niche space within the Great Basin that promoted a niche shift in these lizards.

## Introduction

The burgeoning awareness of global climate change has focused the attention of biologists on species responses to past, current, and potential future climatic changes (Coetzee *et al.* 2009; Galbreath *et al.* 2009; Peterson *et al.* 2004; Thuiller *et al.* 2008; Waltari *et al.* 2007). A general assumption underlying the interplay between current and past models of species response has been that during climate change, the climatic niche of a species (the range of climate conditions utilized by the species) remains relatively stable and thus its geographic range ‘shifts’ by tracking the climate to which it is adapted (Davis & Shaw 2001; Pearman *et al.* 2008; Wiens & Graham 2005). Such niche conservatism *sensu stricto*, also called niche identity (Warren *et al.* 2008), has been a working assumption for climatic niche assessments at time scales across the Last Glacial Maximum, Middle Holocene, the pre-industrial era, and into the foreseeable future. The departure from niche conservatism is a niche shift, a response that has been implicated in episodes of rapid species diversification (i.e., upon entering a new geographic space or adaptive zone), but that has often been considered negligible in species responses to recent climatic changes (Ackerly 2003b).

In general, a niche shift can result from a shift in the fundamental niche (the full spectrum of environmental factors that can be potentially utilized by an organism) or the realized niche (a subset of the fundamental niche actually used by the organism, restricted by historical and biotic factors; Araujo & Guisan 2006; Pearman *et al.* 2008). Niche shifts resulting from an alteration of the realized niche (Jackson & Overpeck 2000; Rodder & Lotters 2009) do not require evolutionary adaptation as species responses

may result from shifts in the realized environment , plasticity of traits (e.g. physiological, behavioral) or through spatial segregation of individuals with certain functional traits into populations following a process of ecological sorting (Ackerly 2003a, b).

Alternatively, the fundamental niche of a species can be shifted or broadened to meet new environmental conditions through evolutionary adaptation (Davis & Shaw 2001; Davis *et al.* 2005; Urban *et al.* 2007). The shift of fundamental niche (evolutionary adaptation) can be difficult to distinguish from realized niche shift (plasticity or spatial segregation) as both can arise from the same underlying processes and may even accompany each other (Csuti 1979).

Niche shifts (either realized or fundamental) have been shown to be triggered by changes in biotic interactions through time or space, such as ecological release from predators or competitors (Holt *et al.* 2005) or by access to novel combinations of environmental variables (Ackerly 2003b; Jackson & Overpeck 2000; Nogues-Bravo 2009; Rodder & Lotters 2009). Invasive species often undergo niche shifts, which may result from differences in biotic and abiotic conditions experienced between native and introduced ranges (Broennimann *et al.* 2007; Rodder & Lotters 2009; Urban *et al.* 2007). Niche shift has also been shown in native populations responding to climatic changes (Davis & Shaw 2001; Davis *et al.* 2005). Novel environmental conditions with no known historical analog are often formed during climate change, creating a spectrum of new habitats to individuals capable of exploiting the novel environment through shifts in their realized or fundamental niches. Environmental changes resulting from glacial-interglacial climatic oscillations throughout the Late Pleistocene are known to have had

pronounced impacts on species ranges and genetic structure (Hewitt 1996, 2000); however, little is known about whether these climatic changes promoted niche shifts, and whether certain areas, time periods, or taxa are more or less prone to such responses in a predictable fashion.

In this study, I evaluate whether climatic changes at the end of the Last Glacial Maximum (LGM; ca. 21,000 years ago; Harrison 2000), which expanded arid habitats within the western deserts of North America, promoted a climatic niche shift in the desert horned lizard (*Phrynosoma platyrhinos*). I follow up on two previous studies of this species in which recent (post-LGM) northward range expansion into the Great Basin was indicated from the more southern Mojave and Sonoran deserts (Jones 1995; Mulcahy *et al.* 2006). I assess the phylogeography of *P. platyrhinos* to evaluate the extent of population persistence within regions throughout the LGM and to detail the recent expansion northwards. I then conduct a climatic niche analysis based on temperature and precipitation variables to explore whether the recently expanding populations occupy a different climatic niche than more southern populations that likely persisted throughout climatic changes. Importantly, *P. platyrhinos* could have expanded its range from areas of the Mojave and Sonoran deserts into the Great Basin without a niche shift, if its current climatic niche within the Great Basin existed in the south during the LGM and then expanded northward at the end of this major climatic event. This would have allowed *P. platyrhinos* to expand its geographic range by tracking the climate to which it was already adapted. To explore this possibility, I project climatic niche models (CNMs) of recently expanding populations to climatic simulations of LGM



conditions (Harrison 2000), and evaluate the prospect that current climatic niches in the Great Basin was mirrored by equivalent conditions during the LGM within more southern latitudes. Finally, I compare climatic niches of persistent and expanding populations of *P. platyrhinos* to the parapatric congeneric species, *P. douglasii* and *P. hernandesi*, that also occupy portions of the Great Basin. I predict that when novel environmental conditions form new potential habitat after a major climatic event, species inhabiting the ecologically closest, and geographically most proximal, niche would be most likely to exploit the new space. In other words, I predict that the current CNMs of the northward expanding populations of *P. platyrhinos* will be more similar to the CNM of the persistent, southern populations of *P. platyrhinos* than to those of *P. douglasii* and *P. hernandesi*. In this study, I generally do not distinguish between shifts in fundamental niche from shifts in realized niche; however, I do discuss and propose hypotheses about both types of a niche shift when appropriate.

## Methods

### Study organism

*Phrynosoma platyrhinos* is one of 17 currently recognized species within the horned lizard genus (Luxbacher & Knouft 2009), all of which occur in western North America. In general, species of *Phrynosoma* tend not to occur in sympatry (Dumas 1964; Leache & McGuire 2006; Luxbacher & Knouft 2009; Montanucci 1981). *Phrynosoma platyrhinos* appears to be part of a species complex with taxa distributed from the northwestern Sonoran Desert through the Mojave and Great Basin deserts (Fig. 3.1). Based on

mitochondrial (mtDNA) sequence data and morphology, Mulcahy *et al.* (2006) found that southeastern populations of *P. platyrhinos* south of the Gila River represented a distinct lineage, which they referred to as *P. goodei*. Mulcahy *et al.* (2006) also identified another population quite divergent from *P. platyrhinos* from the Yuma Proving Grounds (La Paz County, Arizona) along the east side of the Lower Colorado River Valley, but this lineage was not assigned any particular taxonomic status. *Phrynosoma goodei* was shown to hybridize with the flat-tailed horned lizard, *P. mcallii*, and as *P. goodei* appears morphologically intermediate between *P. mcallii* and *P. platyrhinos*, Mulcahy *et al.* (2006) speculated on a potential ancient hybrid origin for *P. goodei*.

#### Taxon sampling

For genetic assessments I acquired samples from 216 individuals of my target taxa from 104 localities: 198 samples from 96 localities of *P. platyrhinos*, five samples from one locality of the divergent lineage at the Yuma Proving Grounds, and 13 samples from seven localities of *P. goodei* (Fig. 3.1A; Appendix 2). I included a sample of *P. mcallii* as an outgroup based on its mtDNA similarity to *P. platyrhinos* (Leache & McGuire 2006; Mulcahy *et al.* 2006). Overall, 30 samples were represented by sequences from GenBank (Mulcahy *et al.* 2006), 19 samples came from museums, and the remaining samples were obtained from specimens specifically captured for this study or accessed from the previous study of Jones (1995). I mostly obtained samples from tail- or toe-clips and then released the animals at the capture site (following a protocol approved by the Animal Care and Use Committee, University of Nevada, Las Vegas).

For niche assessments and niche modeling, I compiled additional occurrence information for *P. platyrhinos* as well as the parapatric species *P. douglasii*, and *P. hernandesi* (Fig. 3.1A, B). Museum records for the occurrence of *P. platyrhinos* were obtained (on 12 February 2009) from HerpNET (<http://herpnet.org>) and the Collections Database of the Museum of Vertebrate Zoology, University of California (<http://mvzarcos.berkeley.edu>). All records missing geographic coordinates and records with geographic uncertainty greater than 5 km were excluded. Occurrence records for *P. hernandesi* and *P. douglasii* followed those reported by Jezkova *et al.* (2009a). Because *P. hernandesi* is widespread and occupies a range of different habitats, the dataset was restricted to samples from a monophyletic mtDNA clade occurring in the Great Basin and Colorado Plateau as identified by Zamudio *et al.* (1997).

#### Laboratory methods

I isolated total genomic DNA from tissues following a phenol-chloroform protocol or using a DNeasy Extraction Kit (Qiagen Inc.). I amplified and sequenced the mitochondrial NADH dehydrogenase subunit 4 (ND4) gene and adjacent tRNA for all samples using primers ND4 and Leu (Arevalo *et al.* 1994). For the phylogenetic analyses, I also amplified and sequenced a portion of the Cytochrome B (cytB) for a subset of samples using primers MVZ 49 and MVZ 14 (Roe *et al.* 1985). I updated amplification protocols through time, but most were accomplished at a 55°C annealing temperature using Takara Ex Taq Polymerase Premix (Takara Mirus Bio, Inc.) followed by purification using ExoSap-IT (USB Corp.). I conducted double-stranded cycle sequencing using fluorescence-based chemistry (BigDye Terminator v. 3.1 Cycle Sequencing Kit) with

electrophoresis and visualization on an ABI Prism 3130 automated sequencer (Applied Biosystems, Inc.). I aligned sequences using SEQUENCHER (v. 4.6; Gene Codes Corp.) and verified the alignment visually.

### Phylogenetic analyses

I conducted phylogenetic analyses using Maximum likelihood (ML) and Bayesian inference (BI) on a combined dataset of 1089 base pairs (bp) of *cytB*, 684 bp of *ND4*, and 128 (or 129) bp of adjacent tRNA for a subset of samples including 21 *P. platyrhinos*, 3 specimens from the Yuma Proving Grounds, 2 *P. goodei*, and a sample of *P. mcallii*. These samples were chosen to represent the major nested clades of *P. platyrhinos* identified from the haplotype network (see below). Prior to BI and ML analyses, I partitioned the dataset variously by gene and codon position (including unpartitioned) and identified best-fitting models for each partition using MRMODELTEST (v. 2.2(Nylander 2004)under the Akaike Information Criterion (Posada & Buckley 2004). To evaluate partitions, I ran each partition in MRBAYES (v. 3.1.2; Ronquist & Huelsenbeck 2003) for 5 million generations and compared marginal likelihood scores using Bayes factors (Kass & Raftery 1995; Table 3.1). For final analyses of BI and ML, I used the HKY+I+ $\Gamma$  model for the 1<sup>st</sup>+2<sup>nd</sup> codon positions of *ND4* and *cytB*, GTR+  $\Gamma$  model for the 3<sup>rd</sup> codon position of *ND4* and *cytB*, and the HKY+I model for the tRNA regions. I assessed the tree topology and clade support in MRBAYES from consensus trees and posterior probabilities of 3 final runs of 10 million generations each, sampled every 100 generations, with the first 2.5 million generations (25,000 trees) discarded as burn-in after assessing stationarity by plotting log-likelihood scores against generations (Leache & Reeder 2002). I conducted

ML analyses in TREEFINDER (Jobb *et al.* 2004) using the “Bootstrap Analysis” option with 1,000 replicates and consensus level 50 to assess nodal support.

I constructed a median-joining network (a maximum parsimony approach) on the concatenated data (812 bp) of partial ND4 and adjacent partial tRNA using the program NETWORK (v. 4.5.1.0; Bandelt *et al.* 1999) for 216 sequences of *P. platyrhinos* including samples from the Yuma Proving Grounds and *P. goodei*. To construct the final network, I weighted transversions twice as high as transitions. After generating the network, I employed the MP option (Polzin & Daneshmand 2003) to remove excessive links from the network. For visualization purposes, I nested the haplotype network (Templeton *et al.* 1987; Templeton & Sing 1993). I also calculated haplotype and nucleotide diversity for the clades of *P. platyrhinos* using ARLEQUIN (v. 3.11; Excoffier *et al.* 2005).

#### Historical demography

I defined the ‘core clade’ of *P. platyrhinos* to exclude the more distantly related samples from the Yuma Proving Grounds and *P. goodei* (see Phylogenetic analyses results), and then analyzed past demographic changes. I assessed the concatenated ND4 and tRNA data using the Bayesian skyline plot (BSP) coalescent model (Drummond *et al.* 2005) implemented in the program BEAST (v. 1.5.3; Drummond & Rambaut 2007). I employed the Bayes factor test (Newton *et al.* 1994; Suchard *et al.* 2001) implemented in TRACER (v. 1.5; Rambaut & Drummond 2007) to assess partitioning and chose to partition by genes and codon positions. Following assessment in MRMODELTEST, I selected substitution models GTR+I+ $\Gamma$  for the 1<sup>st</sup>+2<sup>nd</sup> codon positions of ND4, GTR+ $\Gamma$  for the 3<sup>rd</sup> codon position of ND4, and HKY+I for the tRNA. I used a strict molecular clock

after assessing for clock-like behavior (Drummond *et al.* 2007), a general mtDNA substitution rate of 1% /lineage/million years available from the literature (Macey *et al.* 1999), 5 skyline groups, and a 2-year generation time (Pianka & Parker 1975). I conducted several independent Markov Chain Monte Carlo (MCMC) runs of 80 million generations, each with sampling every 8000 generations and a burn-in of 10%. For final analysis, four MCMC runs (all with similar results) were combined using LOGCOMBINER (distributed with BEAST). I checked convergence (effective sample sizes > 200) and visualized the median and 95% highest posterior density intervals using TRACER.

#### Niche similarity comparisons between persistent and expanding populations

I conducted climatic niche comparisons following the methodology of Rissler and Apodaca (2007) and Kozak and Wiens (2006) among geographically persistent versus recently expanded populations of *P. platyrhinos* in order to evaluate whether the latter exhibit a significantly different climatic niche. I assigned each sampling locality to persistent or expanding groups of populations based on genetic patterns (see Results for patterns). I also used museum records, assigning specimens to a particular persistent or recently expanded group if the record was geographically located within the minimum convex polygon of all genetically sampled localities that belonged to that group. Museum records that fell outside all polygons were discarded from the analysis. Minimum convex polygons for groups partly overlapped (Fig. 3.1B), thus not all localities were independent in the analyses. From each occurrence record I extracted values from 19 bioclimatic variables (Kozak & Wiens 2006; Waltari *et al.* 2007) derived from the WorldClim dataset (v. 1.4) with resolution of 2.5 minutes (Hijmans *et al.* 2005).

In SYSTAT (v.12; Hilbe 2008), I conducted a Principal Component Analysis (PCA) to reduce the 19 bioclimatic variables, some of which are correlated (Kozak & Wiens 2006), to principal components (PCs). Factor scores of PCs with eigenvalues >1 were saved. After confirming that PC scores were normally distributed, I ran unbalanced ANOVA based on Type III sums of squares with PCA axis scores as dependent variables and the groups as fixed factors to test for the separation of population groups into different climatic niches. I used Post hoc Tukey tests to compare the least square means of each PC.

#### Evaluation of a niche shift in *P. platyrhinos*

In order to evaluate whether range shifts in *P. platyrhinos* were accompanied by niche shifts, I used climatic niche modeling (more broadly referred to as ecological niche modeling) to assess whether the climatic niche currently occupied by the expanding populations of *P. platyrhinos* had an equivalent during the LGM within the more persistent southern areas. This analysis was conducted under the assumption that the modeling approach was capturing climatic niches at a scale of resolution sufficiently relevant to species biology and ecology (see Discussion). I used the software package MAXENT (v. 3.2.1; Phillips *et al.* 2006) to reconstruct current climatic niches and project those onto the climatic reconstructions of the LGM. MAXENT implements a maximum entropy algorithm to model the niche of a species from data on species occurrence (presence-only data) and environmental variables. For occurrence records, I used the same persistent and expanding groups as in the climatic niche similarity assessment. For environmental layers of current climatic conditions, I started with the same 19

bioclimatic variables used in the previous assessment that were clipped to the extent of western North America. Because some of the 19 bioclimatic variables are correlated (Kozak & Wiens 2006), I evaluated pairwise correlations on values extracted from the presence records, and considered any two variables highly correlated when the correlation coefficient was  $\geq 0.9$ . Subsequently, I selected the following 13 variables for building the CNMs: Bio2 - mean diurnal range, Bio3 - isothermality, Bio4 - temperature seasonality, Bio5 - maximum temperature of warmest month, Bio6 - minimum temperature of coldest month, Bio8 - mean temperature of wettest quarter, Bio9 - mean temperature of driest quarter, Bio10 - mean temperature of warmest quarter, Bio13 - precipitation of wettest month, Bio15 - precipitation seasonality, Bio17 - precipitation of driest quarter, Bio18 - precipitation of warmest quarter, Bio19 - precipitation of coldest quarter. I also explored selection of variables based on perceived biological relevance to *P. platyrhinos* and model importance, but overall model predictions were similar in all cases.

For environmental layers representing the climatic conditions of the LGM, I used the current set of coupled ocean-atmosphere simulations (Harrison 2000) available through the Paleoclimatic Modelling Intercomparison Project (PMIP; Braconnot *et al.* 2007). PMIP has established a protocol followed by participating modeling groups for LGM simulations regarding concentration of greenhouse gases, ice sheet coverage, insolation, or change in topography caused by lowering sea levels. Of more than 15 available climatic models, I used the following two: Community Climate System Model (CCSM v. 3; Otto-Bliesner *et al.* 2006) with a resolution of  $1^\circ$ , and the Model for



Interdisciplinary Research on Climate (MIROC v. 3.2; Hasumi & Emori 2004) with an original spatial resolution of  $1.4^{\circ} \times 0.5^{\circ}$  (Braconnot *et al.* 2007). The original climatic variables used in these models have been downscaled to the spatial resolution of 2.5 minutes under the assumption of high spatial autocorrelation and converted to bioclimatic variables (Hijmans *et al.* 2005; Peterson & Nyari 2008).

I used the default parameters of MAXENT (i.e. 500 maximum iterations, convergence threshold of 0.00001, regularization multiplier of 1, 10,000 background points) with the following user-selected features: application of random seed, duplicate presence records removal, logistic probabilities for the output (Phillips & Dudik 2008). Since I were interested in the extent of analogous climates during two time periods, I used the 'do not interpolate' option in MAXENT. This option insures that the probability values for the LGM variables are not going to be extrapolated beyond the range of values used for model calibration. Therefore, my LGM models should not be interpreted as areas of persistence for *P. platyrhinos*, but only as the extent of specific climatic combinations during two time periods. I ran 10 replicates in MAXENT for each group (projected 10 times on each of the CCSM and MIROC climatic reconstructions), but present an average model. Because the analyses are greatly dependent on the chosen threshold cut-off values used for interpreting the CNMs, I evaluated across two extreme thresholds corresponding to 0 and 10% omission error.

#### Niche overlap between *P. platyrhinos* and parapatric species

I used ENM tools (Warren *et al.* 2008) to assess the niche overlap between persistent and expanding populations of *P. platyrhinos* with respect to *P. douglasii*, and

parapatric populations of *P. hernandesi*. The CNMs for all species were generated in MAXENT as described above. I measured niche overlap among CNMs of taxa pairs using the coefficient of niche similarity  $D$ , a metric for quantifying similarity of CNMs (Warren *et al.* 2008). Values of  $D$  range from zero to one, where zero represents no overlap between the CNMs of two taxa and one represents total (absolute) overlap. I then conducted randomization tests of 50 pseudoreplicates where the records of each pair are randomly partitioned and overlap is calculated using  $D$ . From the 50 pseudoreplicates, I built a null distribution of  $D$  that was compared to the observed value of  $D$ . A one-tailed test was used to evaluate the hypothesis that observed  $D$  is drawn from a statistical population that has a mean that is significantly different than that of the random  $D$ .

## Results

### Phylogenetic and network analyses

Assessments using ML and BI methods based on a subset of haplotypes produced identical tree topologies with similar nodal support based on posterior probabilities and bootstrap values (Fig. 3.2). The inferred topology shows a strongly supported core clade of *P. platyrhinos* with *P. goodei* as the sister clade, and samples from the Yuma Proving Grounds appearing as a clade outside this group. The basal position of the Yuma Proving Grounds clade, however, was not strongly supported, and the relationship could be interpreted as a polytomy of three strongly supported clades. These results are consistent with patterns identified by Mulcahy *et al.* (2006) for the same taxa. Within

the core clade of *P. platyrhinos*, I identified three well-supported subclades which I refer to as the Western, Northeastern, and Southeastern clades (Fig. 3.2, 3.3).

I identified 120 unique haplotypes from 216 sequences (812 bp of ND4+tRNA) representing *P. platyrhinos*, *P. goodei*, and samples from the Yuma Proving Grounds. Most haplotypes were represented by 1–4 samples with the exceptions of the abundant and widespread haplotypes Hw1, Hw2, He1, and He2 which give rise to numerous satellite (newly evolved) haplotypes (Fig. 3.3). Using nesting rules, I identified 15 four-step clades (Fig. 3.4). Clade 1 corresponds to the 5 animals collected from the Yuma Proving Grounds, and clades 2 and 3 correspond to individuals of *P. goodei*. Clade 4 represents all other samples from Arizona east of Colorado River with the exception of the sample from locality 12 (Fig. 3.1A, 3.4). Clades 5, 7–10, 12–14 are mainly distributed within the Mojave and northwestern Sonoran deserts, west of the Colorado River, in California, Nevada, Utah and northern Baja California (Fig. 3.4). Clade 15 was represented by one individual from the House Rock Valley in Arizona. The clades in the Mojave and northwestern Sonoran deserts overlap geographically and each clade exhibits numerous diverse and geographically restricted haplotypes. Within the southern deserts, haplotype and nucleotide diversities were relatively high (0.992 and 0.022, respectively) indicative of long-term persistence within the area (Hewitt 1996, 2000).

Clade 6 can be found within the northern Mojave Desert and within the western Great Basin. The haplotype Hw2 in clade 6 and its satellite haplotypes comprise all individuals from the southwestern Great Basin, and Hw1 and its satellite haplotypes

comprise all individuals in the northwestern Great Basin (Fig. 3.4). Similarly, clade 11 extends from the eastern Mojave Desert into the eastern Great Basin with widespread haplotypes He1 and He2, and their satellite haplotypes, comprising those in the northeastern and southeastern Great Basin respectively (Fig. 3.4). This structure is most often indicative of a recent range expansion, when a subset of haplotypes (in this case Hw1, Hw2, He1, He2) expands across large areas from the source haplotype group followed by emergence of new mutations (see Discussion). Accordingly, these areas dominated by one of the four abundant and widespread haplotypes were likely colonized by *P. platyrhinos* only recently. Across the Great Basin, haplotype and nucleotide diversities were lower (0.877 and 0.016, respectively), despite the presence of the two quite divergent clades (6 and 11) within this overall region.

#### Historical demography

To estimate the starting time for growth of genetic diversity relative to the last glacial period (Late Pleistocene) I used the BSP analysis and a general mutation rate of 1% /lineage/million years. The plot (Fig. 3.5) indicates a rapid and recent increase in genetic diversity, which is consistent with recent population expansion. This increase in genetic diversity appears to have begun accumulating sometime during the latest glacial period, but I cannot rule out a post-glacial expansion.

#### Niche similarity comparisons between persistent and expanding populations

I assessed climatic niche similarity between areas occupied by persistent populations and the areas that exhibit signals of recent range expansion. I analyzed the eastern and western expanding populations separately as each represents a different genetic clade

(Fig. 3.4). In the expanding western (Ew) group, I included haplotypes Hw1, Hw2 and their satellite haplotypes from clade 6, and in the expanding eastern (Ee) group, I included haplotypes He1, He2 and their satellite haplotypes from clade 11 (Fig. 3.1B). Although satellite haplotypes can be identified visually as they cluster around ancestral nodes, I also used a star contraction method in NETWORK to confirm such clusters (Forster *et al.* 2001). Other haplotypes in clades 6 and 11 from more southern areas were excluded from the analysis, as these did not exhibit a clear signal of range expansion. The remaining clades from the southern deserts that exhibit high genetic diversity were assigned to the Persistent group (excluding clades 1-3, as these represent more distantly related *P. goodei* and Yuma Proving Grounds animals). The final occurrence dataset included 165 records in the Persistent group, 65 occurrence records in the Ew group, and 25 occurrence records in the Ee group.

PCA reduced the 19 bioclimatic variables to four PCs with eigenvalues  $\geq 1$  that explained 50.9, 25.0, 9.6, and 5.8 percent of the total variance, respectively. The scatterplot for the first two PCs shows separation between the persistent and the expanding groups with the latter experiencing (on average) lower temperatures, greater temperature seasonality, greater annual range in temperature, higher precipitation, and lesser seasonality in precipitation (Fig. 3.6A). The ANOVA showed significant differences in least square means for all four PCs (for all tests,  $P < 0.05$ ; Fig. 3.6B), and the pairwise Tukey tests (Table 3.2) revealed that the PCs could be used to easily differentiate among the groups.

### Evaluation of a niche shift in *P. platyrhinos*

The CNM for the Persistent group in general did not over-predict the current climatic niche into areas now occupied by the expanding populations, although some areas within the southwestern Great Basin were predicted as suitable using the more liberal (0% omission) threshold (Fig. 3.7A). The model based on a zero omission threshold also over-predicted into areas currently occupied by congeneric species (i.e. *P. goodei*, *P. mcallii*, and *P. blainvillii*; Fig. 3.7B). For the expanding groups within the Great Basin (Ew and Ee), the thresholds used resulted in little difference in overall outcomes of the models. The CNM for the Ew group captures the distribution of the group within the western part of the Great Basin (Fig. 3.7B). The CNM based on the Ee group correctly captures the eastern Great Basin, but over-predicts across the western Great Basin, as well as to the east into the Colorado Plateau where *P. platyrhinos* is not known to occur (Tanner 1999).

The LGM projections for the Persistent group (Fig. 3.7D,G) show retention of the climatic niche within areas of current range, although the overall extent is substantially restricted, especially in the MIROC model (Fig. 3.7G). The LGM projections of climatic niches for the Ew and Ee groups were greatly reduced in size (Fig. 3.7E,F,H,I). The LGM projections for the Ew group shows retention of the current climatic niche only within small patches in the western Great Basin (CCSM, Fig. 3.7E) and Columbia Plateau (MIROC, Fig. 3.7H). Only the CCSM model projects the Ew climatic niche to a small area in the Sonoran Desert currently occupied by the Persistent group, but within the area covered by the Pleistocene Lake Cahuilla (Fig. 3.7E). The LGM projections for the Ee

group shows displacement of the current climatic niche to the western Great Basin (CCSM, Fig. 3.7F) and some additional areas within the northeastern Great Basin (MIROC with the less conservative threshold only), although this area was covered predominately by the pluvial Lake Bonneville. The Ee climatic niche does not project to the areas currently inhabited by the Persistent group.

#### Niche overlap between *P. platyrhinos* and parapatric species

Using the measure D, I could reject the hypothesis of niche identity ( $P < 0.001$ ) for all pairs of taxa. In general, the climatic niche overlaps (Table 3.3) between the persistent and expanding groups of *P. platyrhinos* were relatively low ( $D = 0.11$  and  $0.24$ ), indicating dissimilarity in climatic regimes. Niche overlap between the two expanding groups of *P. platyrhinos* was relatively high ( $D = 0.44$ ), despite the fact that there is no geographic overlap between the ranges. The overlaps of the expanding groups with parapatric *P. hernandesi* were also high ( $D = 0.53$  and  $0.32$ ), while overlaps with *P. douglasii* were slightly lower ( $D = 0.27$  and  $0.32$ ).

### Discussion

#### Phylogeographic analyses

Within areas of the Mojave and Sonoran deserts, *P. platyrhinos* exhibits pronounced genetic structuring and high genetic diversity (Fig. 3.4) consistent with population persistence. The levels of genetic diversity and structure in *P. platyrhinos* appear higher in comparison to genetic patterns documented in several other warm-desert taxa that were interpreted as an evidence of geographically restricted glacial refugia in northern

regions during the Late Pleistocene (Castoe *et al.* 2007; Douglas *et al.* 2006; Jezkova *et al.* 2009b; Murphy *et al.* 2006). The genetic patterns of *P. platyrhinos* indicate that the species probably remained quite widespread throughout much of the northern Sonoran Desert and maintained populations within portions of the Mojave Desert despite the climatic changes of the Late Pleistocene.

Populations of *P. platyrhinos* within the Great Basin exhibit low genetic diversity and little genetic structure indicative of recent range expansion from areas of persistence in the south. Within the Great Basin, the haplotype network shows a few central, abundant, and widespread haplotypes surrounded by numerous, low frequency haplotypes separated by one or a few mutational steps (Fig. 3.3), with the satellite haplotypes being geographically restricted. This pattern is consistent with leading edge colonization where the range expansion involves random subsets of individuals from the populations at the colonization front (Excoffier *et al.* 2009; Hampe & Petit 2005; Hewitt 1993; Hewitt 2000). The genetic evidence suggests that *P. platyrhinos* expanded along two different low elevation colonization routes into the Great Basin, one along an eastern corridor into the Bonneville Basin and the other along a western corridor into the Lahontan Basin. These expansions involved individuals from two different clades, with clade 6 expanded along the western front and clade 11 along the eastern front. As envisioned for leading edge colonization, genetic diversity typically decreases in the direction of expansion due to loss of haplotypes through founder effects (thinning of haplotypes; Hewitt 1996). The northward expansion in *P. platyrhinos* clearly exhibits this pattern with only haplotypes from clades 6 and 11 evident in samples from above 38° of



latitude and only single ancestral haplotypes, Hw1 and He1, found above 39° of latitude along each of the two expansion routes (Fig. 3.4). Under this expansion scenario, populations grow exponentially and private mutations typically become fixed with higher frequency in the newly invaded areas. Such newly evolved satellite haplotypes are evident around each of the four ancestral haplotypes associated with the range expansions (Fig. 3.3). Recent population expansion in *P. platyrhinos* is also consistent with the pattern exhibited in the coalescent assessment of the sequence data presented in the BSP (Fig. 3.5).

The expansion of *P. platyrhinos* into the Great Basin likely followed warming climate and desiccation of pluvial lakes at the end of the LGM. Such northward expansion of warm-desert organisms is not unique to *P. platyrhinos* as similar patterns have been documented or proposed in several plant and animal taxa (Hockett 2000; Mulcahy 2008; Pavlik 1989). My coalescent assessment of the sequence data under the presumed mutation rate estimates the timing of the expansion to the latest glacial period (Fig. 3.5), but prior to the LGM time frame. Only a slightly faster mutation rate, however, would need to be assumed for the assessment to be consistent with post-Pleistocene expansion (Galbreath *et al.* 2009).

#### Niche similarity comparisons between persistent and expanding populations

Climatic niche comparisons revealed that the recently expanding populations of *P. platyrhinos* within the western and eastern Great Basin occupy a different climatic niche than that of the persistent populations in the Mojave and Sonoran deserts. Individuals from the expanding populations within the Great Basin experience on average higher

precipitation and lower temperatures than individuals from the southern populations (Fig. 3.6; 3.8). *Phrynosoma platyrhinos* within the eastern Great Basin experience the most extreme conditions with the lowest temperatures, largest annual temperature range, and highest precipitation (Fig. 3.8). These climatic differences likely impact latitudinal shifts in ecological, behavioral, and physiological traits. For example, northern animals may compensate for a colder climate behaviorally (Kearney *et al.* 2009) by shifting activity periods and occupying microhabitats with suitable thermal conditions (e.g. basalt formations), or physiologically by lowering active body temperatures (Monasterio *et al.* 2009). Changes in mean body temperature may in turn affect life history traits such as growth rate, body size, and reproductive rate (Sears & Angilletta 2004). Indeed, there is evidence that populations of *P. platyrhinos* in the Great Basin exhibit increased sexual size dimorphism, shorter breeding season, smaller number of clutches per season, and possibly larger clutch size than populations within the southern deserts (Pianka & Parker 1975). Such latitudinal shifts in natural history traits are not unique and have been documented in many reptilian taxa (Fitch 1985; Iverson *et al.* 1993).

Whether *P. platyrhinos* has reached its tolerance limits for climatic variables is not entirely clear. Interestingly, I found *P. platyrhinos* near Elko, NV (Fig. 3.1, locality 58), although only *P. hernandesi* has been reported from this higher elevation area in the past (Jezkova *et al.* 2009a). I speculate that *P. platyrhinos* has invaded this area only within the last few decades, either following a further shift in tolerance limits for climatic variables or in response to a subtle but favorable shift in regional climate. The

haplotype (He1) identified in the two samples from Elko belong to the eastern clade, even though the area is geographically closer to known occurrence records for *P. platyrhinos* from lower elevations to the west that are likely occupied by individuals from the western clade (Fig. 3.1, locality 68 and occurrence records between localities 58 and 68) and that are connected to the Elko site by a potential dispersal corridor along the Humboldt River. I speculate that the individuals from the eastern clade, which on average inhabit the most regionally extreme climatic conditions for the species (see above), might have been better suited to invade the cold Elko area.

#### Evaluation of a niche shift in *P. platyrhinos*

The LGM models presented herein could possibly be interpreted as indicating areas of potential persistence for *P. platyrhinos* within the Great Basin assuming that the climatic niche between the two time periods was conserved. My genetic data, however, indicate that *P. platyrhinos* did not persist within the northern regions during the LGM. I use these models unconventionally to evaluate whether climatic conditions currently occupied by the expanding populations of *P. platyrhinos* in the Great Basin had equivalents during the LGM within the southern areas currently occupied by persistent populations. If such patterns were found, then *P. platyrhinos* could have expanded its range from the Mojave and Sonoran deserts into the Great Basin without a niche shift. My projections, however, indicate that the climatic niches within the Great Basin currently occupied by *P. platyrhinos* do not appear to have been predominately shifted southwards during the LGM (Fig. 3.7). These results are in agreement with paleo-environmental reconstructions, based on pollen and macrofossil data from packrat

middens, indicating that temperature and precipitation averages within large areas of the southern deserts never reached the climate extremes that *P. platyrhinos* currently experiences within the Great Basin. The paleo-environmental reconstructions suggest that the temperature in the Mojave Desert was on average 6°C colder and precipitation 40 % higher during the LGM than today (Spaulding 1990). In comparison, my niche assessment indicates that the current average annual temperature experienced by populations within the Great Basin is roughly 9°C lower than that of populations in the Mojave and Sonoran deserts, and annual precipitation within the areas occupied in the eastern Great Basin (Ee group) is almost 70 % higher than that for the areas occupied by the persistent populations (2.8). These patterns further support my interpretations from the CNMs and the hypothesis that *P. platyrhinos* did not just shift range in response to warming climate but that some *P. platyrhinos* experienced a niche shift in certain climatic variables that allowed the northward range expansions.

An important question to ask here is whether the postulated niche shift experienced by populations invading the Great Basin is biologically meaningful? As mentioned above, variability in ecological traits has been recorded between the northern and southern populations of *P. platyrhinos* (Pianka & Parker 1975) indicating that the species could have adapted to novel conditions through genetic differentiation and thereby expand its fundamental niche (Csuti 1979; Davis & Shaw 2001; Urban *et al.* 2007). Rapid climatic shifts at the end of the LGM are similar to conditions that species encounter during invasions to new areas. Novel niches with less competition promote rapid dispersal and expansion of populations, possibly followed by directional selection favoring those

individuals that are better adapted to the new environment (Ackerly 2003b; Broennimann *et al.* 2007; Davis & Shaw 2001; Pearman *et al.* 2008). Climatic changes at the end of LGM indeed appear to have produced new combinations of climatic variables within areas of the Great Basin, which had no substantial analog in southern regions during the previous glacial period. Therefore, these developing novel habitats may have promoted adaptation (Ackerly 2003b; Davis & Shaw 2001).

Alternatively, *P. platyrhinos* may exhibit phenotypic plasticity in various life history traits and could have simply expanded its realized niche to take advantage of changing environmental conditions that may well be within its existing range of tolerances (Jackson & Overpeck 2000; Rodder & Lotters 2009). Such a shift could also have been intertwined with a shift in realized environment (i.e. environmental conditions available at any given time; Ackerly 2003a,b; Hoffmann 2005; Jackson & Overpeck 2000). These shifts would not have necessarily required any significant adaptive changes in the biology of the species; common garden and reciprocal transplant experiments could be used to test between these alternatives. In any case, the niche shift in *P. platyrhinos* is interesting as several currently sympatric lizards within the Mojave and Sonoran deserts have not expanded into the Great Basin (e.g. *Dipsosaurus dorsalis*, *Sauromalus obesus*, *Heloderma suspectum*), and while there may be several factors related to these more limited distributions, possibly some species simply were not capable of the scale of niche shift documented in *P. platyrhinos*.

In this study, I focused on the niche shift of *P. platyrhinos* within the Great Basin; however, the niche of persistent populations within the Mojave and Sonoran deserts

has also likely shifted. In particular, following the end of the last glacial period, *P. platyrhinos* did not disappear from the low elevation valleys that now represent the hottest and driest parts of western North America. These areas are currently dominated by novel plant communities, creosote bush-white bursage desertscrub, that did not develop until Holocene times (Hunter *et al.* 2001; Spaulding 1990). This combination of extreme desert climate and novel plant communities apparently had no equivalent during the LGM (Spaulding 1990), although these areas now represent prime habitat occupied by *P. platyrhinos*, as well as many other warm-desert species.

The niche shift assessments conducted here critically depend on accurate climatic reconstructions of the LGM. The differences in the two LGM simulation models (CCSM and MIROC) generate quite different projections, indicating that current understanding of LGM climate as applied broadly is limited. The MIROC model predicts higher precipitation than the CCSM model (which appears to generate a more restricted climatic niche for the Persistent group within the Mojave Desert) but warmer temperatures (which appear to generate more extensive climatic niches for the expanding groups within the Great Basin). In general, interpolations and downscaling of the originally coarse climatic datasets may have produced severe errors or inaccuracies in the base data layers, especially in topographically diverse terrains such as the Great Basin. Further, neither the CCSM nor MIROC models take into account impacts of the large pluvial lakes on local or regional temperature and precipitation. Despite these potential inaccuracies, my interpretations of the overall results are consistent regardless of the model used, which indicates some robustness in my general findings.

Unfortunately, my approach only searches for the presence of climatic variable combinations available during calibration and does not reveal which variables may have actually shifted, to what extent, and the biological relevance of the shifts on *P. platyrhinos*.

#### Niche overlap between *P. platyrhinos* and parapatric species

When a novel niche space opens up after a major climatic event, I might expect the species inhabiting the ecologically closest and geographically most proximal niche to be most likely to colonize the new environment. I predicted that the climatic niche of the recently expanding populations of *P. platyrhinos* would be more similar to the persistent southern populations than to *P. douglasii* and *P. hernandesi* from the Great Basin. Conversely, my results indicate that the climatic niches currently occupied by the expanding populations are more similar to that of *P. hernandesi* and *P. douglasii* than to that of the persistent populations of *P. platyrhinos* (Table 3.3). In other words, given my prediction, *P. hernandesi* and *P. douglasii* should have been more likely than *P. platyrhinos* to populate parts of the Great Basin currently occupied by *P. platyrhinos*.

I do not have a direct explanation for this seeming contradiction, but offer several (not mutually exclusive) hypotheses. First, it is possible that *P. platyrhinos* directly outcompetes these other lizards. Competitive exclusion between *P. platyrhinos* and *P. hernandesi*, and *P. douglasii* has not been tested, but some studies suggest that competition between *Phrynosoma* species might exist despite different reproductive strategies or diet preferences (Montanucci 1981; Pianka & Parker 1975). Second, genetic constraints on adaptation (Davis *et al.* 2005) could possibly prevent *P.*

*hernandesi* and *P. douglasii* from adapting to warming climate or to some other (non-measured) factors that may favor *P. platyrhinos*. Lastly, ranges of many species expand at high latitudes and elevations, but contract at the warm margins in response to warming climate (Hewitt 2000; Merrill *et al.* 2008). The northward expansion of *P. platyrhinos* at the end of the LGM placed the northern populations on the leading edge of novel habitat where adaptation may be enhanced by the increased variability facilitated by gene flow from the centers of the range (Davis & Shaw 2001). Conversely, *P. hernandesi* and *P. douglasii* within the Great Basin were on the trailing edge during warming climate, where adaptation to the changing environment depends predominantly on variation within local populations (Davis & Shaw 2001). As has been hypothesized, changing environment could have produced deteriorating conditions for the trailing-edge populations, leading to population contraction or even extirpation (Davis & Shaw 2001; Hampe & Petit 2005). This leading versus trailing edge effect could have been most pronounced during the warm Middle Holocene when climate would have favored expansion of warm-desert species moving northward along the latitudinal gradients or upward along elevational gradients (Grayson 2000).

### Conclusions

My analyses indicate that *P. platyrhinos* likely persisted within areas of the Mojave and Sonoran deserts through climatic oscillations of the Late Pleistocene. Conversely, this species only recently expanded into the Great Basin, likely following the warming climate along two low elevation corridors at the end of the LGM. The expanding populations did not just track expansion of suitable habitat, but appear to have



experienced niche shift that allowed populations of *P. platyrhinos* to exploit novel (colder and wetter) environmental conditions. The biological mechanisms behind the niche shift, however, remain unclear.

**Table 3.1.** Bayes factors for partitionings of the ND4+cytB+tRNA sequences of *Phrynosoma* taxa calculated in MrBAYES.

| Partition  | N of partitions | Model  | Mean Marginal Likelihood Score | Bayes Factor |
|--|-----------------|--|--------------------------------|--------------|
| All together   | 1               | GTR+I+ $\Gamma$                                      | 5919.62                        |              |
| By gene  | 3               | GTR+I+ $\Gamma$ (cytB)                               | 5914.47                        | 10.3         |
|  |                 | HKY+I+ $\Gamma$ (ND4)                                |                                |              |
|  |                 | HKY+I (tRNA)   |                                |              |
| By 1 <sup>st</sup> + 2 <sup>nd</sup> , 3 <sup>rd</sup> codon             | 3               | HKY+I+ $\Gamma$ (1 <sup>st</sup> + 2 <sup>nd</sup> ) | 5709.16                        | 410.62       |
| (tRNA unpartitioned)   |                 | GTR+ $\Gamma$ (3 <sup>rd</sup> )                     |                                |              |
|  |                 | HKY+I (tRNA)   |                                |              |
| By gene and by 1 <sup>st</sup> + 2 <sup>nd</sup> , 3 <sup>rd</sup> codon | 5               | HKY+I (1 <sup>st</sup> + 2 <sup>nd</sup> cytB)       | 5760.04                        | -101.76      |
| (tRNA unpartitioned)   |                 | GTR+ $\Gamma$ (3 <sup>rd</sup> cytB)                 |                                |              |
|  |                 | HKY+I (1 <sup>st</sup> + 2 <sup>nd</sup> ND4)        |                                |              |
|  |                 | GTR+ $\Gamma$ (3 <sup>rd</sup> ND4)                  |                                |              |
|  |                 | HKY+I (tRNA)   |                                |              |

**Table 3.2.** Significant differentiation of the climatic niche represented by four principal components based on Tukey tests between the Persistent, expanding eastern (Ee) and expanding western (Ew) groups of *Phrynosoma platyrhinos*. Significant values are indicated in bold.

|                        | Persistent   |              |              |              | East expanding (Ee) |       |              |              |
|------------------------|--------------|--------------|--------------|--------------|---------------------|-------|--------------|--------------|
|                        | PC1          | PC2          | PC3          | PC4          | PC1                 | PC2   | PC3          | PC4          |
| East expanding<br>(Ee) | <b>0.000</b> | <b>0.000</b> | 0.956        | <b>0.001</b> |                     |       |              |              |
| West expanding<br>(Ew) | <b>0.000</b> | <b>0.000</b> | <b>0.000</b> | 0.923        | <b>0.000</b>        | 0.358 | <b>0.003</b> | <b>0.001</b> |

**Table 3.3.** Niche overlap measure using the index  $D$  between evaluated *Phrynosoma* taxa, with those of *P. platyrhinos* shown for the Persistent, expanding eastern (Ee), and expanding western (Ew) groups.  $N$  represents the number of occurrence records used to build climatic niche models.

|                       | $N$ | <i>P. douglasii</i> | <i>P. hernandesii</i> | Ee   | Persistent | Ew |
|-----------------------|-----|---------------------|-----------------------|------|------------|----|
| <i>P. douglasii</i>   | 17  | x                   |                       |      |            |    |
| <i>P. hernandesii</i> | 41  | 0.49                | x                     |      |            |    |
| Ee                    | 25  | 0.27                | 0.53                  | x    |            |    |
| Persistent            | 165 | 0.12                | 0.07                  | 0.11 | x          |    |
| Ew                    | 65  | 0.32                | 0.32                  | 0.44 | 0.24       | x  |

**Figure 3.1.** (A) General sample sites of *Phrynosoma platyrhinos*, *P. goodei* (localities 7, 13–17, 102) and *P. platyrhinos* from Yuma Proving Grounds (locality 2); closely situated samples without intervening physical barriers are grouped for visual interpretation and discussion. Grey circles represent occurrence records based on museum specimens and grey shadings represent ecoregions as adapted from (Olson *et al.* 2001). (B) Occurrence records used in the niche similarity comparison of *P. platyrhinos* and parapatric species *P. douglasii* and *P. hernandesi*. The shaded polygons represent Persistent, expanding western (Ew), and expanding eastern (Ee) groups of *P. platyrhinos* (see text for details).

**Figure 3.2.** Maximum likelihood tree for combined ND4, tRNA and cytB sequences for 22 unique haplotypes of *Phrynosoma platyrhinos*, 2 haplotypes of *P. goodei*, and 3 haplotypes of *Phrynosoma platyrhinos* from Yuma Proving Grounds. *Phrynosoma mcallii* was used as an outgroup. The three major clades of *P. platyrhinos* (Western, Southeastern, and Northeastern) are indicated. Nodal support from nonparametric bootstrap values are shown (numbers above), as are posterior probabilities from Bayesian inference (numbers below). The locality number for each sample is in parentheses.

**Figure 3.3.** Median-joining network of concatenated ND4 and tRNA sequences for *Phrynosoma platyrhinos*, *P. goodei*, and *P. platyrhinos* from Yuma Proving Grounds. Circle size and shading reflect the number of samples exhibiting a given haplotype ranging from 1 to 4, with several abundant haplotypes indicated by large circles labeled

internally with the number of samples. The length of connection lines between haplotypes is proportional to the number of mutational changes, with the shortest connection line representing a single mutational change.

**Figure 3.4.** Distribution of major nested clades (left) identified from the median-joining network (right) of the concatenated ND4 and tRNA sequences for *Phrynosoma platyrhinos*, *P. goodei*, and *P. platyrhinos* from Yuma Proving Grounds. Pie graph sizes in the map reflect the sample size at each location progressing from smallest ( $N = 1$ ) to largest ( $N = 6$ ). The clade numbers and color correspond to those on the median-joining network. The polygons represent the extent of the contracted expanding haplotypes in the western Great Basin (Hw1 and Hw2 indicated in green), and those in the eastern Great Basin (He1 and He2 indicated in pink).

**Figure 3.5.** Bayesian skyline plot derived from concatenated ND4 and tRNA sequences of *Phrynosoma platyrhinos*. The x-axis shows the time and the y-axis shows an index of genetic diversity assuming a 2-year generation time. The black line is the median for genetic diversity and the grey area represents the 95% upper and lower highest posterior density limits. The plot is presented truncated on the right as the extended region showed no evidence of change in genetic diversity.

**Figure 3.6.** (A) Scatter plot for the first two principal components derived from 19 bioclimatic variables for the Persistent group (white dots), expanding eastern (Ee) group (black dots), and expanding western (Ew) group (grey dots) of *Phrynosoma platyrhinos*.

(B) Least square means for the first four principal components derived from 19 bioclimatic variables for the three groups of *P. platyrhinos*.

**Figure 3.7.** Climatic niche models of the climatic niche for the Persistent group (left column), expanding western (Ew) group (center column) and expanding eastern (Ee) group (right column) of *Phrynosoma platyrhinos*. Rows represent the current climatic condition (first row), and two reconstructions of climatic conditions at the Last Glacial Maximum — CCSM (second row) and MIROC (third row). The light areas represent predicted suitable habitats using the minimum presence threshold (light green) and 10% omission (darker green). Dark blue areas represent the extent of pluvial lakes (Raines *et al.* 1996).

**Figure 3.8.** Least square means of four bioclimatic variables for the expanding western, expanding eastern and Persistent groups of *Phrynosoma platyrhinos*.

Figure 3.1






Figure 3.2




Figure 3.3





Figure 3.4



**Figure 3.5**




Figure 3.6





Figure 3.7





Figure 3.8



## Bibliography

- Ackerly D (2003a) Canopy gaps to climate change - extreme events, ecology and evolution. *New Phytologist*, **160**, 2-4.
- Ackerly DD (2003b) Community assembly, niche conservatism, and adaptive evolution in changing environments. *International Journal of Plant Sciences*, **164**, S165-S184.
- Araujo MB, Guisan A (2006) Five (or so) challenges for species distribution modelling. *Journal of Biogeography*, **33**, 1677-1688.
- Arevalo E, Davis SK, Sites JW (1994) Mitochondrial DNA sequence divergence and phylogenetic relationships among 8 chromosome races of the *Sceloporus grammicus* complex (Phrynosomatidae) in central Mexico. *Systematic Biology*, **43**, 387-418.
- Bandelt HJ, Forster P, Rohl A (1999) Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution*, **16**, 37-48.
- Braconnot P, Otto-Bliesner B, Harrison S, *et al.* (2007) Results of PMIP2 coupled simulations of the Mid-Holocene and Last Glacial Maximum - Part 1: experiments and large-scale features. *Climate of the Past*, **3**, 261-277.
- Broennimann O, Treier UA, Muller-Scharer H, *et al.* (2007) Evidence of climatic niche shift during biological invasion. *Ecology Letters*, **10**, 701-709.
- Castoe TA, Spencer CL, Parkinson CL (2007) Phylogeographic structure and historical demography of the western diamondback rattlesnake (*Crotalus atrox*): A perspective on North American desert biogeography. *Molecular Phylogenetics and Evolution*, **42**, 193-212.
- Coetzee BWT, Robertson MP, Erasmus BFN, van Rensburg BJ, Thuiller W (2009) Ensemble models predict Important Bird Areas in southern Africa will become less effective for conserving endemic birds under climate change. *Global Ecology and Biogeography*, **18**, 701-710.
- Csuti BA (1979) *Patterns of adaptation and variation in the Great Basin kangaroo rat (Dipodomys microps)*. University of California Publications in Zoology 111.
- Davis MB, Shaw RG (2001) Range shifts and adaptive responses to Quaternary climate change. *Science*, **292**, 673-679.
- Davis MB, Shaw RG, Etterson JR (2005) Evolutionary responses to changing climate. *Ecology*, **86**, 1704-1714.



- Douglas ME, Douglas MR, Schuett GW, Porras LW (2006) Evolution of rattlesnakes (Viperidae; *Crotalus*) in the warm deserts of western North America shaped by Neogene vicariance and Quaternary climate change. *Molecular Ecology*, **15**, 3353-3374.
- Drummond AJ, Ho SYH, Rawlence HN, Rambaut A (2007) A rough guide to BEAST 1.4. ([http://beast-mcmc.googlecode.com/files/BEAST14\\_Manual\\_6July2007.pdf](http://beast-mcmc.googlecode.com/files/BEAST14_Manual_6July2007.pdf))
- Drummond AJ, Rambaut A (2007) BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology*, **7**, 214.
- Drummond AJ, Rambaut A, Shapiro B, Pybus OG (2005) Bayesian coalescent inference of past population dynamics from molecular sequences. *Molecular Biology and Evolution*, **22**, 1185-1192.
- Dumas PC (1964) Species-pair allopatry in genera *Rana* + *Phrynosoma*. *Ecology*, **45**, 178-181.
- Excoffier L, Foll M, Petit RJ (2009) Genetic consequences of range expansions. *Annual Review of Ecology Evolution and Systematics*, **40**, 481-501.
- Excoffier L, Laval G, Schneider S (2005) Arlequin (version 3.0): An integrated software package for population genetics data analysis. *Evolutionary Bioinformatics*, **1**, 47-50.
- Fitch HS (1985) *Variation in clutch and litter size in New World reptiles* University of Kansas, Lawrence, KS.
- Forster P, Torroni A, Renfrew C, Rohl A (2001) Phylogenetic star contraction applied to Asian and Papuan mtDNA evolution. *Molecular Biology and Evolution*, **18**, 1864-1881.
- Galbreath KE, Hafner DJ, Zamudio KR (2009) When cold is better: climate-driven elevation shifts yield complex patterns of diversification and demography in an alpine specialist (American pika, *Ochotona princeps*). *Evolution*, **63**, 2848-2863.
- Grayson DK (2000) Mammalian responses to Middle Holocene climatic change in the Great Basin of the western United States. *Journal of Biogeography*, **27**, 181-192.
- Hampe A, Petit RJ (2005) Conserving biodiversity under climate change: the rear edge matters. *Ecology Letters*, **8**, 461-467.
- Harrison SP (2000) Palaeoenvironmental data sets and model evaluation in PMIP, 9-25.

- Hasumi H, Emori S (2004) K-1 Coupled Model (MIROC) Description. Center for Climate System Research, University of Tokyo, Tokyo, Japan.
- Hewitt G (1993) Postglacial distribution and species substructure: Lessons from pollen, insects and hybrid zones. In: *Evolutionary patterns and processes* (eds. Lees DR, Edwards D), pp. 97-123. Linnean Soc. 14. Academic Press, London, U. K.
- Hewitt G (2000) The genetic legacy of the Quaternary ice ages. *Nature*, **405**, 907-913.
- Hewitt GM (1996) Some genetic consequences of ice ages, and their role in divergence and speciation. *Biological Journal of the Linnean Society*, **58**, 247-276.
- Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005) Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, **25**, 1965-1978.
- Hilbe JM (2008) Systat 12.2: An overview. *American Statistician*, **62**, 177-178.
- Hockett BS (2000) Paleobiogeographic changes at the Pleistocene-Holocene boundary near Pintwater Cave, southern Nevada. *Quaternary Research*, **53**, 263-269.
- Hoffmann MH (2005) Evolution of the realized climatic niche in the genus *Arabidopsis* (Brassicaceae). *Evolution*, **59**, 1425-1436.
- Holt RD, Keitt TH, Lewis MA, Maurer BA, Taper ML (2005) Theoretical models of species' borders: single species approaches. *Oikos*, **108**, 18-27.
- Hunter KL, Betancourt JL, Riddle BR, *et al.* (2001) Ploidy race distributions since the Last Glacial Maximum in the North American desert shrub, *Larrea tridentata*. *Global Ecology and Biogeography*, **10**, 521-533.
- Iverson JB, Balgooyen CP, Byrd KK, Lyddan KK (1993) Latitudinal variation in egg and clutch size in turtles. *Canadian Journal of Zoology*, **71**, 2448-2461.
- Jackson ST, Overpeck JT (2000) Responses of plant populations and communities to environmental changes of the late Quaternary. *Paleobiology*, **26**, 194-220.
- Jezkova T, Jaeger JR, Conrad P (2009a) Distribution and Habitat Associations of Short-horned Lizards in Nevada. In: 2009. *Species Investigations: Job Performance Report: State Wildlife Grant* eds. Atkinson SJ, Lamp R, Neel LA, Richards L, Tomlinson CR). Nevada Department of Wildlife, Diversity Division, Reno, NV.

- Jezkova T, Jaeger JR, Marshall ZL, Riddle BR (2009b) Pleistocene impacts on the phylogeography of the desert pocket mouse (*Chaetodipus penicillatus*). *Journal of Mammalogy*, **90**, 306-320.
- Jobb G, von Haeseler A, Strimmer K (2004) TREEFINDER: a powerful graphical analysis environment for molecular phylogenetics. *BMC Evolutionary Biology*, **4**.
- Jones KB (1995) *Phylogeography of the desert horned lizard (Phrynosoma platyrhinos) and the short-horned lizard (Phrynosoma douglassii): patterns of divergence and diversity* Dissertation, University of Nevada Las Vegas.
- Kass RE, Raftery AE (1995) Bayes Factors. *Journal of the American Statistical Association*, **90**, 773-795.
- Kearney M, Shine R, Porter WP (2009) The potential for behavioral thermoregulation to buffer "cold-blooded" animals against climate warming. *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 3835-3840.
- Kozak KH, Wiens JJ (2006) Does niche conservatism promote speciation? A case study in North American salamanders. *Evolution*, **60**, 2604-2621.
- Leache AD, McGuire JA (2006) Phylogenetic relationships of horned lizards (*Phrynosoma*) based on nuclear and mitochondrial data: Evidence for a misleading mitochondrial gene tree. *Molecular Phylogenetics and Evolution*, **39**, 628-644.
- Leache AD, Reeder TW (2002) Molecular systematics of the eastern fence lizard (*Sceloporus undulatus*): A comparison of parsimony, likelihood, and Bayesian approaches. *Systematic Biology*, **51**, 44-68.
- Luxbacher AM, Knouft JH (2009) Assessing concurrent patterns of environmental niche and morphological evolution among species of horned lizards (*Phrynosoma*). *Journal of Evolutionary Biology*, **22**, 1669-1678.
- Macey JR, Wang YZ, Ananjeva NB, Larson A, Papenfuss TJ (1999) Vicariant patterns of fragmentation among gekkonid lizards of the genus *Teratoscincus* produced by the Indian collision: A molecular phylogenetic perspective and an area cladogram for Central Asia. *Molecular Phylogenetics and Evolution*, **12**, 320-332.
- Merrill RM, Gutierrez D, Lewis OT, *et al.* (2008) Combined effects of climate and biotic interactions on the elevational range of a phytophagous insect. *Journal of Animal Ecology*, **77**, 145-155.

- Monasterio C, Salvador A, Iraeta P, Diaz JA (2009) The effects of thermal biology and refuge availability on the restricted distribution of an alpine lizard. *Journal of Biogeography*, **36**, 1673-1684.
- Montanucci RR (1981) Habitat separation between *Phrynosoma douglasii* and *Phrynosoma orbiculare* (Lacertilia: Iguanidae) in Mexico. *Copeia*, **1**, 147-153.
- Mulcahy DG (2008) Phylogeography and species boundaries of the western North American nightsnake (*Hypsiglena torquata*): Revisiting the subspecies concept. *Molecular Phylogenetics and Evolution*, **46**, 1095-1115.
- Mulcahy DG, Spaulding AW, Mendelson JR, Brodie ED (2006) Phylogeography of the flat-tailed horned lizard (*Phrynosoma mcallii*) and systematics of the *P. mcallii-platyrhinos* mtDNA complex. *Molecular Ecology*, **15**, 1807-1826.
- Murphy RW, Trepanier TL, Morafka DJ (2006) Conservation genetics, evolution and distinct population segments of the Mojave fringe-toed lizard, *Uma scoparia*. *Journal of Arid Environments*, **67**, 226-247.
- Newton MA, Raftery AE, Davison AC, *et al.* (1994) Approximate Bayesian inference with the weighted likelihood bootstrap. *Journal of the Royal Statistical Society Series B-Methodological*, **56**, 3-48.
- Nogues-Bravo D (2009) Predicting the past distribution of species climatic niches. *Global Ecology and Biogeography*, **18**, 521-531.
- Nylander JAA (2004) MrModeltest v.2. Evolutionary Biology Centre, Uppsala University, Uppsala, Sweden.
- Olson DM, Dinerstein E, Wikramanayake ED, *et al.* (2001) Terrestrial ecoregions of the worlds: A new map of life on Earth. *Bioscience*, **51**, 933-938.
- Otto-Bliesner BL, Brady EC, Clauzet G, *et al.* (2006) Last Glacial Maximum and Holocene climate in CCSM3. *Journal of Climate*, **19**, 2526-2544.
- Pavlik BM (1989) Phytogeography of sand dunes in the Great Basin and Mojave deserts. *Journal of Biogeography*, **16**, 227-238.
- Pearman PB, Guisan A, Broennimann O, Randin CF (2008) Niche dynamics in space and time. *Trends in Ecology & Evolution*, **23**, 149-158.
- Peterson AT, Martinez-Meyer E, Gonzalez-Salazar C (2004) Reconstructing the Pleistocene geography of the *Aphelocoma* jays (Corvidae). *Diversity and Distributions*, **10**, 237-246.

- Peterson AT, Nyari AS (2008) Ecological niche conservatism and Pleistocene refugia in the thrush-like mourner, *Schiffornis* sp., in the neotropics. *Evolution*, **62**, 173-183.
- Phillips SJ, Anderson RP, Schapire RE (2006) Maximum entropy modeling of species geographic distributions. *Ecological Modelling*, **190**, 231-259.
- Phillips SJ, Dudik M (2008) Modeling of species distributions with Maxent: new extensions and a comprehensive evaluation. *Ecography*, **31**, 161-175.
- Pianka ER, Parker WS (1975) Ecology of horned lizards: A review with special reference to *Phrynosoma platyrhinos*. *Copeia*, **1**, 141-162.
- Polzin T, Daneshmand SV (2003) On Steiner trees and minimum spanning trees in hypergraphs. *Operations Research Letters*, **31**, 12-20.
- Posada D, Buckley TR (2004) Model selection and model averaging in phylogenetics: Advantages of akaike information criterion and Bayesian approaches over likelihood ratio tests. *Systematic Biology*, **53**, 793-808.
- Raines GL, Sawatzky DL, Connors KA (1996) Great Basin geoscience data base. In: *U.S. Geological Survey Digital Data Series DDS-41*.
- Rambaut A, Drummond AJ (2007) Tracer v1.4. <http://beast.bio.ed.ac.uk/Tracer>.
- Rissler LJ, Apodaca JJ (2007) Adding more ecology into species delimitation: Ecological niche models and phylogeography help define cryptic species in the black salamander (*Aneides flavipunctatus*). *Systematic Biology*, **56**, 924-942.
- Rodder D, Lotters S (2009) Niche shift versus niche conservatism? Climatic characteristics of the native and invasive ranges of the Mediterranean house gecko (*Hemidactylus turcicus*). *Global Ecology and Biogeography*, **18**, 674-687.
- Roe BA, Ma DP, Wilson RK, Wong JFH (1985) The complete nucleotide sequence of the *Xenopus laevis* mitochondrial genome. *Journal of Biological Chemistry*, **260**, 9759-9774.
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics*, **19**, 1572-1574.
- Sears MW, Angilletta MJ (2004) Body size clines in *Sceloporus* lizards: Proximate mechanisms and demographic constraints. *Integrative and Comparative Biology*, **44**, 433-442.

- Spaulding AW (1990) Vegetational and climatic development of the Mojave Desert: The last glacial maximum to the Present. In: *Packrat middens: The last 40,000 years of biotic change* (eds. Betancourt JL, Van Devender TR, Martin PS), pp. 200-239. The University of Arizona Press, Tucson.
- Suchard MA, Weiss RE, Sinsheimer JS (2001) Bayesian selection of continuous-time Markov chain evolutionary models. *Molecular Biology and Evolution*, **18**, 1001-1013.
- Tanner WW (1999) Distribution of the species *Phrynosoma platyrhinos* in Utah. *Great Basin Naturalist*, **59**, 295-296.
- Templeton AR, Boerwinkle E, Sing CF (1987) A cladistic analysis of phenotypic associations with haplotypes inferred from restriction endonuclease mapping. I. Basic theory and an analysis of alcohol dehydrogenase activity in *Drosophila*. *Genetics*, **117**, 343-351.
- Templeton AR, Sing CF (1993) A cladistic analysis of phenotypic associations with haplotypes inferred from restriction endonuclease mapping. IV. Nested analyses with cladogram uncertainty and recombination. *Genetics*, **134**, 659-669.
- Thuiller W, Albert C, Araujo MB, *et al.* (2008) Predicting global change impacts on plant species' distributions: Future challenges. *Perspectives in Plant Ecology Evolution and Systematics*, **9**, 137-152.
- Urban MC, Phillips BL, Skelly DK, Shine R (2007) The cane toad's (*Chaunus* [*Bufo*] *marinus*) increasing ability to invade Australia is revealed by a dynamically updated range model. *Proceedings of the Royal Society B-Biological Sciences*, **274**, 1413-1419.
- Varela S, Rodríguez J, Lobo JM (2009) Is current climatic equilibrium a guarantee for the transferability of distribution model predictions? A case study of the spotted hyena. *Journal of Biogeography*, **36**, 1645-1655.
- Waltari E, Hijmans RJ, Peterson AT, *et al.* (2007) Locating Pleistocene refugia: comparing phylogeographic and ecological niche model predictions. *PLoS ONE*, **2**, e563.
- Warren DL, Glor RE, Turelli M (2008) Environmental niche equivalency versus conservatism: quantitative approaches to niche evolution. *Evolution*, **62**, 2868-2883.
- Wiens JJ, Graham CH (2005) Niche conservatism: integrating evolution, ecology, and conservation biology. *Annual Review of Ecology Evolution and Systematics*, **36**, 519-539.

Zamudio KR, Jones KB, Ward RH (1997) Molecular systematics of short-horned lizards: biogeography and taxonomy of a widespread species complex. *Systematic Biology*, **46**, 284-305.

## CHAPTER 4

### SOME MORE CHALLENGES FOR ECOLOGICAL NICHE MODELS: EXPLORING THE CHALLENGES OF TRANSFERAL MODELING IN BIOGEOGRAPHY

#### Abstract

I expand on challenges for ecological niche modeling (ENM) discussed by Araújo & Guisan (2006) into the realm of transferal modeling. Transferal modeling involves projecting ecological niche models (ENMs) onto different geographic areas or different time periods and has been used for predicting species responses to global warming, reconstructing palaeo-ranges of organisms, or modeling the spread of invasive species. Transferability of models in time and space however has been widely questioned because of conceptual and algorithmic assumptions and uncertainties associated with the methodology. Herein, I discuss several high-priority challenges in transferal modeling: (1) model-based variability in climatic reconstructions; (2) selection of variables; (3) projection outside the calibration range; (4) niche shifting; and (5) transferal model evaluations. Through theoretical background and sets of experiments, I demonstrate how these challenges can affect resulting models and, when possible, offer suggestions on how uncertainties might be diminished. I focus mainly on projections in time to climatic conditions of the last glacial maximum, but most of the issues are applicable when projecting to other time periods or in space. Our intent is to provide better understanding of conceptual and algorithmic methodology behind transferal modeling in order to facilitate critical evaluation of existing models, and to stimulate further exploration of these and other challenges associated with transferal modeling.



Ultimately, I hope our experiments and discussion will provide for general improvements in the development of transferal models.

## Introduction

Ecological niche models (ENMs, also referred to as climate envelope models [CEMs], species distribution models [SDMs], or bioclimatic models) are important tools for biogeographers (Araújo & Guisan 2006; Elith *et al.* 2006; Guisan & Zimmermann 2000; Jeschke & Strayer 2008; Peterson & Soberon 2005). Araújo & Guisan (2006) explored several of the fundamental challenges inherent in use of ecological niche modeling, including: (1) niche concepts; (2) data sampling and model building; (3) model parameterization; (4) model selection and predictor contribution; and (5) model evaluation. An additional topic not explored by Araújo & Guisan (2006) concerns the use of ENMs to predict species distributions across geographic space or under changing environmental conditions through time. Herein, I address the challenges related to use of ENMs in the realm of *transferal modeling*.

Transferal modeling involves projecting ENMs onto different geographic areas, *i.e.* transferal in space, often used for modeling invasive species (Fitzpatrick *et al.* 2007; Peterson *et al.* 2003; Randin *et al.* 2006) or onto a focal geographic area under past or future environmental conditions, *i.e.* transferal in time, often used for studying species responses to global climate change, or reconstructing palaeo-ranges of organisms (Carnaval & Moritz 2008; Carstens & Richards 2007; Martinez-Meyer *et al.* 2004; Pearson *et al.* 2006; Peterson *et al.* 2004; Peterson & Nyari 2008; Waltari & Guralnick

2009; Waltari *et al.* 2007). A typical approach to transferal modeling entails building (calibrating) an ecological niche model (ENM) using one of the available statistical methods (e.g. logistic regression, climatic envelope, neural network, or generic algorithms) that utilize presence-only or presence-absence records of species occurrence and a set of environmental variables [for summary see Elith *et al.* (2006) and Jeschke & Strayer (2008)]. Using the rules developed during *calibration*, the ENM is then projected onto a set of environmental variables representing a different area or time period.

Transferability of models in space and time has been widely questioned because of real or perceived obstacles, including: *extrapolation* of environmental variables, *transferability* of variables, existence of non-analogous climates between two time periods, existence of ecotypes or phenotypic plasticity within a modeled taxon, changes in biotic interactions in space and time, or inconsistencies among modeling approaches (Araujo & New 2007; Fitzpatrick *et al.* 2007; Jeschke & Strayer 2008; Nogues-Bravo 2009; Pearman *et al.* 2008; Pearson *et al.* 2002; Randin *et al.* 2006). Below, I discuss several of these high-priority challenges and show how they might affect (or bias) resulting models. I mainly focus on *projections* in time, in particular to climatic conditions of the last glacial maximum (LGM), but most of the issues I address are applicable when projecting in space or to other time periods. When possible, I offer suggestions on how particular uncertainties might be diminished, although I make no claim that such approaches will suffice for any particular dataset or question. I do not address general ENM challenges or comparisons among different modeling approaches,

as these issues have been reviewed elsewhere (Araujo & Guisan 2006; Elith *et al.* 2006; Peterson *et al.* 2007; Phillips 2008; Phillips *et al.* 2006). Although most of our analyses were done using the program Maxent 3.2 (Phillips *et al.* 2006; Phillips & Dudik 2008; Phillips *et al.* 2004), which has been reported to have better predictive accuracy than other current approaches (Elith *et al.* 2006; Phillips 2008; Phillips *et al.* 2006; Phillips & Dudik 2008), most issues discussed herein are applicable, or at least relevant to, other modeling methods, e.g., Desktop GARP (Stockwell & Peters 1999), BIOCLIM (Hijmans *et al.* 2005b).

## Methods

The ENMs (used in Challenges 1-4) were built in the software package MAXENT v. 3.2.1 (Phillips *et al.* 2006), a program that calculates relative probabilities of the species' presence in the defined geographic space (Phillips *et al.* 2004). I used the default parameters in MAXENT (500 maximum iterations, convergence threshold of 0.00001, and regularization multiplier of 1, 10,000 background points) with the application of random seed and logistic probabilities (approximating niche suitability) for the output (Phillips & Dudik 2008). Occurrence records for individual species were obtained from HerpNet (<http://herpnet.org>) or MaNIS (<http://manisnet.org>). All records missing geographic coordinates, lacking a value for geographic uncertainty, and records with geographic uncertainty greater than 5 km were excluded. Current climate was represented (unless otherwise stated) by 19 bioclimatic variables (herein referred to as Bio 1 - 19) from the WorldClim dataset v. 1.4 with resolution of 2.5 minutes

(<http://www.worldclim.org/>; Hijmans *et al.* 2005a). The bioclimatic variables are derived from monthly temperature and precipitation data, and represent biologically meaningful aspects of climate (Jezkova *et al.* 2009; Waltari *et al.* 2007). For environmental layers representing the climatic conditions of the LGM, I used the Community Climate System Model v. 3 (CCSM) and the Model for Interdisciplinary Research on Climate v. 3.2 (MIROC; see below for more details).

#### Challenge 1: Model-based variability in climatic reconstructions

When projecting in space, the environmental variables used in the new area are usually of a similar quality to those used for calibration, *i.e.* the same global dataset can be used for calibration on one continent and projected onto another continent. However, when projecting to a different time period (past or future) the projection variables are often derived very differently from those used for calibration. For example, current climatic (temperature and precipitation) conditions are derived directly from data recorded at numerous worldwide weather stations (Hijmans *et al.* 2005a) and then interpolated over the study region, while the past (and future) climatic conditions are derived using sophisticated simulations under specific assumptions. The current set of coupled ocean-atmosphere simulations for the climatic conditions of the last glacial maximum (LGM: ca 21,000 calendar yr B.P., equivalent to ca 18,000 14C yr B.P.; Harrison 2000) is available through the Paleoclimatic Modeling Intercomparison Project (PMIP phases II; Braconnot *et al.* 2007). PMIP has established a protocol followed by participating modeling groups for LGM simulations regarding concentration of

greenhouse gases, ice sheet coverage, insolation, change in topography (mainly caused by lowering sea levels), etc. Based on this general protocol, more than 15 different climatic models for the LGM are currently available from various modeling groups (<http://pmip2.lsce.ipsl.fr/>; Braconnot *et al.* 2007).

Simulation of the LGM started as an experiment to examine climate response to the presence of ice sheets and lowered greenhouse gas (mainly CO<sub>2</sub>) concentrations in order to provide a credibility test for different future scenarios of increased CO<sub>2</sub> (Otto-Bliesner *et al.* 2006). Increasingly, climatic variables derived from these simulations are being used to reconstruct species paleo-distributions and evaluate range shifts, contractions, and expansions associated with climatic changes since the LGM (Carstens & Richards 2007; Jezkova *et al.* 2009; Martinez-Meyer *et al.* 2004; Peterson *et al.* 2004; Peterson & Nyari 2008; Rodriguez-Robles *et al.* 2010; Waltari *et al.* 2007). Of the 15 commonly available climatic models, the following two are most often utilized for transferal modeling: Community Climate System Model Version 3 (CCSM; Otto-Bliesner *et al.* 2006) with a resolution of 1°, and the Model for Interdisciplinary Research on Climate Version 3.2 (MIROC; Hasumi & Emori 2004) with an original spatial resolution of 1.4° x 0.5° (Braconnot *et al.* 2007). The original climatic variables used in these models have been downscaled to the spatial resolution of 2.5 minutes (Peterson & Nyari 2008; Waltari & Guralnick 2009; Waltari *et al.* 2007) under the assumption that changes in climate are relatively stable over space (high spatial autocorrelation; Hijmans *et al.* 2005a).

The CCSM and MIROC climatic models provide quite different reconstructions of the climatic conditions for the LGM. For example, Figure 4.1 shows a single bioclimatic variable, Annual Precipitation, reconstructed under each climatic model over a portion of western North America (Fig. 4.1A,B), and the absolute difference for this variable between these two models (Fig. 4.1C). The greatest inconsistencies between models are centered within the southern Sonoran Desert (northwestern Mexico) and western Sierra Nevada. Differences between these two climatic models might explain why resulting transferal models of species climatic niches are often quite different. For example, Figure 4.1 also shows a ENM for Merriam's kangaroo rat (*Dipodomys merriami*) projected onto the CCSM (Fig. 4.1D) and MIROC (Fig. 4.1E) models (calibration model and data not shown) using the 19 bioclimatic variables (Hijmans *et al.* 2005a; Waltari *et al.* 2007). The absolute difference in logistic probabilities between these two models (Fig. 4.1F) shows a major area of inconsistency in the Sonoran Desert, in the same region as the area of inconsistency for Annual Precipitation (Fig. 4.1C).

Previous studies that used more than one climatic model for transferal modeling approached these inconsistencies differently. Some authors reconciled discrepancies by summing, intersecting, or averaging all models (e.g.; Waltari & Guralnick 2009; Waltari *et al.* 2007); whereas others presented each model separately (e.g.; Peterson & Nyari 2008; Rodriguez-Robles *et al.* 2010). I currently have no information about the accuracies of the past and future climatic models, and it is likely that model accuracy varies across space. I suggest that the purpose of each specific study should drive the decision on how the model-based variability is treated. For example, if the purpose of

the study is to identify “hotspots” for conservation purposes based on species persistence through time, the intersection of all ENMs can be used to increase confidence in area selection, but when assessing the area of impact for an emerging pathogen or invasive species, all possibilities (sum of all models) might be of interest. Additionally, if the generation of testable hypotheses is the objective (e.g., for evolutionary or phylogeographic studies) models might be considered separately and each tested using independent approaches.

#### Challenge 2: Selection of variables

Selection of environmental variables to be used in ecological niche modeling has been a center of debate, with concerns about how to target environmental variables that actually define a species range, the number of environmental variables that should be used, and the influence of over-parameterization (too many variables) on model performance (Nogues-Bravo *et al.* 2008; Peterson *et al.* 2007; Stockwell 2006). In transferal modeling, variable selection faces an additional challenge since some variables are less transferable than others because of unstable covariation with other variables through time (Nogues-Bravo 2009). For example, elevation (which is known to limit the current distributions of many species) correlates with temperature and precipitation differently at different time periods. In particular, certain elevations in the Great Basin of North America were associated with cooler temperatures and higher precipitation during the LGM relative to current conditions. As a result, these *non-analogous climatic conditions* (i.e. variable combinations nonexistent during the

calibration) might not be correctly projected simply because I have no information in terms of presence/absence of the species under these conditions during the calibration. The presence of non-analogous climates likely results in the underestimation of suitable habitat in transferal models (Nogues-Bravo 2009).

To document this effect, I built ENMs for the Great Basin pocket mouse (*Perognathus parvus*), an endemic species from the Great Basin that in the fossil record shows elevational shifts as a response to late Pleistocene/Holocene climatic changes (Grayson 2000). The models were created using the 19 bioclimatic variables, first without elevation included as a variable, and then with elevation included. While the OK (calibration) models built with and without elevation do not differ substantially (Fig. 4.2A,B), the two transferal LGM models (MIROC) were very different from each other. The inclusion of elevation (Fig. 4.2C) resulted in approximately a two-third reduction in predicted suitable habitat from the model with elevation omitted (Fig. 4.2D). Additionally, histograms of predicted suitable elevations throughout the area (in this example with logistic probabilities higher than 0.2) revealed that only the model excluding elevation predicted species shift to lower elevations during the LGM, in accord with evidence from the fossil record (Fig. 4.3). Although this example might seem trivial, the lack of transferability in other variables might not be so obvious.

The transferability of variables can be partly evaluated in Maxent using the jackknife plot of *test gain* and area under curve (AUC) on test data (see the Maxent manual for better understanding of these methods) because variables that do not perform well on the test data (test data being occurrence points set aside during calibration for



subsequent evaluation of the model) are not likely to perform well when transferred in space or time. Covariance among variables at different time periods can also be compared using standard statistical approaches (Nogues-Bravo 2009). Finally, program algorithms that utilize covariance between environmental variables for deriving ENMs often can be modified and certain features can be excluded (e.g. Product features in Maxent may be one of these) when non-analogous climatic conditions are of concern (Steven J. Phillips, personal communication).

### Challenge 3: Projection outside the calibration range (Extrapolation)

A common issue associated with transferal modeling and is that the calibration layers often have a different range of values than the transferal (projection) layers (Randin *et al.* 2006; Thuiller *et al.* 2004). For example, the climatic conditions of the LGM in western North America are believed to have been colder and wetter than those of today. Under such conditions, I would expect the values of associated climatic variables to be shifted towards lower temperatures and higher precipitation. In particular, 17 of 19 bioclimatic variables within western North America have projection (LGM) values outside the *calibration* (OK) *range* (Table 4.1). Consequently, the habitat suitability across values absent from the calibration dataset cannot be directly projected and some extrapolation method must be employed to complete the projection.

Figure 4.4 shows how different extrapolation methods, as well as the shape of a response curve within the calibration range, effect probability values (representing occurrence probability) within the *projection range*. Both graphs in this figure represent

a unimodal response of a hypothetical organism to a particular environmental variable.

In the left graph, the calibration layer captures the unimodal distribution of the response curve but is missing the right tail (high values) of the projection layer. To complete the projection, I employ three different extrapolation methods. Line A represents a nonlinear regression method and results in an extrapolated curve approximate to the real response curve. Line B represents a method employed by Maxent that truncates the projection layer to the same extent as the calibration layer, with all values outside the calibration range set to the maximum or minimum value of the calibration layers. This latter method results in an overestimate (although not severe) of the probability values outside the calibration range. Line C is the most conservative approach, where probability is set to zero outside the calibration range, which results in an underestimate of occurrence probability. Overall, however, none of these extrapolation methods resulted in severe errors under the hypothetical conditions shown.

A more severe concern is shown in the right graph of Figure 4.4. This scenario shows a unimodal response curve within the calibration range that does not reach the peak, giving the illusion during the calibration that the response to this variable is linear (when in reality it is unimodal). A nonlinear regression (line A) results in a monotonic increase of suitability throughout the projection range, whereas the truncation method (line B) maintains equally high suitability. Both these methods result in increasing overestimates of habitat suitability towards maximum values of the variable. In Maxent, this case can be detected as large areas of unrealistically high probabilities, often in areas with

extreme climatic conditions. Setting the probability to zero outside the calibration range (Line C) has an opposite trend and underestimates values close to the calibration range. This method avoids the strong monotonic increase of probabilities described above (Randin *et al.* 2006), but overall, none of these extrapolation methods was able to approximate the true response of our hypothetical variable.

One way to decrease the amount of extrapolation in transferal modeling is to maximize the range of each variable during the calibration process. This can be achieved when the area for calibration is larger than the area used for projection (Pearson *et al.* 2002). For example, only 7 (for CCSM) and 9 (for MIROC) out of 19 bioclimatic variables will have a projection range outside the calibration range when models are calibrated using climatic variables extending across the entire world and projected onto the variables clipped to an area of western North America (Table 4.1B). The extrapolation differences between calibrating and projecting on the same extent versus calibrating using a much broader area can be evaluated in Maxent. As mentioned above, Maxent restricts the projected variables to the range of values encountered during training (i.e. a technique called *clamping*). The clamping values generated in Maxent give the absolute change in logistic probability caused by the clamping process. To demonstrate how maximizing the range of each variable during the calibration affects clamping values, I generated ENMs for the striped whipsnake (*Masticophis taeniatus*). Figure 4.5A shows where clamping occurred when calibration and projection variables were clipped to western North America. Figure 4.5B shows how clamping values decreased when the model was calibrated using bioclimatic variables from the entire world and

then projected on the truncated western North American dataset. However, large *commission errors* can be a problem with maximizing the area during calibration when utilizing pseudo-absences (the method utilized in Maxent) because the absence data are randomly taken from the entire calibration area (Pearson *et al.* 2002; Randin *et al.* 2006; Thuiller *et al.* 2004). For example, when I modeled the striped whipsnake for LGM using the information reflected in Figure 4.5B (World to western North America) the resulting predicted suitability area was over 40% larger than that modeled using information reflected in Figure 4.5A (western North America to western North America; data and models not shown).

Another option to decrease the amount of extrapolation is to exclude variables that have a projection range that extends extensively beyond the calibration range. In our example, when I exclude all variables that have a projection range for the MIROC model more than 10% beyond the calibration range (Bio 1, 5, 6, 8, 9, 10, 11, 13, 14, 17, 18, and 19; Table 4.1A), the remaining seven variables do not require any extrapolation and consequently clamping values remain zero. The disadvantage is that in many cases, too many variables are often discarded (in our example, 12 from the MIROC model and 16 from the CCSM model).

Importantly, the level and extent of clamping does not necessarily equal the error in the ENM caused by extrapolation. As shown in Figure 4.4, a variable with a unimodal response curve that is captured within the calibration range will likely not result in a severe error. I experienced this while developing models for the common side-blotched lizard (*Uta stansburiana*). Figure 4.6 shows a set of response curves from 19 bioclimatic

variables for this species produced by Maxent (model shown in Figure 4.7). Five variables (Bio 1, 3, 4, 6, and 11) have a unimodal response curve and all values outside the calibration range will have a probability value close to zero. Six variables (Bio 2, 5, 8, 9, 10, and 18) have a non-zero probability value (herein referred to as an *open end*) on the upper end of the value ranges. The remaining nine variables (Bio 7, 12, 13, 14, 15, 16, 17, 18, and 19) have non-zero probability values on the lower end. Extrapolation of these variables can result in overestimates of probabilities (as seen in Fig. 4.4); however, when these response curves are compared with the CCSM value ranges in Table 4.1, only a single variable (Bio 2) has an open-end where extrapolation is needed (*i.e.* upper values). Figure 4.7 shows CCSM models for the common side-blotched lizard when all 19 bioclimatic variables were included (Fig. 4.7A, and associated clamping values 7b) and with the variable Bio 2 (mean diurnal temperature range) excluded (Fig. 4.7C, and associated clamping values 4.7D). The model (Fig. 4.7A) shows high probabilities within the northern and north-eastern areas that are biologically unrealistic (these areas were under ice during the LGM and could not have been habitat for this species); also notice the high clamping values in Fig. 4.7B. Excluding Bio 2 improves the model considerably with the large overpredicted areas disappearing (Fig. 4.7C) and clamping values reduced (Fig. 4.7D).

To assess and minimize the problems of extrapolation, I recommend exploring a range of values and response curves for the calibration and projection variables. If necessary, I suggest decreasing the need for extrapolation by excluding problematic variables, using asymmetrical geographic extent of variables (as described above), or

masking the models to the area of interest. If extrapolation is still required, I suggest exploring various extrapolation methods (which are usually unique to each modeling algorithm) and deciding on which method is most appropriate given the response curves for the variables under consideration.

#### Challenge 4: Ecological niche shift

The vast majority of transferal models are built under the assumption of niche conservatism (Warren *et al.* 2008; Wiens & Graham 2005), which means that the niche currently occupied by a species is identical to the niche that was occupied in the past and that will be occupied in the future. In reality, the assumption of niche conservatism will almost always be violated as species niches shift even over short periods of time (Nogues-Bravo 2009). Evolutionary adaptations in response to the changing environment can be responsible for niche shifts, in which case the fundamental niche of the species is altered (Fig. 4.8). Not all niche shifts, however, result from evolutionary adaptations. In many cases, the realized niche of a species shifts through time while the species fundamental niche remains stable (Fig. 4.8; Pearman *et al.* 2008). As indicated above, realized niche shifts can result from shifts in the realized environment, when non-analogous climatic conditions occur between two time periods (Ackerly 2003a; Ackerly 2003b; Hoffmann 2005; Jackson & Overpeck 2000; Nogues-Bravo 2009). Such niche shifts do not require plastic or evolutionary adaptive change but non-analogous climatic conditions may be omitted during model projection. A species may also occupy different portions of its fundamental niche (i.e. shift their realized niche) because of

varying biotic interactions through time or because of plasticity in various (behavioral, phonological, morphological physiological) traits that allow them to adjust to changing environments and persist in place despite environmental changes (Gibbs & Breisch 2001; Huppopp & Huppopp 2003; Post *et al.* 1999; Smith & Betancourt 2003).

Niche shifts have been documented from fossil records as well as from direct observations of species responses to recent climatic change. For example, only 14 out of 28 species showed elevational shifts in response to warming climate within the last century in California (Moritz *et al.* 2008), and only 3 out of 19 in northern Nevada (Rowe *et al.* 2010). Fossil records show that cooling climates lead to larger body mass in woodrats (genus *Neotoma*), consistent with Bergman's Rule, that allowed them to persist *in situ* throughout the oscillating climate of the late Pleistocene, rather than track their niche up and down in elevation, or north and south, as would be predicted if their niches remained conserved (Smith & Betancourt 2003). Similarly, fossil records of spotted hyena dated to the last interglacial were discovered within areas that exhibited substantially different environmental conditions than those the species inhabits today (Varela *et al.* 2009). Niche shifts have been documented in several species that expanded northwards after the LGM and that show novel adaptations to the newly encountered environment (Davis & Shaw 2001; Davis *et al.* 2005) and niche shifts are also quite common in invasive species where the environmental conditions and biotic interactions often differ between native and introduced ranges (Broennimann *et al.* 2007; Colautti *et al.* 2010; Rodder & Lotters 2009; Urban *et al.* 2007). Species currently occupying the driest and hottest areas within the Mojave desert of western North

America (e.g. Death Valley) likely experienced a niche shift at the end of the last glacial period as the current extreme conditions apparently had no equivalents during the LGM (Spaulding 1990). The average annual temperature within the low elevational parts of the Mojave desert is currently 7-8° C warmer than during the LGM and these areas are currently dominated by novel plant communities, including the dominant and widespread creosote bush - white bursage (*Larrea-Artemisia*) desert scrub, that did not develop until Holocene times (Hunter *et al.* 2001; Spaulding 1990). Following the warming climate at the end of the last glacial period, the species living on the valley bottoms of the Mojave desert did not shift upwards as would be predicted under the assumption of niche conservatism, but rather persisted and adapted to the newly developing, water-stressed environment, possibly through changes in their physiology, diet, and behavior (Tracy & Walsberg 2002).

As an example, I demonstrate the problem of niche shifts on four species (*Lemmiscus curtatus*, *Marmota flaviventris*, *Neotoma lepida*, *Tamias minimus*) by reconstructing their current and LGM climatic niches using ENMs. I used fossil records of the target species from Faunmap (Graham *et al.* 1994) identified as Full Glacial (ca. 14,500-20,500 14C yr B.P.) that should roughly correspond to the climatic conditions of the LGM (Waltari & Guralnick 2009). Values from 19 bioclimatic variables were extracted from both the CCSM and MIROC LGM models and compared with extracted variables from current observation records. The dimensionality of the dataset was reduced using principal component analysis (PCA) and the factor scores from the first two principal components were plotted (Fig 4.9). The results show that some historical records (LGM)



for three of the species (*L. curtatus*, *M. flaviventris*, and *T. minimus*; Fig. 4.9A,B,D) fall outside the ranges for current conditions (0K) for at least one principal component. Furthermore, the LGM records of *N. lepida* as well as one record of *T. minimus* are characterized by principle component scores on the very edge of those derived from current climatic niches (Fig. 4.9C,D). Only two LGM records of *M. flaviventris* fall well within the values representing the current climatic niche (Fig. 4.9B). When ENMs for these four species were constructed and projected on reconstructions of the LGM, many of the fossil occurrence records were outside or on the edge of the climatic niche interpreted as suitable during LGM (in agreement with the PCA analysis; Fig. 4.10).

Niche shifts are tied directly to the previous two challenges and in some cases can be alleviated by extrapolation. Extrapolation, however, is not necessarily biologically meaningful and can result in nonrealistic predictions as seen in Fig. 4.7. Nogues-Bravo (2009) strongly suggested that researchers always test for presence of non-analogous climates between the two evaluated time periods. Fossil records from different time periods can also be reviewed and a potential niche reconstructed by summing the niches occupied during all time slices (Nogues-Bravo *et al.* 2008). Finally, presence of niche shifts may be revealed from genetic data as populations that shifted their niche *in situ* and persisted in place will exhibit a different genetic signal than populations that shifted their distributions in concert with the oscillating climate (see chapter 2).

## Challenge 5: Transferal model evaluations

Model evaluation is important (Jeschke & Strayer 2008; Nogues-Bravo *et al.* 2008) but unlike current models (and to some extent transferal models in space) that can be evaluated by splitting the data into training and testing datasets, evaluation of models projected in time must be done using independent and often surrogate data. Herein I discuss three evaluation approaches and their possible drawbacks – (1) population genetic data, (2) direct fossil records, and (3) indirect pollen and plant macrofossil records representing biome changes.

### Genetics

Genetic signals imprinted in species genomes has been shown to trace species geographic and demographic histories (Excoffier *et al.* 2009; Hewitt 2000; Hewitt 1996; Hewitt 2004), and various molecular markers have been used to evaluate processes such as range shift, range expansion, range contraction, population expansion or population bottlenecks (Avice 2000). Recently, genetic markers also have been used to evaluate *a priori* hypotheses based on transferal models (Carstens & Richards 2007; Waltari *et al.* 2007). Although genetic analyses can be valuable for assessing transferal models, there are several issues that must be considered to validate this approach.

First, pronounced phylogeographic structure, in the form of genetic divergence among geographically separated populations dated to the mid or late Pleistocene, often has been attributed to climatic changes, providing evidence that during certain climatic periods (e.g. glacial periods) populations have been fragmented in two or more isolated refugia that persisted long enough to generate a phylogeographic signal (Knowles 2001).

In such cases, the transferal model is expected to predict isolated patches of suitable habitat for the climatic conditions representing the glacial period. Genetic divergence, however, can originate without any obvious geographic or ecological barriers (Graham *et al.* 2006; Irwin 2002; Jansson 2003; Losos *et al.* 2006; Neigel & Avise 1993), even when habitat remains relatively contiguous through time. This is particularly true for mtDNA which may exhibit rapid and stochastic lineage sorting (Avise 2000). Divergence without geographic barriers can be particularly pronounced in species with low mobility and small effective population sizes, where genetic drift can override gene flow (Irwin 2002; Rodriguez-Robles *et al.* 2010). Consequently, genetic divergence (especially when only a single molecular marker is being evaluated) does not necessarily indicate former presence of a barrier of unsuitable habitat among isolated refugia. To distinguish between stochastic divergence and that caused by past habitat fragmentation, multiple independent molecular markers and, ideally, multiple co-distributed taxa with similar ecological niches should be evaluated in search of congruent patterns of molecular and geographic divergence.

Second, extinction of clades is almost impossible to track using genetic information from extant taxa (Calvignac *et al.* 2008) which can complicate the use of genetic data. Let us assume a situation (Fig. 4.11) where a transferal model suggests presence of a hypothetical organism during an initial time period in the past (T1) in two separate refugia (Areas 1 and 2). The genetic analysis at current time (T3), however, recovers relatively high genetic diversity within Area 1 (indicative of some persistence of the organism within this area) while that for Area 2 exhibits very low genetic diversity with

most (all) individuals possessing genotypes that are also present in Area 1. One possibility (Fig. 4.11, scenario A) is that the transferal model incorrectly predicts persistence of the species within Area 2 which was in reality not suitable during the initial period (T1) – Area 2 was consequently colonized (during T2) from the genotypes present in Area 1. The second possibility (Fig. 4.11, Scenario B) is that the Area 2 was occupied during the past (T1) by the species (as suggested by a transferal model) but subsequently the original population (or at least the genotypes) went extinct and Area 2 was, again, colonized from Area 1 during some subsequent time (T2). In both cases, the genetic structure recovered currently (T3) would be similar (identical) but the histories (and consequently the suitability of Area 2 at T1 - which is of interest) would be different. In summary, an extinction event can destroy a genetic signal of species persistence and can be incorrectly interpreted as an error in the transferal model.

The third issue with using genetic data to evaluate transferal models is pseudo-congruence between the models and genetic data. Persistence of a taxon within an area, divergence of populations into refugia, as well as new colonization of a species into a previously uninhabited area can seemingly be supported by genetic data, while the genetic structure could have been generated during different times and for different reasons. Time of divergence among and within populations can be calculated from genetic data based on genetic divergence (typically percent sequence divergence) and a known, or estimated, mutation rate (Bromham & Penny 2003). These time estimates can then allow genetic patterns to be equated to the time period represented by the transferal model. Genetic divergence, however, greatly depends on effective population

sizes and on population size fluctuations through time (Edwards & Beerli 2000) and is highly stochastic due to the stochastic nature of genetic drift (Carstens *et al.* 2005; Knowles & Maddison 2002; Maddison & Knowles 2006). Also, mutation rate can be difficult to estimate as it is variable among different genes, among organisms (Spradling *et al.* 2001), and through time (Ho *et al.* 2005). As a result, confidence intervals around divergence times derived from genetic data often widely span the time period for which the transferal model was built.

### Fossil record

When appropriately dated, direct fossil records of the target organisms can document presence, or imply absence, of a species and can be used in the evaluation of transferal models (Martinez-Meyer *et al.* 2004; Waltari & Guralnick 2009). There are several drawbacks and challenges that should be considered when using fossil records for evaluation. First, fossils records can be dated with precision from few hundred to thousands, or tens of thousands, of years, while transferal models usually represent a climatic state for a particular time period (e.g. height of the LGM). Under the periods of time represented by the fossil record, an area could have experienced significant climatic fluctuations (Grootes & Stuiver 1997; Thompson *et al.* 1993) and a particular fossil might not represent presence of the species exactly at the time during which the environmental conditions are reflected in the transferal model.

Second, fossil identification may represent a substantial problem. Some species are difficult to identify to species level. For example, pocket mice in the family Heteromyidae comprise two reciprocally monophyletic and deeply divergent genera,

*Perognathus* and *Chaetodipus*, yet are not recognized as separate genera in the fossil record (Alexander & Riddle 2005; Hafner *et al.* 2007). Furthermore, extinct taxa that morphologically resemble extant studied organisms but occupied different niches could exist cryptically in the fossil record (Varela *et al.* 2009).

Finally, local versus regional shifts in species distributions can be difficult to distinguish from the fossil record. Small patches of suitable habitat available locally to the organism may not be recovered in coarse transferal models. In such cases, the fossil record documents presence while the transferal models predict absence of the organisms within a particular area. This can also be a problem when evaluation of the transferal model is based on species absence from the fossil record. For example, while a species can be locally extinct within the small area from which the fossil records originated (e.g. within the home range of the raptor that deposited pellets with prey remains), it could persist regionally within the resolution of the transferal model. In such cases, the fossil record seemingly does not support the transferal model.

#### Palaeo-environmental reconstructions

The PMIP experiments of simulated past climates have been evaluated using palaeo-environmental reconstructions. Several global palaeo-environmental datasets have been used for PMIP experiment evaluation (Farrera *et al.* 1999; Harrison 2000; Kohfeld & Harrison 2000; Prentice & Jolly 2000; Prentice & Webb 1998; Qin *et al.* 1998). For western North America, BIOME 6000 (Thompson & Anderson 2000) represents a pollen and plant macrofossil dataset, with the latter primarily derived from numerous packrat middens (*i.e.* woodrats, rodent species within the genus *Neotoma*). Interestingly, the

pollen and macrofossil data do not necessarily provide the same information. The fossil pollen generally represents a larger, regional picture of the palaeo-vegetation but with less detail (lower species richness), while the packrat middens provide more detailed information (higher species richness) but only locally, within the home range of a woodrat (Mehring & Wigand 1990). Consequently, attention must be paid when comparing palaeo-vegetation from two areas reconstructed from these different data sources. Packrat midden data should also be interpreted with caution because different species of woodrats (which are difficult to identify by their middens) have different preferences for collection of plant material and consequently their midden contents can be different even within the same environment. This problem may be exacerbated through time if one woodrat species replaces another within the same midden, which could result in an incorrect interpretation of environmental change when none occurred (Dial & Czaplewski 1990).

### Concluding remarks

In this paper, I identified several general challenges, both algorithmic and conceptual, that can complicate the use of transferal modeling. Our major goal was to stimulate more objective evaluations of transferal models with better understandings of the assumptions and uncertainties. I hope that our experiments and discussion will motivate researchers to further explore these and other challenges associated with transferal modeling, as our list was certainly not intended to be complete nor have I provided complete solutions to the challenges I describe. While some of the challenges

are substantial, and should raise concerns regarding the accuracy of this approach, I believe that transferal modeling offers strong potential to provide biogeographers with an important approach for increasing my understanding of evolutionary and biogeographic processes.



**Table 4.1.** Range of values of 19 bioclimatic variables for the OK and LGM (CCSM and Miroc) datasets, when (A) both the OK and LGM variables are clipped to the geographic extent of the western North America, and (B) the OK dataset represents the entire world while the LGM variables are clipped to the extent of western North America. Highlighted are LGM values that are outside the OK range (light grey < 10%, dark grey ≥ 10%).

| A     | OK (western) |       | CCSM (western) |       | MIROC (western) |       |
|-------|--------------|-------|----------------|-------|-----------------|-------|
|       | Low          | High  | Low            | High  | Low             | High  |
| Bio01 | -71          | 294   | -314           | 277   | -388            | 282   |
| Bio02 | 32           | 213   | -31            | 360   | 32              | 219   |
| Bio03 | 19           | 86    | -9             | 86    | 21              | 34    |
| Bio04 | 449          | 14199 | 452            | 14195 | 436             | 14200 |
| Bio05 | 60           | 456   | -108           | 422   | -128            | 389   |
| Bio06 | -295         | 215   | -557           | 200   | -667            | 205   |
| Bio07 | 118          | 522   | 99             | 628   | 99              | 549   |
| Bio08 | -135         | 332   | -252           | 299   | -341            | 293   |
| Bio09 | -196         | 328   | -481           | 273   | -553            | 278   |
| Bio10 | 5            | 360   | -143           | 299   | -216            | 307   |
| Bio11 | -221         | 283   | -507           | 267   | -581            | 272   |
| Bio12 | 35           | 4524  | 13             | 5990  | 59              | 3750  |
| Bio13 | 7            | 881   | 4              | 1151  | 9               | 1059  |
| Bio14 | 0            | 112   | 0              | 167   | 0               | 172   |
| Bio15 | 5            | 138   | 5              | 141   | 5               | 138   |
| Bio16 | 18           | 2378  | 9              | 3150  | 24              | 2401  |
| Bio17 | 0            | 366   | 0              | 541   | 0               | 547   |

|          |            |       |                |       |                 |       |
|----------|------------|-------|----------------|-------|-----------------|-------|
| Bio18    | 1          | 1608  | 0              | 1854  | 5               | 2359  |
| Bio19    | 1          | 1455  | 3              | 1350  | 0               | 1901  |
| <b>B</b> | OK (world) |       | CCSM (western) |       | MIROC (western) |       |
|          | Low        | High  | Low            | High  | Low             | High  |
| Bio01    | -278       | 319   | -314           | 277   | -388            | 282   |
| Bio02    | 9          | 213   | -31            | 360   | 32              | 219   |
| Bio03    | 8          | 96    | -9             | 86    | 21              | 34    |
| Bio04    | 64         | 22704 | 452            | 14195 | 436             | 14200 |
| Bio05    | -86        | 489   | -108           | 422   | -128            | 389   |
| Bio06    | -559       | 258   | -557           | 200   | -667            | 205   |
| Bio07    | 53         | 725   | 99             | 628   | 99              | 549   |
| Bio08    | -278       | 376   | -252           | 299   | -341            | 293   |
| Bio09    | -501       | 365   | -481           | 273   | -553            | 278   |
| Bio10    | -127       | 382   | -143           | 299   | -216            | 307   |
| Bio11    | -506       | 289   | -507           | 267   | -581            | 272   |
| Bio12    | 0          | 2115  | 13             | 5990  | 59              | 3750  |
| Bio13    | 0          | 2437  | 4              | 1151  | 9               | 1059  |
| Bio14    | 0          | 697   | 0              | 167   | 0               | 172   |
| Bio15    | 0          | 265   | 5              | 141   | 5               | 138   |
| Bio16    | 0          | 6586  | 9              | 3150  | 24              | 2401  |
| Bio17    | 0          | 2319  | 0              | 541   | 0               | 547   |
| Bio18    | 0          | 5040  | 0              | 1854  | 5               | 2359  |
| Bio19    | 0          | 4580  | 3              | 1350  | 0               | 1901  |

**Figure 4.1.** Difference between the two climatic models for the western region of North America: The left column represents Bio 12 (Annual Precipitation) for (A) the CCSM model, (B) MIROC model, and (C) the absolute difference between the two models. The values are divided using the natural (Jenks) breaks into five classes that maximize differences among classes. The right column represents the ENMs for the Merriam's kangaroo rat (*Dipodomys merriami*) using 19 bioclimatic variables projected on (C) the CCSM and (E) MIROC climatic reconstructions of the LGM, and (F) the absolute difference of logistic probabilities between these two models. Shading represents different categories of logistic probabilities.

**Figure 4.2.** ENMs for the Great Basin Pocket Mouse (*Perognathus parvus*) created using 19 bioclimatic variables and elevation for (A) current and (B) LGM conditions. ENMs with elevation excluded are shown for (C) current and (D) LGM conditions. While the models calibrated on current variables do not differ significantly whether the elevation was included or not, the LGM model with (B) elevation included has less than 1/3 of suitable habitat than (D) the model with elevation omitted.

**Figure 4.3.** Histograms of elevation for the areas with logistic probabilities higher than 0.2 extracted from the ENMs of *Perognathus parvus*. The current model was created using 19 bioclimatic variables and elevation (top). The LGM models were created using all bioclimatic variables (MIROC) but with elevation included or excluded (middle and bottom, respectively). While both LGM models predict extinction of the species at

higher elevations (blue rectangles), only the LGM model that excludes elevation predicts that the species shifted to lower elevations (red rectangle).

**Figure 4.4.** Schematic picture showing different approaches when projecting a model outside the calibration range (extrapolation). The curve represents a hypothetical unimodal response of organism to an environmental variable. In each picture, line A represents interpolation of values using a regression method; line B represents a method utilized by Maxent where the values outside the training range are assigned the probability value of the last (highest or lowest) value of the training range; and line C represents a method where all values of habitat suitability outside the calibration range are set to zero. Although both figures represent the same response of the organism, the extrapolation differs based on what part of the response curve is available during calibration (see text for details).

**Figure 4.5.** The clamping values for ENMs of the striped whipsnake (*Masticophis taeniatus*) when calibration and projection was performed on 19 bioclimatic variables (A) clipped to the western North America and (B) when the calibration of the model was performed using bioclimatic variables representing the entire world and then projected on the truncated dataset of the western North America. The clamping values range from 0 to 1 because they represent change in logistic probabilities caused by clamping (see text for details).

**Figure 4.6.** A set of response curves of 19 bioclimatic variables for the common side-blotched lizard (*Uta stansburiana*) produced by Maxent. The X-axis represents a range of values of each variable for the OK climatic conditions, and the Y-axis represents the logistic probability value interpreted as occurrence probability or habitat suitability.

**Figure 4.7.** (A) ENM for the common side-blotched lizard (*Uta stansburiana*) for the LGM when all 19 bioclimatic variables are included, and (B) the associated clamping values. (C) The LGM model derived with variable Bio 2 (mean diurnal temperature range) excluded, and (D) the associated clamping values.

**Figure 4.8.** Representation of a niche shift in a hypothetical species from time T1 to time T2. The shift in T2A was due to an evolutionary adaptation that resulted in an expansion of the species fundamental niche, while in T2B, the fundamental niche remained conserved while the realized niche shifted (because of changes in biotic factors, plasticity of the organism that allowed it to persist in place, or because of the appearance of novel environmental conditions that were not available at time T1).


**Figure 4.9.** Comparison of principal components scores derived from bioclimatic niches occupied by four species of rodents at OK (empty symbols) and during the LGM (red and green colored symbols). The axes represent the first two principal components of the 19 bioclimatic variables after data reduction using principal component analysis. The occurrence records representing the OK climatic conditions were downloaded from

MaNIS while the occurrence records representing the LGM were obtained from the Faunmap database, constrained to the full glacial (ca. 14,500-20,500 years B.P.).

**Figure 4.10.** Ecological niche models for four species of rodents using the LGM reconstructions of CCSM (left column) and MIROC (right column). The red dots represent fossil records dated to the full glacial (ca. 14,500-20,500 years B.P.) obtained from the Faunmap database. The shades represent different categories of logistic probabilities.

**Figure 4.11.** Two scenarios of a hypothetical species history leading to the same genetic signal. In scenario A, the species persists only in Area 1 during an early time (T1), generating high genetic diversity. When habitat becomes suitable at Area 2 during a subsequent time (T2), a small subset of the genotypes from Area 1 colonizes Area 2. In scenario B, both Areas 1 and 2 are suitable during T1. During T2, genotypes from Area 2 go extinct while a small subset of the genotypes from Area 1 colonizes Area 2. In both cases, the genetic structure identified during current time (T3) is identical.

Figure 4.1











Figure 4.2








Figure 4.3



**Figure 4.4**




Figure 4.5





Figure 4.6




Figure 4.7




Figure 4.8






Figure 4.9





Figure 4.10








Figure 4.11



## Bibliography

- Ackerly D (2003a) Canopy gaps to climate change - extreme events, ecology and evolution. *New Phytologist* **160**, 2-4.
- Ackerly DD (2003b) Community assembly, niche conservatism, and adaptive evolution in changing environments. *International Journal of Plant Sciences* **164**, S165-S184.
- Alexander LF, Riddle BR (2005) Phylogenetics of the new world rodent family heteromyidae. *Journal of Mammalogy* **86**, 366-379.
- Araujo MB, Guisan A (2006) Five (or so) challenges for species distribution modelling. *Journal of Biogeography* **33**, 1677-1688.
- Araujo MB, New M (2007) Ensemble forecasting of species distributions. *Trends in Ecology & Evolution* **22**, 42-47.
- Avice JC (2000) *Phylogeography: The History and Formation of Species* Harvard University Press, Cambridge, MA.
- Braconnot P, Otto-Bliesner B, Harrison S, Joussaume S, Peterchmitt JY, Abe-Ouchi A, Crucifix M, Driesschaert E, Fichet T, Hewitt CD, Kageyama M, Kitoh A, Laine A, Loutre MF, Marti O, Merkel U, Ramstein G, Valdes P, Weber SL, Yu Y, Zhao Y (2007) Results of PMIP2 coupled simulations of the Mid-Holocene and Last Glacial Maximum - Part 1: experiments and large-scale features. *Climate of the Past* **3**, 261-277.
- Broennimann O, Treier UA, Muller-Scharer H, Thuiller W, Peterson AT, Guisan A (2007) Evidence of climatic niche shift during biological invasion. *Ecology Letters* **10**, 701-709.
- Bromham L, Penny D (2003) The modern molecular clock. *Nature Reviews Genetics* **4**, 216-224.
- Calvignac S, Hughes S, Tougaard C, Michaux J, Thevenot M, Philippe M, Hamdine W, Hanni C (2008) Ancient DNA evidence for the loss of a highly divergent brown bear clade during historical times. *Molecular Ecology* **17**, 1962-1970.
- Carnaval AC, Moritz C (2008) Historical climate modelling predicts patterns of current biodiversity in the Brazilian Atlantic forest. *Journal of Biogeography* **35**, 1187-1201.
- Carstens BC, Brunsfeld SJ, Demboski JR, Good JM, Sullivan J (2005) Investigating the evolutionary history of the Pacific Northwest mesic forest ecosystem: Hypothesis

- testing within a comparative phylogeographic framework. *Evolution* **59**, 1639-1652.
- Carstens BC, Richards CL (2007) Integrating coalescent and ecological niche modeling in comparative phylogeography. *Evolution* **61**, 1439-1454.
- Colautti RI, Eckert CG, Barrett SCH (2010) Evolutionary constraints on adaptive evolution during range expansion in an invasive plant. *Proceedings of the Royal Society B-Biological Sciences* **277**, 1799-1806.
- Davis MB, Shaw RG (2001) Range shifts and adaptive responses to Quaternary climate change. *Science* **292**, 673-679.
- Davis MB, Shaw RG, Etterson JR (2005) Evolutionary responses to changing climate. *Ecology* **86**, 1704-1714.
- Dial KP, Czaplewski NJ (1990) Do Woodrat Middens Accurately Represent the Animal's Environments and Diets? The Woodhouse Mesa Study. In: *Packrat Middens: The last 40,000 Years of Biotic Change* (eds. Betancourt JL, Van Devender TR, Martin PS), p. 468. The University of Arizona Press, Tucson.
- Edwards SV, Beerli P (2000) Perspective: Gene divergence, population divergence, and the variance in coalescence time in phylogeographic studies. *Evolution* **54**, 1839-1854.
- Elith J, Graham CH, Anderson RP, Dudik M, Ferrier S, Guisan A, Hijmans RJ, Huettmann F, Leathwick JR, Lehmann A, Li J, Lohmann LG, Loiselle BA, Manion G, Moritz C, Nakamura M, Nakazawa Y, Overton JM, Peterson AT, Phillips SJ, Richardson K, Scachetti-Pereira R, Schapire RE, Soberon J, Williams S, Wisz MS, Zimmermann NE (2006) Novel methods improve prediction of species' distributions from occurrence data. *Ecography* **29**, 129-151.
- Excoffier L, Foll M, Petit RJ (2009) Genetic consequences of range expansions. *Annual Review of Ecology Evolution and Systematics* **40**, 481-501.
- Farrera I, Harrison SP, Prentice IC, Ramstein G, Guiot J, Bartlein PJ, Bonnefille R, Bush M, Cramer W, von Grafenstein U, Holmgren K, Hooghiemstra H, Hope G, Jolly D, Lauritzen SE, Ono Y, Pinot S, Stute M, Yu G (1999) Tropical climates at the Last Glacial Maximum: a new synthesis of terrestrial palaeoclimate data. I. Vegetation, lake levels and geochemistry. *Climate Dynamics* **15**, 823-856.
- Fitzpatrick MC, Weltzin JF, Sanders NJ, Dunn RR (2007) The biogeography of prediction error: why does the introduced range of the fire ant over-predict its native range? *Global Ecology and Biogeography* **16**, 24-33.

- Gibbs JP, Breisch AR (2001) Climate warming and calling phenology of frogs near Ithaca, New York, 1900-1999. *Conservation Biology* **15**, 1175-1178.
- Graham CH, Moritz C, Williams SE (2006) Habitat history improves prediction of biodiversity in rainforest fauna. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 632-636.
- Graham RW, Lundelius EL, Jr. , Group FW (1994) FAUNMAP: a database documenting Late Quaternary distributions of mammal species in the United States. *Illinois State Museum Scientific Papers* **XXV**, 1-690.
- Grayson DK (2000) Mammalian responses to Middle Holocene climatic change in the Great Basin of the western United States. *Journal of Biogeography* **27**, 181-192.
- Grootes PM, Stuiver M (1997) Oxygen 18/16 variability in Greenland snow and ice with 10(-3)- to 10(5)-year time resolution. *Journal of Geophysical Research-Oceans* **102**, 26455-26470.
- Guisan A, Zimmermann NE (2000) Predictive habitat distribution models in ecology. *Ecological Modelling* **135**, 147-186.
- Hafner JC, Light JE, Hafner DJ, Hafner MS, Reddington E, Rogers DS, Riddle BR (2007) Basal clades and molecular systematics of heteromyid rodents. *Journal of Mammalogy* **88**, 1129-1145.
- Harrison SP (2000) Palaeoenvironmental data sets and model evaluation in PMIP, 9-25.
- Hasumi H, Emori S (2004) K-1 Coupled Model (MIROC) Description. Center for Climate System Research, University of Tokyo, Tokyo, Japan.
- Hewitt G (2000) The genetic legacy of the Quaternary ice ages. *Nature* **405**, 907-913.
- Hewitt GM (1996) Some genetic consequences of ice ages, and their role in divergence and speciation. *Biological Journal of the Linnean Society* **58**, 247-276.
- Hewitt GM (2004) Genetic consequences of climatic oscillations in the Quaternary. *Philosophical Transactions of the Royal Society of London Series B-Biological Sciences* **359**, 183-195.
- Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005a) Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* **25**, 1965-1978.

- Hijmans RJ, Guarino L, Bussink C, Barrantes I, Rojas E (2005b) Diva-GIS version 5.2. Manual.
- Ho SYW, Phillips MJ, Cooper A, Drummond AJ (2005) Time dependency of molecular rate estimates and systematic overestimation of recent divergence times. *Molecular Biology and Evolution* **22**, 1561-1568.
- Hoffmann MH (2005) Evolution of the realized climatic niche in the genus *Arabidopsis* (Brassicaceae). *Evolution* **59**, 1425-1436.
- Hunter KL, Betancourt JL, Riddle BR, Van Devender TR, Cole KL, Spaulding WG (2001) Ploidy race distributions since the Last Glacial Maximum in the North American desert shrub, *Larrea tridentata*. *Global Ecology and Biogeography* **10**, 521-533.
- Huppopp O, Huppopp K (2003) North Atlantic Oscillation and timing of spring migration in birds. *Proceedings of the Royal Society of London Series B-Biological Sciences* **270**, 233-240.
- Irwin DE (2002) Phylogeographic breaks without geographic barriers to gene flow. *Evolution* **56**, 2383-2394.
- Jackson ST, Overpeck JT (2000) Responses of plant populations and communities to environmental changes of the late Quaternary. *Paleobiology* **26**, 194-220.
- Jansson R (2003) Global patterns in endemism explained by past climatic change. *Proceedings of the Royal Society of London Series B-Biological Sciences* **270**, 583-590.
- Jeschke JM, Strayer DL (2008) Usefulness of bioclimatic models for studying climate change and invasive species. In: *Year in Ecology and Conservation Biology 2008*, pp. 1-24.
- Jezkova T, Jaeger JR, Marshall ZL, Riddle BR (2009) Pleistocene impacts on the phylogeography of the desert pocket mouse (*Chaetodipus penicillatus*). *Journal of Mammalogy* **90**, 306-320.
- Knowles LL (2001) Did the Pleistocene glaciations promote divergence? Tests of explicit refugial models in montane grasshoppers. *Molecular Ecology* **10**, 691-701.
- Knowles LL, Maddison WP (2002) Statistical phylogeography. *Molecular Ecology* **11**, 2623-2635.



- Kohfeld KE, Harrison SP (2000) How well can I simulate past climates? Evaluating the models using global palaeoenvironmental datasets. *Quaternary Science Reviews* **19**, 321-346.
- Losos JB, Glor RE, Kolbe JJ, Nicholson K (2006) Adaptation, speciation, and convergence: A hierarchical analysis of adaptive radiation in Caribbean *Anolis* lizards. *Annals of the Missouri Botanical Garden* **93**, 24-33.
- Maddison WP, Knowles LL (2006) Inferring phylogeny despite incomplete lineage sorting. *Systematic Biology* **55**, 21-30.
- Martinez-Meyer E, Townsend Peterson A, Hargrove WW (2004) Ecological niches as stable distributional constraints on mammal species, with implications for Pleistocene extinctions and climate change projections for biodiversity. *Global Ecology and Biogeography* **13**, 305-314.
- Mehring PJJ, Wigand PE (1990) Comparison of Late Holocene Environments from Woodrat Middens and Pollen: Diamond Craters, Oregon. In: *Packrat Middens: The Last 40,000 Years of Biotic Change* (eds. Betancourt JL, Van Devender TR, Martin PS), p. 468. University of Arizona Press, Tucson.
- Moritz C, Patton JL, Conroy CJ, Parra JL, White GC, Beissinger SR (2008) Impact of a century of climate change on small-mammal communities in Yosemite National Park, USA. *Science* **322**, 261-264.
- Neigel JE, Avise JC (1993) Application of Random-walk Model to Geographic Distributions of Animal Mitochondrial-DNA Variation. *Genetics* **135**, 1209-1220.
- Nogues-Bravo D (2009) Predicting the past distribution of species climatic niches. *Global Ecology and Biogeography* **18**, 521-531.
- Nogues-Bravo D, Rodriguez J, Hortal J, Batra P, Araujo MB (2008) Climate change, humans, and the extinction of the woolly mammoth. *Plos Biology* **6**, 685-692.
- Otto-Bliesner BL, Brady EC, Clauzet G, Tomas R, Levis S, Kothavala Z (2006) Last Glacial Maximum and Holocene climate in CCSM3. *Journal of Climate* **19**, 2526-2544.
- Pearman PB, Guisan A, Broennimann O, Randin CF (2008) Niche dynamics in space and time. *Trends in Ecology & Evolution* **23**, 149-158.
- Pearson RG, Dawson TP, Berry PM, Harrison PA (2002) SPECIES: A Spatial Evaluation of Climate Impact on the Envelope of Species. *Ecological Modelling* **154**, 289-300.

- Pearson RG, Thuiller W, Araujo MB, Martinez-Meyer E, Brotons L, McClean C, Miles L, Segurado P, Dawson TP, Lees DC (2006) Model-based uncertainty in species range prediction. *Journal of Biogeography* **33**, 1704-1711.
- Peterson AT, Martinez-Meyer E, Gonzalez-Salazar C (2004) Reconstructing the Pleistocene geography of the *Aphelocoma* jays (Corvidae). *Diversity and Distributions* **10**, 237-246.
- Peterson AT, Nyari AS (2008) Ecological niche conservatism and Pleistocene refugia in the thrush-like mourner, *Schiffornis* sp., in the neotropics. *Evolution* **62**, 173-183.
- Peterson AT, Papes M, Eaton M (2007) Transferability and model evaluation in ecological niche modeling: a comparison of GARP and Maxent. *Ecography* **30**, 550-560.
- Peterson AT, Papes M, Kluza DA (2003) Predicting the potential invasive distributions of four alien plant species in North America. *Weed Science* **51**, 863-868.
- Peterson AT, Soberon J (2005) Interpretation of models of fundamental ecological niches and species' distributional areas. *Biodiversity Informatics* **2**, 1-10.
- Phillips SJ (2008) Transferability, sample selection bias and background data in presence-only modelling: a response to Peterson et al. (2007). *Ecography*.
- Phillips SJ, Anderson RP, Schapire RE (2006) Maximum entropy modeling of species geographic distributions. *Ecological Modelling* **190**, 231-259.
- Phillips SJ, Dudik M (2008) Modeling of species distributions with Maxent: new extensions and a comprehensive evaluation. *Ecography* **31**, 161-175.
- Phillips SJ, Dudik M, Schapire RE (2004) A maximum entropy approach to species distribution modeling, 655-662.
- Post E, Langvatn R, Forchhammer MC, Stenseth N (1999) Environmental variation shapes sexual dimorphism in red deer. *Proceedings of the National Academy of Sciences of the United States of America* **96**, 4467-4471.
- Prentice IC, Jolly D (2000) Mid-Holocene and glacial-maximum vegetation geography of the northern continents and Africa. *Journal of Biogeography* **27**, 507-519.
- Prentice IC, Webb T (1998) BIOME 6000: reconstructing global mid-Holocene vegetation patterns from palaeoecological records. *Journal of Biogeography* **25**, 997-1005.

- Qin BQ, Harrison SP, Kutzbach JE (1998) Evaluation of modelled regional water balance using lake status data: A comparison of 6 ka simulations with the NCAR CCM. *Quaternary Science Reviews* **17**, 535-548.
- Randin CF, Dirnbock T, Dullinger S, Zimmermann NE, Zappa M, Guisan A (2006) Are niche-based species distribution models transferable in space? *Journal of Biogeography* **33**, 1689-1703.
- Rodder D, Lotters S (2009) Niche shift versus niche conservatism? Climatic characteristics of the native and invasive ranges of the Mediterranean house gecko (*Hemidactylus turcicus*). *Global Ecology and Biogeography* **18**, 674-687.
- Rodriguez-Robles JA, Jezkova T, Leal M (2010) Climatic stability and genetic divergence in the tropical insular lizard *Anolis krugi*, the Puerto Rican 'Lagartijo Jardinero de la Montana'. *Molecular Ecology* **19**, 1860-1876.
- Rowe R, Finarelli J, Rickart E (2010) Range dynamics of small mammals along an elevational gradient over an 80-year interval. *Global Change Biology* **early view**, 1-14.
- Smith FA, Betancourt JL (2003) The effect of Holocene temperature fluctuations on the evolution and ecology of *Neotoma* (woodrats) in Idaho and northwestern Utah. *Quaternary Research* **59**, 160-171.
- Spaulding AW (1990) Vegetational and climatic development of the Mojave Desert: The last glacial maximum to the Present. In: *Packrat middens: The last 40,000 years of biotic change* (eds. Betancourt JL, Van Devender TR, Martin PS), pp. 200-239. The University of Arizona Press, Tucson.
- Spradling TA, Hafner MS, Demastes JW (2001) Differences in rate of cytochrome-b evolution among species of rodents. *Journal of Mammalogy* **82**, 65-80.
- Stockwell D, Peters D (1999) The GARP modelling system: problems and solutions to automated spatial prediction. *International Journal of Geographical Information Science* **13**, 143-158.
- Stockwell DRB (2006) Improving ecological niche models by data mining large environmental datasets for surrogate models. *Ecological Modelling* **192**, 188-196.
- Thompson RS, Anderson KH (2000) Biomes of western North America at 18,000, 6000 and 0 C-14 yr BP reconstructed from pollen and packrat midden data. *Journal of Biogeography* **27**, 555-584.

- Thompson RS, Whitlock C, Bartlein PJ, Harrison SP, Spaulding WG (1993) Climatic changes in the western United States since 18,000 yr B.P. . In: *Global climates since the last glacial maximum* (eds. Wright Jr. HE, Kutzbach JE, Webb III T, *et al.*), pp. 468-513. University of Minnesota Press, Minneapolis, MN.
- Thuiller W, Brotons L, Araujo MB, Lavorel S (2004) Effects of restricting environmental range of data to project current and future species distributions. *Ecography* **27**, 165-172.
- Tracy RL, Walsberg GE (2002) Kangaroo rats revisited: re-evaluating a classic case of desert survival. *Oecologia* **133**, 449-457.
- Urban MC, Phillips BL, Skelly DK, Shine R (2007) The cane toad's (*Chaunus* [*Bufo*] *marinus*) increasing ability to invade Australia is revealed by a dynamically updated range model. *Proceedings of the Royal Society B-Biological Sciences* **274**, 1413-1419.
- Varela S, Rodríguez J, Lobo JM (2009) Is current climatic equilibrium a guarantee for the transferability of distribution model predictions? A case study of the spotted hyena. *Journal of Biogeography* **36**, 1645-1655.
- Waltari E, Guralnick RP (2009) Ecological niche modelling of montane mammals in the Great Basin, North America: examining past and present connectivity of species across basins and ranges. *Journal of Biogeography* **36**, 148-161.
- Waltari E, Hijmans RJ, Peterson AT, Nyari AS, Perkins SL, Guralnick RP (2007) Locating Pleistocene refugia: comparing phylogeographic and ecological niche model predictions. *PLoS ONE* **2**, e563.
- Warren DL, Glor RE, Turelli M (2008) Environmental niche equivalency versus conservatism: quantitative approaches to niche evolution. *Evolution* **62**, 2868-2883.
- Wiens JJ, Graham CH (2005) Niche conservatism: integrating evolution, ecology, and conservation biology. *Annual Review of Ecology Evolution and Systematics* **36**, 519-539.

## APPENDIX 1

### SAMPLES OF *DIPODOMYS MICROPS*

Samples of *Dipodomys microps* sorted by locality identification number. Ear-clips are identified by a tissue number (LVT – Las Vegas Tissues), vouchers are identified by a tissue and voucher number (MVZ – Museum of Vertebrate Zoology, TTU – Texas Tech University, NMMNH – New Mexico Museum of Natural History). Latitude and longitude are in decimal degrees and WGS 84 datum.

Arizona: Coconino County: **1** – House Rock Valley, 36.4940, -111.9623 [LVT-9375, LVT-9377, LVT-9380, LVT-9382, LVT-9383, LVT-9384 (NMMNH 5612)]; 36.6294, -111.9747 [LVT-9361, LVT-9363, LVT-9365, LVT-9368]; 36.7256, -111.7583 [LVT-9370]; Mohave County: **2** – 10 mi N Wolf's Hole, 36.8847, -113.5625 [LVT-10399, LVT-10400, LVT-10401, LVT-10402, LVT-10403, LVT-10404, LVT-10405, LVT-10406, LVT-10407]; 36.8939, -113.5615 [LVT-10392, LVT-10393, LVT-10394, LVT-10395, LVT-10396, LVT-10397]; California: Inyo County: **3** – Coso Junction, 36.0517, -117.9370 [LVT-8699 (NMMNH 5601)]; Haiwee, 36.1532, -117.9781 [LVT-8688, LVT-8689, LVT-8692, LVT-8695]; Olancho, 36.2516, -117.9867 [LVT-8683, LVT-8684 (NMMNH 5599)]; **4** – Owens Valley, 36.5364, -117.9256 [LVT-7795 (NMMNH 5519)]; **5** – Saline Valley, 36.8388, -117.9139 [LVT-9795]; **6** – Deep Spring Valley, 37.3038, -118.0562, [LVT-8133 (NMMNH 5580), LVT-8134, LVT-8135, LVT-8136]; Kern County: **7** – Koehl Lake, 35.2890, -117.8800 [LVT-8700, LVT-8700, LVT-8701, LVT-8702, LVT-8703, LVT-8704, LVT-8705, LVT-8706, LVT-8707,

LVT-8708, LVT-8709, LVT-8710, LVT-8712, LVT-8713]; Modoc County: **8** – 8 mi S Eagleville, 41.1621, -119.9928 [LVT-8900, LVT-8901, LVT-8903, LVT-8905]; 5 mi E, 3 mi N Eagleville, 41.3649, -120.0007 [LVT-8911, LVT-8914, LVT-8915, LVT-8917, LVT-8918, LVT-8919, LVT-8920, LVT-8921, LVT-8922]; Riverside County: **9** – Stubby Spring, 33.9863, -116.2305 [MVZ-114058, MVZ-114059, MVZ-148434, MVZ-148435, MVZ-148436, MVZ-149483, MVZ-149484, MVZ-149569]; San Bernardino County: **10** – 12 mi NNE Boron, 35.1493, -117.5718 [LVT-2058]; **11** – Yucca Valley, 34.1450, -116.4684 [MVZ-159304, MVZ-159305]; **12** – Superior Lake, 35.2259, -117.0300 [LVT 10465, LVT 10466]; **13** – Clark Mountain, 35.5855, -115.6357 [LVT-10432, LVT-10433, LVT-10434, LVT-10435, LVT-10436, LVT-10437, LVT-10438, LVT-10439, LVT-10440, LVT-10441, LVT-10442, LVT-10443, LVT-10444, LVT-10445]; Idaho: Owyhee County: **14** – Bruneau Canyon, 42.7629, -115.7564 [LVT-10368, LVT-10369, LVT-10370, LVT-10371]; **15** – 5 mi W Murphy, 43.1902, -116.6445 [LVT-8969, LVT-8970, LVT-8972, LVT-8983]; 2 mi W Murphy, 43.2012, -116.5861 [LVT-8976, LVT-8978, LVT-8979, LVT-8980, LVT-8981, LVT-8985]; Nevada: Churchill County: **16** – Fallon (Sand Creek Road), 39.3765, -118.8837 [LVT-9587]; 10 mi W Fallon, 39.4988, -118.9875 [LVT-9573]; **17** – Hot Springs Mountains, 39.7581, -118.8735 [LVT-10203, LVT-10204, LVT-10205]; Clark County: Cottonwood Valley, 36.0015, -115.4496; **18** – Cottonwood Valley, 36.0015, -115.4496 [LVT-10684, LVT-10685]; **19** – Kyle Canyon, 36.2643, -115.4788 [LVT-10446, LVT-10448, LVT-10449, LVT-10450, LVT-10451, LVT-10452, LVT-10453, LVT-10454, LVT-10455, LVT-10456, LVT-10457, LVT-10458, LVT-10459, LVT-10460]; Elko County: **20** – Montello, 41.2886, -114.1579 [LVT-9659, LVT-9660, LVT-9661, LVT-9662]; Tecoma, 41.3196, -114.0804 [LVT-

9653, LVT-9655, LVT-9656, LVT-9657, LVT-9658]; Esmeralda County: **21** – Fish Lake Valley, 37.7805, -118.0980 [LVT-8128 (NMMNH 5577)]; **22** – 10 mi S Tonopah, 37.9120, -117.2235 [LVT-10161, LVT-10162, LVT-10164, LVT-10165, LVT-10166, LVT-10167]; **23** – White River Valley, 38.3865, -115.1460 [LVT 08232 (NMMNH 5595), LVT 08234]; 38.4084, -115.1199 [LVT 08231 (NMMNH 5594)]; Eureka County: **24** – 5 mi SSW Beowawe, 40.5203, -116.5138 [LVT-9640, LVT-9641, LVT-9642]; 3 mi SSW Beowawe, 40.5614, -116.4881 [LVT-9646]; Humboldt County: **25** – 10 mi S Battle Mountain, 40.4751, -117.0621 [LVT-8986, LVT-8987, LVT-8988, LVT-8989, LVT-8990]; **26** – ~5 mi W Sulphur Landing, 40.8432, -118.7577 [LVT-9610, LVT-9611, LVT-9612, LVT-9613, LVT-9614, LVT-9615, LVT-9616]; **27** – 4 mi E Golconda, 40.9586, -117.4243 [LVT-9635, LVT-9636]; **28** – 30 mi S Denio, 41.6474, -118.4404 [LVT-9635, LVT-9636]; Lander County: **29** – Monitor Valley, 39.2288, -116.6990 [LVT-10175, LVT-10176, LVT-10177, LVT-10178, LVT-10179, LVT-10180]; **30** – Big Smoky Valley, 39.3893, -116.9322 (LVT-10193); Lincoln County: **31** – Coyote Spring, 37.1481, -114.7228 [LVT-9664, LVT-9665]; 7 mi E Coyote Spring, 37.1481, -114.7228 [LVT-9785]; **32** – Pahrnagat NWR, 37.2044, -115.0662 [TTU-161809]; 37.2343, -115.1019 [TTU-161826, TTU-161827, TTU-161828]; 37.2343, -115.1010 [TTU-148048]; 37.2415, -115.1144 [TTU-161832, TTU-161833]; 37.2829, -115.1272 [TTU-161836]; **33** – Kane Spring Valley, 37.2993, -114.5806 [LVT-10536, LVT-10537, LVT-10538, LVT-10539, LVT-10540, LVT-10541, LVT-10542, LVT-10543, LVT-10544, LVT-10545, LVT-10546, LVT-10547, LVT-10548, LVT-10549, LVT-10550, LVT-10551, LVT-10552, LVT-10553]; **34** – Tickaboo Valley, 37.3195, -115.4383 [LVT-8163]; 37.3696, -115.4697 [LVT-9452]; 37.3969, -115.4625 [LVT-9458, LVT-9459]; **35** – Delmar

Valley, 37.5211, -114.8689 [LVT-7754]; 37.5572, -114.8600 [LVT-7771]; **36** – 6 mi N, 31 mi W Hiko, 37.6200, -115.8303 [LVT 05140]; Rachel, 37.7059, -115.8135 [LVT-8158, LVT-8159, LVT-8160, LVT-8161, LVT-8162]; **37** – Dry Lake Valley, 37.7119, -114.7976 [LVT-7757]; 37.7130, -114.8012 [LVT-7764]; 37.7784, -114.8301 [LVT-7773]; **38** – ~6 mi NE Tempiute, 37.7393, -115.5759 [LVT-1570 (NMMNH-5909)]; **39** – Dry Lake Valley, 37.9428, -114.8320 [LVT-7775]; 38.0694, -114.7744 [LVT-7780 (NMMNH-5511)]; **40** – Cave Valley, 38.3672, -114.8243 [LVT-8060 (NMMNH-5566)]; 38.3773, -114.8428 [LVT-8064 (NMMNH-5570)]; 38.3992, -114.8171 [LVT-8061 (NMMNH-5567)]; **41** – Lake Valley, 38.4299, -114.5834 [LVT-7822 (NMMNH-5533)]; 38.4515, -114.6098 [LVT-7821 (NMMNH-5532), LVT-7835 (NMMNH-5539)]; Mineral County: **42** – Marietta, Teels Marsh, 38.2294, -118.3136 [LVT-8498]; **43** – Tonapah Junction, 38.2670, -118.1085 [LVT-8505, LVT-8506, LVT-8507]; **44** – Smith Valley, 38.8371, -119.3057 [LVT-9571]; Nye County: **45** – Ash Meadows NWR, 36.3773, -116.2977 [TTU-161787, TTU-161788, TTU-161789]; 36.4037, -116.2754 [TTU-150485]; 36.4267, -116.3018 [TTU-161761]; 36.4708, -116.3235 [TTU-148008]; **46** – Beatty, 36.9086, -116.7583 [LVT-9201, LVT-9202, LVT-9203, LVT-9204, LVT-9205, LVT-9206, LVT-9207, LVT-9208]; 9 mi N Beatty, 36.9845, -116.7230 [LVT-4935 (NMMNH-5968)]; **47** – Nevada Test Site, Area 9, 37.1251, -116.0212 [LVT-10518, LVT-10519, LVT-105120]; Nevada Test Site, Area 10, 37.1561, -115.9720 [LVT-10516, LVT-10517]; **48** – Hot Creek Valley, 38.2100, -116.1800 [LVT-9493 (NMMNH-5613), LVT-9497]; **49** – Railroad Valley, 38.5115, -115.7829 [LVT-8649, LVT-8650]; 38.5209, -115.7533 [LVT-8651]; **50** – 2-8 mi SE Carrant, 38.6980, -115.4861 [LVT-9482, LVT-9483]; 38.7203, -115.4624 [LVT-9478, LVT-9479, LVT-94780, LVT-9484];



38.7275, -115.4592 [LVT-9474, LVT-9475, LVT-9476]; 38.7934, -115.5254 [LVT-9469, LVT-9470, LVT-9471]; Washoe County: **51** – Mullen Pass, 39.8730, -119.6210 [LVT-10215, LVT-10216, LVT-10217, LVT-10218, LVT-10219, LVT-10220, LVT-10221, LVT-10222, LVT-10223, LVT-10224, LVT-10225, LVT-10226, LVT-10227]; **52** – Flanigan, 40.1891, -119.8490 [LVT-8525, LVT-8526, LVT-8527]; **53** – 5 mi NE Gerlach, 40.7469, -119.2875 [LVT-9599]; White Pine: **54** – Snake Valley, 38.6303, -114.2701 [LVT-8166 (NMMNH-5587)]; 38.6967, -114.1146 [LVT-8173 (NMMNH-5585); LVT-8175, LVT-8176, LVT-8178]; Spring Valley, 38.6586, -114.3823 [LVT-7846 (NMMNH-5547), LVT-7899]; **55** – Spring Valley, 38.9378, -114.4176 [LVT-7850]; 38.9406, -114.4212 [LVT-7901]; **56** – Spring Valley, 39.1397, -114.4982 [LVT-7902, LVT-8054, LVT-8055]; 39.1975, -114.4983 [LVT-8053 (NMMNH-5563)]; 39.2885, -114.4635 [LVT-7903 (NMMNH 5553), LVT 08057]; 39.3026, -114.3890 [LVT-7906 (NMMNH-5556)]; **57** – Steptoe Valley, S of Ely, 39.2336, -114.8395 [LVT-9804]; **58** – Spring Valley, 39.4867, -114.3878 [LVT-8056]; OR: Harney County: **59** – Fields, 42.3067, -118.6598 [LVT-8958, LVT-8959, LVT-8960, LVT-8965]; **60** – 20 mi S Narrows, 43.0709, -118.8730 [LVT-8943, LVT-8944, LVT-8947, LVT-8948]; Lake County: **61** – Alkali Lake, 42.9433, -120.0274 [LVT-8936, LVT-8937]; 43.0107, -120.0114 [LVT-8925, LVT-8927, LVT-8928, LVT-8929, LVT-8930, LVT-8938, LVT-8940, LVT-8941]; Utah: Boxelder County: **62** – Kelton, 41.7260 , -113.1380 [LVT-8625, LVT-8628]; 41.7410, -113.1260 [LVT-8629, LVT-8630, LVT-8631], 41.7550, -113.1080 [LVT-8632, LVT-8633, LVT-8634, LVT-8635]; Juab County: **63** – Snake Valley, 39.4568, -113.9606 [LVT-8656, LVT-8657, LVT-8658, LVT-8659, LVT-8665, LVT-8666, LVT-8667, LVT-8670, LVT-8671, LVT-8672, LVT-8673, LVT-8675, LVT-8679]; Millard County: **64** – 10 mi S Delta,

39.0580, -112.7640 [LVT-8594]; White Sage Valley, 39.1030, -112.8500 [LVT-8587, LVT-8588, LVT-8603 (NMMNH-5597); Cricket Mountains, 39.1120, -112.7600 (LVT-8589, LVT-8590, LVT-8591, LVT-8592, LVT-8593); Tooele County: **65** – Rush Valley, 40.0780, -112.2560 [LVT-8621]; **66** – Puddle Valley, 40.8910, -112.9330 [LVT-8639]; 40.8950, -112.9460 [LVT-8638]; 40.9100, -112.9040 [LVT-8646, LVT-8647]; Washington County: **67** – 23 mi N Littlefield, 37.0784, -113.9237 [LVT-10409, LVT-10410, LVT-10411, LVT-10412, LVT-10413]; 24 mi N Littlefield, 37.0952, -113.9540 [LVT-10418, LVT-10419, LVT-10420].

## APPENDIX 2

### SAMPLES OF *PHRYNOSOMA PLATYRHINOS*

Descriptions of general sample areas for *Phrynosoma platyrhinos* by country, state, county, site identification number (referenced in Fig. 2.1A), site reference name, latitude, and longitude. Samples associated with the sample area follow in parentheses.

United States: Arizona: Coconino County: **1** – House Rock Valley, 36.6294, -111.9747 (LVT-9367); La Paz County: **2** – Yuma Proving Grounds, 32.8683, -114.3742 (CAS 228862-3, LVT-9951-3); **3** – vicinity of Salome, 33.6379, -113.4419 (LVT-7381); **4** – Bouse-Quarzsie Rd (CAS-228896); **5** – Bouse Dunes, 33.8578, -113.9480 (CAS-228895, LVT-829-32); **6** – Wenden, 33.8819, -113.5475 (LVT-7382-3); Maricopa County: **7** – vicinity of Ajo, 32.4902, -112.8621 (LVT-810, 817-9); **8** – Wittmann, 33.7764, -112.5285 (LVT-361); Mohave County: **9** – vicinity of Golden Shores, 34.8038, -114.4966 (LVT-7728); **10** – vicinity of Oatman, 34.9090, -114.4327 (LVT-6346); **11** – E of Kingman, 35.1059, -113.6698 (LVT-6343-5); **12** – Dolan Spring Rd, 35.6407, -114.2424 (LVT-735); Pima County: **13** – W of Tucson Mtn Park, 32.2231, -111.1143 (LVT-820); Yuma County: **14** – Pinacate Lava Flow, 32.1017, -113.4621 (CAS-228867-9); **15** – vicinity of Copper Mtn, BMG Airforce Range, 32.4375, -113.9663 (CAS-8874); **16** – BMG Airforce Range, 32.4873, -114.4577 (CAS\_228841, 9922); **17** – Mohawk Dunes, 32.6947, -113.8083 (CAS-228865); California: Imperial County: **18** – Ocotillo, 32.7764, -116.0695 (CAS-3601, LVT-6368); **19** – Ogilby Rd, 32.9833, -114.8989 (CAS-228893); Inyo County: **20** – vicinity of

Shoshone, 35.9956, -116.2149 (LVT-6080); **21** – Panamint Valley Rd, 36.2804, -117.3712 (LVT-8379-83); **22** – vicinity of Darwin Falls, 36.3567, -117.5279 (LVT-8374-8); **23** – Saline Valley Rd, 36.4166, -117.6554 (LVT-7386-7); **24** – vicinity of Bishop, 37.2359, -118.2565 (LVT-851-4); **25** – Eureka Valley Rd, 37.2589, -117.9279 (LVT-7385); **103** – vicinity of Benton, 37.6878, -118.4309 (LVT-10514); Modoc County: **26** – Surprise Valley, 41.5646, -120.0235 (LVT-6364-5); Riverside County: **27** – vicinity of Blythe, 33.5388, -114.7849 (LVT-816); **28** – near Joshua Tree NM, 33.8680, -115.6382 (LVT-815); San Bernardino County: **29** – Cadiz Valley, 34.0550, -115.2503 (LVT-7389); **30** – Whipple Mtns, 34.2587, -114.5429 (LVT-7390); **31** – vicinity of Amboy, 34.5377, -115.7236 (LVT-812-4, LVT-6082-3); **32** – Ord Mtns, 34.7113, -116.8321 (LVT-7388); **33** – vicinity of Barstow, 34.8612, -117.1005 (LVT-8366-8); **34** – vicinity of Kelso Dunes, 34.9108, -115.7303 (LVT-8369-73); **35** – Arrowhead Junction, 35.1851, -114.9027 (LVT-7735); **36** – Cima Rd, 35.4108, -115.6418 (LVT-636); **37** – Spring Mtn, 35.5855, -115.6357 (LVT-9807); **38** – Trona, 35.7628, -117.3719 (LVT-8365); San Diego County: **39** – Carrizo Badlands, 32.8548, -116.1775 (DGM-804); **40** – Blair Valley, 33.0293, -116.4146 (DGM-549); **41** – Ocotillo Wells State Vehicle Recreation Area, 33.1544, -116.1675 (CAS-228891); **42** – Borrego Springs, 33.1834, -116.3257 (CAS-228880, 228894); Idaho: Owyhee County: **43** – Bruneau Canyon, 42.7629, -115.7564 (LVT-10373-4); **44** – vicinity of Murphy, 43.1471, -116.4998 (LVT-934); Nevada: Churchill County: **45** – Hot Springs Mtns, 39.7581, -118.8735 (LVT-10194-6); **46** – Stillwater Mtns, 39.9935, -118.1820 (LVT-867); Clark County: **47** – El Dorado Valley, 35.7727, -114.9104 (LVT-0473-5); **48** – S of Sloan, 35.9122, -115.2064 (LVT-801-2); **49** – vicinity of Las Vegas, 36.1256, -115.2612 (LVT-

811); **50** – vicinity of Apex, 36.2977, -114.9625 (LVT-824-6); **51** – Lee Canyon, 36.4085, -115.5668 (CAS-223435); **52** – Whitney Pockets, 36.4988, -114.1519 (CAS-223372); **53** – Overton, 36.5433, -114.4649 (LVT-465); **54** – Arrowhead, 36.6589, -114.5758, (LVT-803-5); **55** – Upper Mormon Mesa, 36.7774, -114.4281 (LVT-0471-2); **56** – Coyote Springs Valley, 36.8312, -114.8646 (LVT-476-8); Elko County: **57** – Currie, 40.2636, -114.7475 (LVT-924); **58** – Elko, 40.8884, -115.8083 (LVT-10382-3); Esmeralda County: **59** – vicinity of Bonnie Claire Lake, 37.1539, -117.1781 (CAS-223438, 8976, LVT-6361); **60** – vicinity of Gold Field, 37.8735, -117.2436 (LVT-6311-5); **61** – vicinity of Tonopah, 37.9120, -117.2235 (LVT-10168-72); **62** – Blow Sand Mtn, 39.1990, -118.7221 (LVT-10464); Humboldt County: **63** – vicinity of Sulphur Landing, 40.8432, -118.7577 (LVT-9625); **64** – Black Rock Desert, 40.8953, -118.5200 (LVT-919); **65** – Golconda Summit, 40.9210, -117.3915 (DGM-1775); **66** – vicinity of Winnemucca, 41.3463, -117.5984 (LVT-941-2); **67** – McGee Mtns, 41.7668, -118.9189 (LVT-10248-53); Lander County: **68** – vicinity of Battle Mtn, 40.1889, -117.1591 (LVT-916-8); Lincoln County: **69** – Tikaboo Valley, 37.3774, -115.4663 (LVT-9451); **70** – Rachel, 37.6715, -115.7578, (LVT-9461); **71** – Dry Lake Valley, 37.9054, -114.8152 (LVT-7760, 9958-9); **72** – vicinity of Hiko, 37.9230, -115.0083 (LVT-0915); Nye County: **73** – Cave Valley, 38.3673, -114.8243 (LVT-9957); **74** – Pahrump Valley, 36.1626, -115.8990 (LVT-6079); **75** – Ash Meadows, 36.4275, -116.3520 (LVT-10246-7); **76** – vicinity of Mercury Hwy and Hwy 95, 36.5971, -115.9536 (LVT-6317); **77** – Crater Flat, 36.8085, -116.6047 (LVT-470); **78** – vicinity of Beatty, 37.0970, -116.7958 (LVT-6316); **79** – vicinity of Hwy 376 and Hwy 6, 38.3896, -117.1595 (LVT-838-40, 843-5); **80** – Round Mtn dump site, 38.7049, -117.0742 (LVT-841-2); **81** – Railroad

Valley near Currant, 38.7243, -115.4783 (LVT-9464-6, 9481); Pershing County: **82** – Black Rock Desert, 40.7978, -119.0297 (LVT-920-2); Washoe County: **83** – Empire Range Rd, 40.6256, -119.4316 (LVT-6366-7); White Pine County: **84** – Hamlin Valley, 38.6599, -114.1502 (DGM-1104); **85** – Snake Valley, 38.9737, -114.0745 (CAS-223386, 3390, LVT-6362-3); **86** – Spring Valley, 39.1462, -114.5084 (LVT-9805); **104** – White River Valley, 38.99379, -115.07311 (LVT-10523); Oregon: Harney County: **87** – vicinity of Fields; 42.2527, -118.6606 (LVT-0936-40); Malheur County: **88** – Jordan Range Rd, 42.2149, -117.7841 (LVT-935); Utah: Box Elder County: **89** – West Desert, 41.4490, -113.6427 (DGM-1027-9); **90** – vicinity of Etna, 41.5426, -113.9629 (LVT-925-6); Iron County: **91** – Lund, 38.0245, -113.4044 (LVT-858-61); Millard County: **92** – Desert Experimental Range, 38.6623, -113.8493 (LVT-931-3); **93** – Sunstone Knoll, 39.1487, -112.7164 (CAS-228887); **94** – Tule Valley, 39.4333, -113.6333 (LVT-10462-3); Tooele County: **95** – vicinity of Callao, 40.1778, -113.8031 (LVT-930); **96** – Skull Valley Rd, 40.6641, -112.6739 (CAS-228889); **97** – vicinity of Wendover, 40.9936, -113.8401 (LVT-929); **98** – N of Wendover, 41.2356, -114.0244 (LVT-927-8); Washington County: **99** – Beaver Dam Slope, 37.0846, -113.9483 (CAS-228888, LVT-855-7); Mexico: Baja California North: **100** – El Moreno, 31.0267, -115.1010 (DGM-481); **101** – Valle Santa Clara, 31.2051, -115.3121, (DGM-477-8); Sonora: **102** – Puerto Penasco, 31.5536, -113.4735 (KVY-0013).

## APPENDIX 3

### GLOSSARY

**Analogous climate:** An association of climatic variables (either in combination or by correlation) used in model calibration that is the same as that used in the model projected either in time or across space.

**Calibration:** Development of species distribution models using a particular algorithm (e.g., General Linear Models, Bioclim, GARP, Maxent), records of species presence or presence-absence, and a set of environmental variables; establishing a baseline for model projection either in time or across space.

**Calibration range:** The range of values for a particular variable across a particular geographic extent used during the calibration of a species distribution model.

**Clamping:** The restriction of variable values during model projection to those encountered during calibration. In Maxent this is done by truncating the value to the minimum or maximum of the calibration range. The logistic probability (i.e. habitat suitability) of the truncated values in the projected model equals that of the minimum or maximum value used in calibrating the model.

**Commission error:** A geographic location identified by a model as suitable when no individuals of the focal species actually occur in the predicted area – over-prediction.

**Extrapolation:** Estimating new values beyond a discrete set of known values based on a hypothesized relationship. In species distribution modeling, this means estimating the response of a species to environmental values outside the range of known values.

**Fundamental Niche:** The full suite of environmental conditions under which populations of a particular species can potentially survive (sensu Hutchinson 1957).

Organisms typically occupy a narrower range of conditions (i.e. realized niche) because of biotic interactions, barriers to dispersal, or non-existence of particular abiotic conditions during a certain time period.

**Non-analogous climate:** An association of climatic variables (either in combination or by correlation) used in model calibration that are NOT the same as those in the projected model (either in time or across space).

**Omission error:** A geographic location identified by a model as unsuitable when individuals of the focal species actually occur at the site – under-prediction.

**Projection:** Application of rules developed during the calibration of a species distribution model to a new set of environmental layers representing either a different time period or different area.

**Projection range:** Range of values for a particular variable across the geographic extent of a location on which a species distribution model will be projected.

**Realized Niche:** The suite of biotic and abiotic environmental conditions defining the actual occurrence of a species. Populations may be able to survive under a wider range of conditions (i.e. fundamental niche), but are limited to the realized niche because of factors such as biotic interactions, barriers to dispersal, or non-existence of particular abiotic conditions during certain time period.

**Realized niche shift:** A shift in the actual or potential occurrence of a species resulting from plasticity, changes in biotic interactions or the development of previously



unavailable (non-existent) environmental conditions. Under such conditions, there is no change in the fundamental niche, only a shift in the portion of the fundamental niche occupied.

VITA

Graduate College  
University of Nevada, Las Vegas

Tereza Jezkova

Degrees:

Master of Science, Biology, 2002  
Charles University, Prague, Czech Republic

Special Honors and Awards:

2004. NSF EPSCoR ACES Grant, \$6,000  
2005. Research Grant, Graduate Student Association, UNLV, \$500  
2005. NSF EPSCoR ACES Grant, \$20,000  
2005. Grant-In-Aid, American Society of Mammalogists, \$1,200  
2005. Research Grant, Graduate Student Association, UNLV, \$500  
2006. Research Grant, Graduate Student Association, UNLV, \$750  
2008. Doctoral Dissertation Improvement Grant, National Science Foundation, \$11,500  
2009. Travel Award, International Biogeography Society Meeting, Mérida, México, \$1,300  
2009. Best Poster Award, International Biogeography Society Meeting, Mérida, México  
2009. BIOS publication award, UNLV  
2009. Travel Grant, Graduate Student Association, UNLV, \$300  
2010. Best oral presentation award, Graduate Student Association, UNLV, \$100

Publications:

Rodríguez-Robles JA, **Jezkova T**, Leal M. 2010. Climatic stability and genetic divergence in the tropical insular lizard *Anolis krugi*, the Puerto Rican "Lagartijo Jardinero de la Montaña." *Molecular Ecology* 19(9):1860-1876.

**Jezkova T**, Jaeger JR, Marshall ZL, Riddle BR. 2009. Pleistocene impacts on the phylogeography of the desert pocket mouse (*Chaetodipus penicillatus*). *Journal of Mammalogy* 90:306–320.

**Jezkova T**, Leal M, Rodríguez-Robles JA. 2009. Living together but remaining apart: comparative phylogeography of *Anolis poncensis* and *A. cooki*, two lizards endemic to the aridlands of Puerto Rico. *Biological Journal of the Linnaean Society* 96: 617–634.

Glaudas X, **Jezkova T**, Rodríguez-Robles JA. 2008. Feeding ecology of the Great Basin Rattlesnake (*Crotalus lutosus*, Viperidae). *Canadian Journal of Zoology* 86:723-734.

Rodríguez-Robles JA, **Jezkova T**, Leal M. 2008. Genetic structuring in the threatened "Lagartijo del Bosque Seco" (*Anolis cooki*) from Puerto Rico. *Molecular Phylogenetics and Evolution* 46:503-514.

Rodríguez-Robles JA, **Jezkova T**, García MA. 2007. Evolutionary relationships and historical biogeography of *Anolis deseichensis* and *A. monensis*, two lizards endemic to small islands in the eastern Caribbean Sea. *Journal of Biogeography* 34:1546-1558.

Dissertation Title: Reconstructing species responses to past climatic changes using niche modeling and genetic data

Dissertation Examination Committee:

Committee Chairperson, Brett Riddle, Ph. D.

Committee Member, John Klicka, Ph. D.

Committee Member, Daniel Thompson, Ph. D.

Committee Member, John Hafner, Ph. D.

Graduate Faculty Representative, Matthew Lachniet, Ph. D.